

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number  
**WO 03/082190 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K** (74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (21) International Application Number: **PCT/US03/08722**
- (22) International Filing Date: 21 March 2003 (21.03.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/367,655 26 March 2002 (26.03.2002) US
- (71) Applicant (for all designated States except US): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HAGMANN, William, K.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **LIN, Linus, S.** [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **SHAH, Shrenik, K.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **GOULET, Mark, T.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **JEWELL, James, P.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **SPIROCYCLIC AMIDES AS CANNABINOID RECEPTOR MODULATORS**

(57) Abstract: Novel compounds of structural formula (I) are antagonists and/or inverse agonists of the Cannabinoid-1 (CB1) receptor and are useful in the treatment, prevention and suppression of diseases mediated by the CB1 receptor. The compounds of the present invention are useful as psychotropic drugs in the treatment of psychosis, memory deficits, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders including multiple sclerosis and Guillain-Barre syndrome and the inflammatory sequelae of viral encephalitis, cerebral vascular accidents, and head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, movement disorders, and schizophrenia. The compounds are also useful for the treatment of substance abuse disorders, the treatment of obesity or eating disorders, as well as, the treatment of asthma, constipation, chronic intestinal pseudo-obstruction, and cirrhosis of the liver.

WO 03/082190 A2

## TITLE OF THE INVENTION

## SPIROCYCLIC AMIDES AS CANNABINOID RECEPTOR MODULATORS

## CROSS REFERENCE TO RELATED APPLICATIONS

5 Not applicable.

## BACKGROUND OF THE INVENTION

Marijuana (*Cannabis sativa L.*) and its derivatives have been used for centuries for medicinal and recreational purposes. A major active ingredient in marijuana and hashish has been determined to be  $\Delta^9$ -tetrahydrocannabinol ( $\Delta^9$ -THC). Detailed research has revealed that the biological action of  $\Delta^9$ -THC and other members of the cannabinoid family occurs through two G-protein coupled receptors termed CB1 and CB2. The CB1 receptor is primarily found in the central and peripheral nervous systems and to a lesser extent in several peripheral organs. The CB2 receptor is found primarily in lymphoid tissues and cells. Three endogenous ligands for the cannabinoid receptors derived from arachidonic acid have been identified (anandamide, 2-arachidonoyl glycerol, and 2-arachidonoyl glycerol ether). Each is an agonist with activities similar to  $\Delta^9$ -THC, including sedation, hypothermia, intestinal immobility, antinociception, analgesia, catalepsy, anti-emesis, and appetite stimulation.

The genes for the respective cannabinoid receptors have each been disrupted in mice. The CB1<sup>-/-</sup> receptor knockout mice appeared normal and fertile. They were resistant to the effects of  $\Delta^9$ -THC and demonstrated a strong reduction in the reinforcing properties of morphine and the severity of withdrawal syndrome. They also demonstrated reduced motor activity and hypoalgesia. The CB2<sup>-/-</sup> receptor knockout mice were also healthy and fertile. They were not resistant to the central nervous system mediated effects of administered  $\Delta^9$ -THC. There were some effects on immune cell activation, reinforcing the role for the CB2 receptor in immune system functions.

Excessive exposure to  $\Delta^9$ -THC can lead to overeating, psychosis, hypothermia, memory loss, and sedation. Specific synthetic ligands for the cannabinoid receptors have been developed and have aided in the characterization of the cannabinoid receptors: CP55,940 (J. Pharmacol. Exp. Ther. 1988, 247, 1046-1051); WIN55212-2 (J. Pharmacol. Exp. Ther. 1993, 264, 1352-1363); SR141716A (FEBS Lett. 1994, 350, 240-244; Life Sci. 1995, 56, 1941-1947); and SR144528 (J.

Pharmacol. Exp. Ther. 1999, 288, 582-589). The pharmacology and therapeutic potential for cannabinoid receptor ligands has been reviewed (Exp. Opin. Ther. Patents 1998, 8, 301-313; Ann. Rep. Med. Chem., A. Doherty, Ed.; Academic Press, NY 1999, Vol. 34, 199-208; Exp. Opin. Ther. Patents 2000, 10, 1529-1538; Trends in  
5 Pharma. Sci. 2000, 21, 218-224). There is at least one CB1 modulator characterized as an inverse agonist or an antagonist, N-(1-piperidiny)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methylpyrazole-3-carboxamide (SR141716A), in clinical trials for treatment of eating disorders at this time. There still remains a need for potent low molecular weight CB1 modulators that have pharmacokinetic and pharmacodynamic  
10 properties suitable for use as human pharmaceuticals.

Treatment of asthma with CB1 receptor modulators (such as CB1 inverse agonists) is supported by the finding that presynaptic cannabinoid CB1 receptors mediate the inhibition of noradrenaline release (in the guinea pig lung) (Europ. J. of Pharmacology, 2001, 431 (2), 237-244).

15 Treatment of cirrhosis of the liver with CB1 receptor modulators is supported by the finding that a CB1 receptor modulator will reverse the low blood pressure observed in rats with carbon tetrachloride-induced liver cirrhosis and will lower the elevated mesenteric blood flow and portal vein pressure (Nature Medicine, 2001, 7 (7), 827-832).

20 US Patents US 5,624,941 and US 6,028,084, PCT Application Nos. WO98/43636, WO98/43635, and WO 02/076945, and EPO Application No. EP-658546 disclose substituted pyrazoles having activity against the cannabinoid receptors.

PCT Application Nos. WO98/31227 and WO98/41519 also disclose  
25 substituted pyrazoles having activity against the cannabinoid receptors.

PCT Application Nos. WO98/37061, WO00/10967, and WO00/10968 disclose diaryl ether sulfonamides having activity against the cannabinoid receptors.


PCT Application Nos. WO97/29079 and WO99/02499 disclose  
30 alkoxy-isoindolones and alkoxy-quinolones as having activity against the cannabinoid receptors.

US Patent US 5,532,237 discloses N-benzoyl-indole derivatives having activity against the cannabinoid receptors.

US Patents US 4,973,587, US 5,013,837, US 5,081,122, and US 5,112,820, US 5,292,736 disclose aminoalkylindole derivatives as having activity against the cannabinoid receptors.


PCT publication WO 01/58869 discloses pyrazoles, pyrroles and imidazole cannabinoid receptor modulators useful for treating respiratory and non-respiratory leukocyte activation-associated disorders.

PCT publication WO 01/09120, assigned to Ortho-McNeil Pharmaceutical, Inc., is directed to monosubstituted cycloalkyl amides of the structural formula:



wherein B<sub>1</sub>, B<sub>2</sub> and R<sub>1</sub> are hydrogen, R<sub>2</sub> is aryl or heteroaryl, m is 1 and L is C<sub>3</sub>-7cycloalkyl; and their use as neuropeptide Y<sub>5</sub> receptor ligands for the treatment of obesity and other disorders associated with NPY receptor subtype Y<sub>5</sub>.


PCT publication WO 96/33159, assigned to Abbot Laboratories, is directed to tetra-substituted cyclobutyl amides of structural formula:



wherein R<sub>6</sub> is hydrogen or lower alkyl, R<sub>6a</sub> and R<sub>6b</sub> are hydrogen, Y is a single bond, R<sub>3</sub> and R<sub>4</sub> are phenyl; and their use as squalene synthase and protein farnesyltransferase inhibitors for the inhibition of cholesterol biosynthesis.

PCT publication WO 98/33765, assigned to E. I. Du Pont De Nemours and Company, is directed to bis(2,2-dichlorocyclopropyl) amides of the structural formula:







wherein  $R^1$  is H, methyl or ethyl,  $R^2$  is phenyl, X is  $-CH_2-$  and Z is phenyl; and their use as fungicides for controlling plant diseases caused by fungal plant pathogen

Lack et al, J. Pharmacol. Exptl. Therap, Vol. 139, p 248-58, 1963

discloses the following two compounds:



and




10

The compounds of the present invention are modulators of the Cannabinoid-1 (CB1) receptor and are useful in the treatment, prevention and suppression of diseases mediated by the Cannabinoid-1 (CB1) receptor. In particular, compounds of the present invention are antagonists or inverse agonists of the CB1

receptor. The invention is concerned with the use of these compounds to modulate the Cannabinoid-1 (CB1) receptor. As such, compounds of the present invention are useful as psychotropic drugs in the treatment of psychosis, memory deficits, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders including multiple sclerosis and Guillain-Barre syndrome and the inflammatory sequelae of viral encephalitis, cerebral vascular accidents, and head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, movement disorders, and schizophrenia. The compounds are also useful for the treatment of substance abuse disorders, particularly to opiates, alcohol, marijuana, and nicotine. The compounds are also useful for the treatment of eating disorders by inhibiting excessive food intake and the resulting obesity and complications associated therewith. The compounds are also useful for the treatment of constipation and chronic intestinal pseudo-obstruction, as well as for the treatment of asthma, and cirrhosis of the liver.

## SUMMARY OF THE INVENTION

The present invention is concerned with substituted amides of the general Formula I:



(I)

and pharmaceutically acceptable salts thereof which are antagonists and/or inverse agonists of the Cannabinoid-1 (CB1) receptor and are useful in the treatment, prevention and suppression of diseases mediated by the Cannabinoid-1 (CB1) receptor. The invention is concerned with the use of these novel compounds to selectively antagonize the Cannabinoid-1 (CB1) receptor. As such, compounds of the present invention are useful as psychotropic drugs in the treatment of psychosis, memory deficits, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders including multiple sclerosis and Guillain-Barre syndrome and the inflammatory sequelae of viral encephalitis, cerebral vascular accidents, and head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, movement disorders,

and schizophrenia. The compounds are also useful for the treatment of substance abuse disorders, particularly to opiates, alcohol, marijuana, and nicotine, including smoking cessation. The compounds are also useful for the treatment of obesity or eating disorders associated with excessive food intake and complications associated therewith. The compounds are also useful for the treatment of constipation and chronic intestinal pseudo-obstruction. The compounds are also useful for the treatment of cirrhosis of the liver. The compounds are also useful for the treatment of asthma.

The present invention is also concerned with treatment of these conditions, and the use of compounds of the present invention for manufacture of a medicament useful in treating these conditions. The present invention is also concerned with treatment of these conditions through a combination of compounds of formula I and other currently available pharmaceuticals.


The invention is also concerned with novel compounds of structural formula I.

The invention is also concerned with pharmaceutical formulations comprising one of the compounds as an active ingredient.

The invention is further concerned with processes for preparing the compounds of this invention.

## DETAILED DESCRIPTION OF THE INVENTION

The compounds used in the methods of the present invention are represented by structural formula I:



(I)

or a pharmaceutically acceptable salt thereof, wherein;

R<sup>1</sup> is selected from:

- (1) C<sub>1-10</sub>alkyl,
- (2) C<sub>3-10</sub>cycloalkyl,

(3) cycloheteroalkyl,

(4) aryl, and

(5) heteroaryl,

5 wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from  $R^a$ , and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from  $R^b$ ;

$R^2$  is selected from:

- 10 (1) C<sub>3-10</sub>cycloalkyl,  
(2) cycloheteroalkyl,  
(3) aryl,  
(4) heteroaryl,  
(5) -OR<sup>d</sup>,  
(6) -NR<sup>c</sup>R<sup>d</sup>, and  
15 (7) -CO<sub>2</sub>R<sup>d</sup>,

wherein each alkyl is optionally substituted with one, two, three or four substituents independently selected from  $R^a$ , and each cycloalkyl, and cycloheteroalkyl aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from  $R^b$ ;

20  $R^3$  is selected from:

- (1) hydrogen, and  
(2) C<sub>1-4</sub>alkyl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from  $R^a$ ;

25  $R^4$  is selected from:

- (1) hydrogen,  
(2) C<sub>1-10</sub>alkyl,  
(3) C<sub>2-10</sub>alkenyl,  
(4) C<sub>2-10</sub>alkynyl,  
30 (5) -OR<sup>c</sup>,  
(6) -CO<sub>2</sub>R<sup>c</sup>,  
(7) -OCOOR<sup>c</sup>,  
(8) -OCONR<sup>c</sup>R<sup>d</sup>,  
(9) -NR<sup>c</sup>R<sup>d</sup>,  
35 (10) -NR<sup>c</sup>(CO)OR<sup>d</sup>,

(11)  $-\text{NR}^{\text{c}}\text{SO}_2\text{R}^{\text{d}}$ ,

(12)  $-\text{S}(\text{O})_m\text{R}^{\text{c}}$ ,

(13) aryl, and

(14) heteroaryl,

5 wherein alkyl, alkenyl, and alkynyl are optionally substituted with one, two, three or four substituents independently selected from  $\text{R}^{\text{a}}$ , and aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from  $\text{R}^{\text{b}}$ ;

$\text{R}^{\text{b}}$  is selected from:

10 (1) hydrogen,

(2)  $\text{C}_{1-4}$ alkyl,

(3)  $\text{C}_{2-4}$ alkenyl,

(4)  $\text{C}_{2-4}$ alkynyl,

(5)  $-\text{OR}^{\text{d}}$ ,

15 (6) halogen,

(7)  $-\text{CN}$ ,

(8)  $-\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $\text{R}^{\text{a}}$ ;

20 **A** is a 3- to 8-membered monocyclic saturated ring incorporating the same carbon atom to which  $\text{R}^{\text{4}}$  is attached, and optionally containing one to two heteroatoms chosen from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted with one, two, three or four substituents independently selected from oxo and  $\text{R}^{\text{b}}$ ;  
each  $\text{R}^{\text{a}}$  is independently selected from:

25 (1)  $-\text{OR}^{\text{c}}$ ,

(2)  $-\text{NR}^{\text{c}}\text{S}(\text{O})_m\text{R}^{\text{d}}$ ,

(3)  $-\text{NO}_2$ ,

(4) halogen,

(5)  $-\text{S}(\text{O})_m\text{R}^{\text{c}}$ ,

30 (6)  $-\text{SR}^{\text{c}}$ ,

(7)  $-\text{S}(\text{O})_2\text{OR}^{\text{c}}$ ,

(8)  $-\text{S}(\text{O})_m\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,

(9)  $-\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,

(10)  $-\text{O}(\text{CR}^{\text{e}}\text{R}^{\text{f}})_n\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,

- (11)  $-\text{C}(\text{O})\text{R}^{\text{c}}$ ,  
 (12)  $-\text{CO}_2\text{R}^{\text{i}}$ ,  
 (13)  $-\text{CO}_2(\text{CR}^{\text{e}}\text{R}^{\text{f}})_n\text{CONR}^{\text{c}}\text{R}^{\text{d}}$ ,  
 (14)  $-\text{OC}(\text{O})\text{R}^{\text{c}}$ ,  
 5 (15)  $-\text{CN}$ ,  
 (16)  $-\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,  
 (17)  $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{R}^{\text{d}}$ ,  
 (18)  $-\text{OC}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,  
 (19)  $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{OR}^{\text{d}}$ ,  
 10 (20)  $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,  
 (21)  $-\text{CR}^{\text{c}}(\text{N}-\text{OR}^{\text{d}})$ ,  
 (22)  $\text{CF}_3$ ,  
 (23)  $-\text{OCF}_3$ ,  
 (24)  $\text{C}_3\text{-8cycloalkyl}$ ,  
 15 (25) cycloheteroalkyl, and  
 (26) oxo;

each  $\text{R}^{\text{b}}$  is independently selected from:

- (1)  $\text{R}^{\text{a}}$ ,  
 (2)  $\text{C}_1\text{-10alkyl}$ ,  
 20 (3)  $\text{C}_3\text{-8cycloalkyl}$ ,  
 (4) cycloheteroalkyl,  
 (5) aryl,  
 (6)  $\text{arylC}_1\text{-4alkyl}$ ,  
 (7) heteroaryl, and  
 25 (8) heteroaryl $\text{C}_1\text{-4alkyl}$ ;

each  $\text{R}^{\text{c}}$  and  $\text{R}^{\text{d}}$  is independently selected from:

- (1) hydrogen,  
 (2)  $\text{C}_1\text{-10alkyl}$ ,  
 (3)  $\text{C}_2\text{-10alkenyl}$ ,  
 30 (4)  $\text{C}_2\text{-10alkynyl}$ ,  
 (5) cycloalkyl,  
 (6) cycloalkyl- $\text{C}_1\text{-10alkyl}$ ,  
 (7) cycloheteroalkyl,  
 (8) cycloheteroalkyl- $\text{C}_1\text{-10alkyl}$ ,

- (9) aryl,
- (10) heteroaryl,
- (11) aryl-C<sub>1-10</sub>alkyl, and
- (12) heteroaryl-C<sub>1-10</sub>alkyl, or

5 R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sub>g</sub>,

each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two, three or four substituents selected from R<sup>h</sup>;

10 each R<sup>e</sup> and R<sup>f</sup> are independently selected from:

- (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>2-10</sub> alkenyl,
- (4) C<sub>2-10</sub>alkynyl;
- 15 (5) cycloalkyl,
- (6) cycloalkyl-C<sub>1-10</sub> alkyl,
- (7) cycloheteroalkyl,
- (8) cycloheteroalkyl-C<sub>1-10</sub> alkyl,
- (9) aryl,
- 20 (10) heteroaryl,
- (11) aryl-C<sub>1-10</sub> alkyl, and
- (12) heteroaryl-C<sub>1-10</sub> alkyl, or

R<sup>e</sup> and R<sup>f</sup> together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and  
25 nitrogen;

each R<sub>g</sub> is independently selected from:

- (1) C<sub>1-10</sub>alkyl,
- (2) C<sub>3-8</sub>cycloalkyl,
- (3) cycloheteroalkyl,
- 30 (4) aryl,
- (5) arylC<sub>1-4</sub>alkyl,
- (6) heteroaryl,
- (7) heteroarylC<sub>1-4</sub>alkyl,
- (8) -S(O)<sub>m</sub>R<sup>e</sup>,
- 35 (9) -C(O)R<sup>e</sup>,

- (10)  $-\text{CO}_2\text{Re}$ ,
- (11)  $-\text{CO}_2(\text{CReRf})_n\text{CONReRf}$ , and
- (12)  $-\text{C}(\text{O})\text{NReRf}$ ;

each  $\text{R}^h$  is independently selected from:

- 5 (1) halogen,
- (2)  $\text{C}_{1-10}$ alkyl,
- (3)  $\text{C}_{3-8}$ cycloalkyl,
- (4) cycloheteroalkyl,
- (5) aryl,
- 10 (6)  $\text{arylC}_{1-4}$ alkyl,
- (7) heteroaryl,
- (8) heteroaryl $\text{C}_{1-4}$ alkyl,
- (9)  $-\text{ORE}$ ,
- (10)  $-\text{NReS}(\text{O})_m\text{Rf}$ ,
- 15 (11)  $-\text{S}(\text{O})_m\text{Re}$ ,
- (12)  $-\text{SRe}$ ,
- (13)  $-\text{S}(\text{O})_2\text{ORE}$ ,
- (14)  $-\text{S}(\text{O})_m\text{NReRf}$ ,
- (15)  $-\text{NReRf}$ ,
- 20 (16)  $-\text{O}(\text{CReRf})_n\text{NReRf}$ ,
- (17)  $-\text{C}(\text{O})\text{Re}$ ,
- (18)  $-\text{CO}_2\text{Re}$ ,
- (19)  $-\text{CO}_2(\text{CReRf})_n\text{CONReRf}$ ,
- (20)  $-\text{OC}(\text{O})\text{Re}$ ,
- 25 (21)  $-\text{CN}$ ,
- (22)  $-\text{C}(\text{O})\text{NReRf}$ ,
- (23)  $-\text{NReC}(\text{O})\text{Rf}$ ,
- (24)  $-\text{OC}(\text{O})\text{NReRf}$ ,
- (25)  $-\text{NReC}(\text{O})\text{ORf}$ ,
- 30 (26)  $-\text{NReC}(\text{O})\text{NReRf}$ ,
- (27)  $-\text{CF}_3$ , and
- (28)  $-\text{OCF}_3$ ;

each  $\text{R}^i$  is independently selected from:

- (1)  $\text{C}_{1-10}$ alkyl,



- (2) C<sub>2-10</sub> alkenyl,  
(3) C<sub>2-10</sub> alkynyl,  
(4) cycloalkyl,  
(5) cycloalkyl-C<sub>1-10</sub> alkyl,  
5 (6) cycloheteroalkyl,  
(7) cycloheteroalkyl-C<sub>1-10</sub> alkyl,  
(8) aryl,  
(9) heteroaryl,  
(10) aryl-C<sub>1-10</sub> alkyl, and  
10 (11) heteroaryl-C<sub>1-10</sub> alkyl,

each R<sup>i</sup> may be unsubstituted or substituted with one, two, or three substituents selected from R<sup>h</sup>;

m is selected from 1 and 2; and

n is selected from 1, 2, and 3;

- 15 and pharmaceutically acceptable salts thereof.

In one embodiment of the present invention, R<sup>1</sup> is selected from:

- (1) C<sub>1-10</sub> alkyl,  
(2) C<sub>3-10</sub> cycloalkyl,  
(3) cycloheteroalkyl,  
20 (4) aryl, and  
(5) heteroaryl,

wherein each alkyl is optionally substituted with one to three substituents independently selected from R<sup>a</sup>, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one to three substituents  
25 independently selected from R<sup>b</sup>.

In one class of this embodiment, R<sup>1</sup> is selected from:

- (1) C<sub>1-4</sub> alkyl,  
(2) C<sub>3-10</sub> cycloalkyl-,  
(3) cycloheteroalkyl,  
30 (4) phenyl, and  
(5) pyridyl,

wherein each alkyl is optionally substituted with one R<sup>a</sup> substituent, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one to three substituents independently selected from R<sup>b</sup>.

In one subclass of this embodiment, R<sup>1</sup> is selected from:

- (1) isopropyl,
- (2) isobutyl,
- (3) n-propyl,
- 5 (4) cyclopropyl,
- (5) cyclobutyl,
- (6) cyclopentyl,
- (7) cyclohexyl,
- (8) piperidiny,
- 10 (9) phenyl, and
- (10) pyridyl,

wherein each alkyl is optionally substituted with one R<sup>a</sup> substituent, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one to three substituents independently selected from R<sup>b</sup>.

- 15 In another embodiment of the present invention, R<sup>1</sup> is selected from aryl and pyridyl, wherein aryl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>.

- In one class of this embodiment, R<sup>1</sup> is phenyl, wherein phenyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>.
- 20

In one subclass, R<sup>1</sup> is selected from:

- (1) phenyl,
- (2) 4-fluorophenyl,
- (3) 2-chlorophenyl,
- 25 (4) 3-chlorophenyl,
- (5) 4-chlorophenyl,
- (6) 3-cyanophenyl,
- (7) 4-cyanophenyl,
- (8) 4-methylphenyl,
- 30 (9) 4-isopropylphenyl,
- (10) 4-biphenyl,
- (11) 4-bromophenyl,
- (12) 4-iodophenyl,
- (13) 2,4-dichlorophenyl, and

(14) 2-chloro-4-fluorophenyl.

In another subclass,  $R^1$  is selected from:

- (1) phenyl, and
- (2) 4-chlorophenyl.

5 In yet another embodiment of the present invention,  $R^2$  is selected from:

- (1)  $C_{1-10}$ alkyl,
- (2)  $C_{3-10}$ cycloalkyl,
- (3) cycloheteroalkyl,
- 10 (4) aryl,
- (5) heteroaryl,
- (6) -OR<sup>d</sup>,
- (7) -NRCR<sup>d</sup>, and
- (8) -CO<sub>2</sub>R<sup>d</sup>, and

15 wherein each alkyl is optionally substituted with one, two or three substituents independently selected from  $R^a$ , and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one, two or three substituents independently selected from  $R^b$ .

In a class of this embodiment,  $R^2$  is selected from:

- 20 (1) cyclobutyl,
- (2) cyclopentyl,
- (3) cyclohexyl,
- (4) pyrrolidinyl,
- (5) pyrimidinyl,
- 25 (6) benzoxazolyl,
- (7) dihydroindolyl,
- (8) dihydroquinolinyl,
- (9) benzotriazolyl,
- (10) thiophenyl,
- 30 (11) indolyl,
- (12) indazolyl,
- (13) pyrrolidinyl,
- (14) pyridazinyl
- (15) triazolyl,
- 35 (16) azaindolyl,

- (17) cyclobutylmethoxy,
- (18) phenyl,
- (19) pyridyl,
- (20) -NR<sup>c</sup>R<sup>d</sup>, and
- 5 (21) -CO<sub>2</sub>R<sup>d</sup>,

wherein each alkyl is optionally substituted with one or two R<sup>a</sup> substituents and each phenyl or pyridyl is independently with one to three R<sup>b</sup> substituents.

In another class of the present invention, R<sup>2</sup> is phenyl,  
wherein phenyl is optionally substituted with one, two, three or four substituents  
10 independently selected from R<sup>b</sup>.

In one subclass, R<sup>2</sup> is selected from:

- (1) phenyl,
- (2) 4-fluorophenyl,
- (3) 2-chlorophenyl,
- 15 (4) 3-chlorophenyl,
- (5) 4-chlorophenyl,
- (6) 3-cyanophenyl,
- (7) 4-cyanophenyl,
- (8) 4-methylphenyl,
- 20 (9) 4-isopropylphenyl,
- (10) 4-biphenyl,
- (11) 4-bromophenyl,
- (12) 4-iodophenyl,
- (13) 2,4-dichlorophenyl, and
- 25 (14) 2-chloro-4-fluorophenyl.

In another subclass of this class of the present invention, R<sup>2</sup> is selected  
from:

- (1) phenyl, and
- (2) 4-chlorophenyl.

30 In another embodiment of the present invention, R<sup>3</sup> is selected from:

- (1) hydrogen,
- (2) methyl, and
- (3) ethyl.

In one class of this embodiment of the present invention, R<sup>3</sup> is selected  
35 from methyl and ethyl.

In a subclass of this class, R<sup>3</sup> is methyl.

In another class of this embodiment of the present invention, R<sup>3</sup> is hydrogen.

In another embodiment of the present invention, R<sup>4</sup> is selected from:

- 5 (1) hydrogen,  
(2) C<sub>1-10</sub>alkyl,  
(3) -OR<sup>c</sup>,  
(4) -NR<sup>c</sup>R<sup>d</sup>,  
(5) -NH(CO)OR<sup>d</sup>,  
10 (6) aryl, and  
(7) heteroaryl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>, and aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>.

15

In one class of this embodiment of the present invention, R<sup>4</sup> is selected from:

- (1) hydrogen,  
(2) methyl,  
20 (3) ethyl,  
(4) propyl,  
(5) butyl,  
(6) pentyl,  
(7) isopropyl,  
25 (8) -OH,  
(9) -O-aryl,  
(10) -NH<sub>2</sub>,  
(11) -NH-aryl,  
(12) -NH(CO)OR<sup>d</sup>,  
30 (13) phenyl, and  
(14) pyridyl,

wherein alkyl groups may be optionally substituted with one, two, or three substituents independently selected from R<sup>a</sup>, and phenyl, aryl and pyridyl are


optionally substituted with one, two, three or four substituents independently selected from  $R^b$ .

In a subclass of this embodiment of the present invention,  $R^4$  is selected from:

- 5 (1) hydrogen,
- (2) methyl,
- (3)  $-OH$ ,
- (4)  $-O$ -phenyl,
- (5)  $-NH_2$ ,
- 10 (6)  $-NH$ -phenyl,
- (7)  $-NH(CO)O$ -alkyl,
- (8) phenyl, and
- (9) pyridyl,

wherein methyl and alkyl are optionally substituted with one, two, three or four substituents independently selected from  $R^a$ , and phenyl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from  $R^b$ .

In one embodiment of the present invention,  $R^6$  is hydrogen. When  $R^6$  is hydrogen, the structural formula I may be represented as structural formula IA:



(IA)

In another embodiment of the present invention,  $R^6$  is selected from:

- 25 (1)  $C_{1-4}$ alkyl,
- (2)  $C_{2-4}$ alkenyl,
- (3)  $C_{2-4}$ alkynyl,
- (4)  $-OR^d$ ,
- (5) halogen, and
- (6)  $-CN$ ,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ .

In a class of this embodiment,  $R^6$  is selected from:

- (1) methyl,
- 5 (2) hydroxyl,
- (3) halogen, and
- (4)  $-\text{CN}$ ;

wherein methyl is optionally substituted with one to three  $R^a$  substituents.

In one subclass of this class,  $R^6$  is selected from:

- 10 (1) methyl,
- (2) hydroxyl,
- (3) halogen, and
- (4)  $-\text{CN}$ .

In one embodiment of the present invention, **A** is selected from the  
 15 group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofurany, and tetrahydropyrany, each optionally substituted with one, two, three or four groups independently selected from oxo and  $R^b$ , with the proviso that when  $R^1$  and  $R^2$  are aryl and **A** is cyclobutyl, the cyclobutyl group is either unsubstituted, or is not  
 20 substituted with an  $R^a$  substituent selected from the group consisting of  $-\text{OCH}_3$ ,  $-\text{C}(\text{O})\text{NR}^c\text{R}^d$ , and  $-\text{NR}^c\text{C}(\text{O})\text{NR}^c\text{R}^d$  and is not substituted with an  $R^h$  substituent selected from the group consisting of: aryl $\text{C}_{1-4}$ alkyl, heteroaryl,  $-\text{NR}^c\text{S}(\text{O})_m\text{R}^d$ ,  $-\text{S}(\text{O})_m\text{NR}^c\text{R}^d$ ,  $-\text{O}(\text{CR}^e\text{R}^f)_n\text{NR}^c\text{R}^d$ ,  $-\text{C}(\text{O})\text{R}^c$ , and  $-\text{C}(\text{O})\text{NR}^c\text{R}^d$ .

In another embodiment of the present invention, **A** is selected from the  
 25 group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, hexahydroazepiny, morpholiny, tetrahydrofurany, tetrahydropyrany, and tetrahydrothienyl, each optionally substituted with one, two, or three groups independently selected from oxo and  $R^b$ .

In one class of this embodiment of the present invention, **A** is selected  
 30 from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofurany, and tetrahydropyrany, each optionally substituted with one or two groups independently selected from oxo and  $R^b$ .

In one subclass of this class of the present invention, **A** is selected from  
 35 the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny,

pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, each optionally substituted with one or two groups independently selected from oxo and  $R^b$ , with the proviso that when **A** is cyclopropyl, the cyclopropyl group is not bis 2,2-dichlorocyclopropyl.

5 In another class of this embodiment of the present invention, **A** is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, azetidyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, each substituted with one group independently selected from  $R^b$ , with the proviso that when **A** is cyclopropyl,  
 10 cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl,  $R^a$  is not selected from the group consisting of  $-NR^cS(O)_mR^d$ ,  $-NR^cC(O)R^d$ ,  $-NR^cC(O)OR^d$ , and  $-NR^cC(O)NR^dR^d$ .

In another embodiment of the present invention,  $R^a$  is selected from:

- (1)  $-OR^c$
- (2) halogen,
- 15 (3)  $-S(O)_mR^c$ ,
- (4)  $-SRC$ ,
- (5)  $-S(O)_2OR^c$ ,
- (6)  $-S(O)_mNR^cR^d$ ,
- (7)  $-NR^cR^d$ ,
- 20 (8)  $-C(O)R^c$ ,
- (9)  $-CO_2R^i$ ,
- (10)  $-CN$ ,
- (11)  $-OC(O)NR^cR^d$ ,
- (12)  $CF_3$ ,
- 25 (13)  $-OCF_3$ ,
- (14)  $C_{3-8}cycloalkyl$ ,
- (15) cycloheteroalkyl, and
- (16) oxo.

30 In one class of this embodiment of the present invention,  $R^a$  is selected from:

- (1)  $-OR^c$ ,
- (2) halogen,
- (3)  $-S(O)_mR^c$ ,
- (4)  $-C(O)R^c$ , and



(5)  $-\text{CO}_2\text{R}^i$ .

In one subclass of this class of the invention,  $\text{R}^a$  is selected from:

- (1)  $-\text{O}-\text{aryl}$ ,
- (2)  $-\text{OH}$ ,
- 5 (3) bromo,
- (4) chloro,
- (5)  $-\text{S}(\text{O})_m-\text{aryl}$ ,
- (6)  $-\text{C}(\text{O})-\text{aryl}$ , and
- (7)  $-\text{CO}_2-\text{alkyl}$ .

10 In another embodiment of the present invention,  $\text{R}^b$  is selected from:

- (1)  $\text{R}^a$ ,
- (2)  $\text{C}_{1-10}\text{alkyl}$ ,
- (3) aryl, and
- (4) heteroaryl.

15 In one class of this embodiment of the present invention,  $\text{R}^b$  is selected from:

- (1)  $\text{R}^a$ ,
- (2) methyl,
- (3) ethyl,
- 20 (4) propyl,
- (5) phenyl,
- (6) benzyl, and
- (7) pyridyl.

25 In another embodiment of the present invention,  $\text{R}^c$  and  $\text{R}^d$  are selected from:

- (1) hydrogen,
- (2)  $\text{C}_{2-10}\text{alkyl}$ ,
- (3)  $\text{C}_{2-10}\text{alkenyl}$ ,
- (4)  $\text{C}_{2-10}\text{alkynyl}$ ,
- 30 (5) cycloalkyl,
- (6) cycloalkyl- $\text{C}_{1-10}\text{alkyl}$ ,
- (7) cycloheteroalkyl,
- (8) cycloheteroalkyl- $\text{C}_{1-10}\text{alkyl}$ ,
- (9) aryl,

- (10) heteroaryl,
- (11) aryl-C<sub>1-10</sub>alkyl, and
- (12) heteroaryl-C<sub>1-10</sub>alkyl, or

R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic  
5 ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected  
from oxygen, sulfur and N-R<sup>g</sup>,  
each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three  
substituents selected from R<sup>h</sup>.

In one class of this embodiment of the present invention, R<sup>c</sup> and R<sup>d</sup>  
10 are selected from:

- (1) hydrogen,
- (2) C<sub>2-10</sub>alkyl,
- (3) aryl, and
- (4) heteroaryl, or

15 R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic  
ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected  
from oxygen, sulfur and N-R<sup>g</sup>,  
each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two, or three  
substituents selected from R<sup>h</sup>.

20 In one subclass of this class of the invention, R<sup>c</sup> and R<sup>d</sup> are  
independently selected from:

- (1) hydrogen,
- (2) ethyl,
- (3) propyl,
- 25 (4) butyl, and
- (5) phenyl,

each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three  
substituents selected from R<sup>h</sup>.

In another embodiment of the present invention, R<sup>g</sup> is selected from:

- 30 (1) C<sub>1-10</sub>alkyl, and
- (2) -C(O)R<sup>e</sup>.

In another embodiment of the present invention, R<sup>h</sup> is selected from:

- (1) halogen,
- (2) C<sub>1-10</sub>alkyl,
- 35 (3) C<sub>3-8</sub>cycloalkyl,

- (4) cycloheteroalkyl,
- (5) aryl,
- (6) heteroarylC<sub>1-4</sub>alkyl,
- (7) -OR<sup>e</sup>,
- 5 (8) -S(O)<sub>m</sub>R<sup>e</sup>,
- (9) -SR<sup>e</sup>,
- (10) -S(O)<sub>2</sub>OR<sup>e</sup>,
- (11) -NR<sup>e</sup>R<sup>f</sup>,
- (12) -CO<sub>2</sub>R<sup>e</sup>,
- 10 (13) -CO<sub>2</sub>(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>CONR<sup>e</sup>R<sup>f</sup>,
- (14) -OC(O)R<sup>e</sup>,
- (15) -CN,
- (16) -NR<sup>e</sup>C(O)R<sup>f</sup>,
- (17) -OC(O)NR<sup>e</sup>R<sup>f</sup>,
- 15 (18) -NR<sup>e</sup>C(O)OR<sup>f</sup>,
- (19) -NR<sup>e</sup>C(O)NR<sup>e</sup>R<sup>f</sup>,
- (20) -CF<sub>3</sub>, and
- (21) -OCF<sub>3</sub>.

In a class of this embodiment, R<sup>h</sup> is selected from:

- 20 (1) halogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>3-8</sub>cycloalkyl,
- (4) cycloheteroalkyl,
- (5) aryl,
- 25 (6) heteroarylC<sub>1-4</sub>alkyl,
- (7) -CN,
- (8) -CF<sub>3</sub>, and
- (9) -OCF<sub>3</sub>.

Particular novel compounds of structural formula I which may be employed in the methods, uses and compositions of the present invention, include:

- 30 (1) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-4-carboxamide;
- (2) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(2-pyridyl)-piperidine-3-carboxamide;

- (3) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-methylmorpholine-2-carboxamide;
- (4) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-pyrrolidine-2(S)-carboxamide;
- 5 (5) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-2-carboxamide;
- (6) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2,2-dimethyl-tetrahydropyran-4-carboxamide;
- (7) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclopentane-carboxamide;
- 10 (8) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- (9) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclohexane-carboxamide;
- 15 (10) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;
- (11) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-cyclopentanone-3-carboxamide;
- (12) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;
- 20 (13) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- (14) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-2-methylazetidine-2-carboxamide;
- 25 (15) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzyl-piperidine-2-carboxamide;
- (16) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-pyrrolidine-3-carboxamide;
- (17) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide;
- 30 (18) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-pyrrolidine-2(S)-carboxamide;
- (19) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-2-methylpyrrolidine-2(S)-carboxamide;

- (20) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-methyltetrahydrofuran-2-carboxamide;
- (21) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-tetrahydrofuran-2-carboxamide;
- 5 (22) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide;
- (23) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-oxopyrrolidine-5-carboxamide;
- (24) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-5-methyl-2-oxopyrrolidine-5-carboxamide;
- 10 (25) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzylpyrrolidine-2-carboxamide;
- (26) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(phenylamino)-cyclopentanecarboxamide;
- 15 (27) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (28) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopropanecarboxamide;
- (29) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopropanecarboxamide;
- 20 (30) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-chlorophenyl)cyclohexanecarboxamide;
- (31) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-2-phenyltetrahydrofuran-2-carboxamide;
- 25 (32) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3,5-difluorophenyl)cyclopentanecarboxamide;
- (33) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(4-pyridyl)cyclopentanecarboxamide;
- (34) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- 30 (35) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (36) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl-N-oxide)cyclopentanecarboxamide;

- (37) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (38) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- 5 (39) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (40) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide;
- (41) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclohexane-carboxamide;
- 10 (42) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclopentane-carboxamide;
- (43) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclohexanecarboxamide;
- 15 (44) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclopentanecarboxamide;
- (45) *N*-[2-(3-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (46) *N*-[2-(3-fluoropyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- 20 (47) *N*-[2-(2-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (48) *N*-[2-(4-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- 25 (49) *N*-[3-(5-chloro-2-pyridyl)-2-phenyl-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (50) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (51) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- 30 (52) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- (53) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;

- (54) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- (55) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- 5 (56) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- (57) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (58) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- 10 (59) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (60) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- 15 (61) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (62) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- (63) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- 20 (64) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- (65) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- 25 (66) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- (67) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (68) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-cyclobutanecarboxamide;
- 30 (69) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylcyclobutanecarboxamide;
- (70) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-ethylcyclobutanecarboxamide;

- (71) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-propylcyclobutanecarboxamide;
- (72) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-benzylcyclobutanecarboxamide;
- 5 (73) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-isopropylcyclobutanecarboxamide;
- (74) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-3-azetidine-3-carboxamide;
- (75) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-10 butoxycarbonyl-2(*S*)-azetidinecarboxamide;
- (76) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 1);
- (77) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 2);
- 15 (78) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (79) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (80) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-20 fluorophenyl)cyclopentanecarboxamide;
- (81) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- (82) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- 25 (83) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- (84) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- (85) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-30 fluorophenyl)cyclopentanecarboxamide;
- (86) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- (87) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;



- (88) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- (89) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- 5 (90) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (91) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (92) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- 10 (93) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (94) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (95) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- 15 (96) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- (97) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- 20 (98) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- (99) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- 25 (100) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- (101) *N*-(2*S*,3*S*)-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylsulfonylcyclobutanecarboxamide;
- (102) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-3-ethylazetidine-3-carboxamide hydrochloride;
- 30 (103) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 1);
- (104) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 2);
- 35 and pharmaceutically acceptable salts thereof.

One subclass of compounds of the present invention includes compounds wherein R<sup>1</sup> is 4-chlorophenyl, R<sup>2</sup> is 4-chlorophenyl and R<sup>3</sup> is methyl. Particular compounds of this subclass include:

- (1) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- (2) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide;
- (3) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide;
- (4) N-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyl)-cyclohexanecarboxamide;
- (5) N-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide;
- (6) N-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclohexane-carboxamide; and
- (7) N-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclopentane-carboxamide;

and pharmaceutically acceptable salts thereof.

The compounds of structural formula I are modulators of the CB1 receptor. In particular, the compounds of structural formula I are antagonists or inverse agonists of the CB1 receptor.

An "agonist" is a compound (hormone, neurotransmitter or synthetic compound) which binds to a receptor and produces a response, such as contraction, relaxation, secretion, change in enzyme activity, etc. An "antagonist" is a compound which attenuates the effect of an agonist. An "inverse agonist" is a compound which acts on a receptor but produces the opposite effect produced by the agonist of the particular receptor.

"Alkyl", as well as other groups having the prefix "alk", such as alkoxy, alkanoyl, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

“Alkynyl” means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

5 “Cycloalkyl” means mono- or bicyclic or bridged saturated carbocyclic rings, each of which having from 3 to 10 carbon atoms. The term also includes monocyclic rings fused to an aryl group in which the point of attachment is on the non-aromatic portion. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, decahydronaphthyl,  
10 indanyl, and the like.

“Aryl” means mono- or bicyclic aromatic rings containing only carbon atoms. The term also includes aryl group fused to a monocyclic cycloalkyl or monocyclic cycloheteroalkyl group in which the point of attachment is on the aromatic portion. Examples of aryl include phenyl, naphthyl, indanyl, indenyl,  
15 tetrahydronaphthyl, 2,3-dihydrobenzofuranyl, dihydrobenzopyranyl, 1,4-benzodioxanyl, and the like.

“Heteroaryl” means a mono- or bicyclic aromatic ring containing at least one heteroatom selected from N, O and S, with each ring containing 5 to 6 atoms. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl,  
20 pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, furo(2,3-b)pyridyl, quinolyl, indolyl, isoquinolyl, and the like. The heteroaryl ring may be substituted on one or more carbon atoms.


25 “Cycloheteroalkyl” means mono- or bicyclic or bridged saturated rings containing at least one heteroatom selected from N, S and O, each of said ring having from 3 to 10 atoms in which the point of attachment may be carbon or nitrogen. The term also includes monocyclic heterocycle fused to an aryl or heteroaryl group in which the point of attachment is on the non-aromatic portion. Examples of  
30 “cycloheteroalkyl” include pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, 2,3-dihydrofuro(2,3-b)pyridyl, benzoxazinyl, tetrahydrohydroquinolyl, morpholinyl, tetrahydroisoquinolyl, dihydroindolyl, and the like. The term also includes partially unsaturated monocyclic rings that are not aromatic, such as 2- or 4-pyridones attached through the nitrogen or N-substituted-(1H, 3H)-pyrimidine-2,4-diones (N-substituted

uracils). The cycloheteroalkyl ring may be substituted on the ring carbons and/or the ring nitrogens.

"Halogen" includes fluorine, chlorine, bromine and iodine.

When any variable (e.g., R<sup>1</sup>, R<sup>d</sup>, etc.) occurs more than one time in  
5 any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

Under standard nomenclature used throughout this disclosure, the  
terminal portion of the designated side chain is described first, followed by the  
10 adjacent functionality toward the point of attachment. For example, a C<sub>1-5</sub>  
alkylcarbonylamino C<sub>1-6</sub> alkyl substituent is equivalent to



In choosing compounds of the present invention, one of ordinary skill  
in the art will recognize that the various substituents, i.e. R<sup>1</sup>, R<sup>2</sup>, etc., are to be chosen  
15 in conformity with well-known principles of chemical structure connectivity and stability.

The term "substituted" shall be deemed to include multiple degrees of  
substitution by a named substituent. Where multiple substituent moieties are  
disclosed or claimed, the substituted compound can be independently substituted by  
20 one or more of the disclosed or claimed substituent moieties, singly or plurally. By  
independently substituted, it is meant that the (two or more) substituents can be the  
same or different.


Compounds of Formula I may contain one or more asymmetric centers  
and can thus occur as racemates and racemic mixtures, single enantiomers,  
25 enantiomeric mixtures, diastereomeric mixtures and individual diastereomers. The  
present invention is meant to comprehend all such isomeric forms of the compounds  
of Formula I.

Some of the compounds described herein contain olefinic double  
bonds, and unless specified otherwise, are meant to include both E and Z geometric  
30 isomers.

Tautomers are defined as compounds that undergo rapid proton shifts  
from one atom of the compound to another atom of the compound. Some of the

compounds described herein may exist as tautomers with different points of attachment of hydrogen. Such an example may be a ketone and its enol form known as keto-enol tautomers. The individual tautomers as well as mixture thereof are encompassed with compounds of Formula I. By way of illustration, tautomers

5 included in this definition include, but are not limited to:



Compounds of the Formula I may be separated into diastereoisomeric pairs of enantiomers by, for example, fractional crystallization from a suitable solvent, for example MeOH or ethyl acetate or a mixture thereof. The pair of enantiomers  
 10 thus obtained may be separated into individual stereoisomers by conventional means, for example by the use of an optically active amine as a resolving agent or on a chiral HPLC column.

Alternatively, any enantiomer of a compound of the general Formula I may be obtained by stereospecific synthesis using optically pure starting materials or  
 15 reagents of known configuration.

Furthermore, some of the crystalline forms for compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds of the instant invention may form solvates with water or common organic solvents. Such  
 20 solvates are encompassed within the scope of this invention.

It is generally preferable to administer compounds of the present invention as enantiomerically pure formulations. Racemic mixtures can be separated

into their individual enantiomers by any of a number of conventional methods. These include chiral chromatography, derivatization with a chiral auxiliary followed by separation by chromatography or crystallization, and fractional crystallization of diastereomeric salts.

5                   The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly  
10 preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-  
15 diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like. The term  
20 "pharmaceutically acceptable salt" further includes all acceptable salts such as acetate, lactobionate, benzenesulfonate, laurate, benzoate, malate, bicarbonate, maleate, bisulfate, mandelate, bitartrate, mesylate, borate, methylbromide, bromide, methylnitrate, calcium edetate, methylsulfate, camsylate, mucate, carbonate, napsylate, chloride, nitrate, clavulanate, N-methylglucamine, citrate, ammonium salt,  
25 dihydrochloride, oleate, edetate, oxalate, edisylate, pamoate (embonate), estolate, palmitate, esylate, pantothenate, fumarate, phosphate/diphosphate, gluceptate, polygalacturonate, gluconate, salicylate, glutamate, stearate, glycolylarsanilate, sulfate, hexylresorcinolate, subacetate, hydrabamine, succinate, hydrobromide, tannate, hydrochloride, tartrate, hydroxynaphthoate, teoclate, iodide, tosylate, trifluoro acetate,  
30 isothionate, triethiodide, lactate, pantoate, valerate, and the like which can be used as a dosage form for modifying the solubility or hydrolysis characteristics or can be used in sustained release or pro-drug formulations.

It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

Compounds of this invention are modulators of the CB1 receptor and as such are useful as psychotropic drugs in the treatment of psychosis, memory deficits, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders including multiple sclerosis and Guillain-Barre syndrome and the inflammatory  
5 sequelae of viral encephalitis, cerebral vascular accidents, and head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, movement disorders, and schizophrenia. The compounds are also useful for the treatment of substance abuse disorders, particularly to opiates, alcohol, marijuana, and nicotine. The compounds are also useful for the treatment of obesity or eating disorders associated with  
10 excessive food intake and complications associated therewith. The compounds are also useful for the treatment of constipation and chronic intestinal pseudo-obstruction. The compounds are also useful for the treatment of cirrhosis of the liver. The compounds are also useful for the treatment of asthma.

The terms "administration of" and or "administering a" compound  
15 should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to the individual in need of treatment.

The administration of the compound of structural formula I in order to practice the present methods of therapy is carried out by administering an effective amount of the compound of structural formula I to the patient in need of such  
20 treatment or prophylaxis. The need for a prophylactic administration according to the methods of the present invention is determined via the use of well known risk factors. The effective amount of an individual compound is determined, in the final analysis, by the physician in charge of the case, but depends on factors such as the exact disease to be treated, the severity of the disease and other diseases or conditions from  
25 which the patient suffers, the chosen route of administration other drugs and treatments which the patient may concomitantly require, and other factors in the physician's judgment.

The utilities of the present compounds in these diseases or disorders may be demonstrated in animal disease models that have been reported in the  
30 literature. The following are examples of such animal disease models: a) suppression of food intake and resultant weight loss in rats (Life Sciences 1998, 63, 113-117); b) reduction of sweet food intake in marmosets (Behavioural Pharm. 1998, 9, 179-181); c) reduction of sucrose and ethanol intake in mice (Psychopharm. 1997, 132, 104-106); d) increased motor activity and place conditioning in rats (Psychopharm. 1998,

135, 324-332; Psychopharmacol 2000, 151: 25-30); e) spontaneous locomotor activity in mice (J. Pharm. Exp. Ther. 1996, 277, 586-594); f) reduction in opiate self-administration in mice (Sci. 1999, 283, 401-404); g) bronchial hyperresponsiveness in sheep and guinea pigs as models for the various phases of asthma (for example, see  
5 W. M. Abraham et al., "α<sub>4</sub>-Integrins mediate antigen-induced late bronchial responses and prolonged airway hyperresponsiveness in sheep." J. Clin. Invest. 93, 776 (1993) and A. A. Y. Milne and P. P. Piper, "Role of VLA-4 integrin in leucocyte recruitment and bronchial hyperresponsiveness in the guinea-pig." Eur. J. Pharmacol., 282, 243 (1995)); h) mediation of the vasodilated state in advanced liver cirrhosis  
10 induced by carbon tetrachloride (Nature Medicine, 2001, 7 (7), 827-832); i) amitriptyline-induced constipation in cynomolgus monkeys is beneficial for the evaluation of laxatives (Biol. Pharm. Bulletin (Japan), 2000, 23(5), 657-9); j) neuropathology of paediatric chronic intestinal pseudo-obstruction and animal models related to the neuropathology of paediatric chronic intestinal pseudo-obstruction  
15 (Journal of Pathology (England), 2001, 194 (3), 277-88).

The compounds of this invention are also useful for the treatment or prevention of conditions associated with, caused by, or resulting from obesity. The compounds are useful for reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy.

20 The magnitude of prophylactic or therapeutic dose of a compound of Formula I will, of course, vary with the nature of the severity of the condition to be treated and with the particular compound of Formula I and its route of administration. It will also vary according to the age, weight and response of the individual patient. In general, the daily dose range lie within the range of from about 0.001 mg to about  
25 100 mg per kg body weight of a mammal, preferably 0.01 mg to about 50 mg per kg, and most preferably 0.1 to 10 mg per kg, in single or divided doses. On the other hand, it may be necessary to use dosages outside these limits in some cases.

For use where a composition for intravenous administration is employed, a suitable dosage range is from about 0.001 mg to about 100 mg  
30 (preferably from 0.01 mg to about 50 mg, more preferably 0.1 mg to 10 mg) of a compound of Formula I per kg of body weight per day.

In the case where an oral composition is employed, a suitable dosage range is, e.g. from about 0.01 mg to about 1000 mg of a compound of Formula I per day, preferably from about 0.1 mg to about 10 mg per day. For oral administration,  
35 the compositions are preferably provided in the form of tablets containing from 0.01



to 1,000 mg, preferably 0.01, 0.05, 0.1, 0.5, 1, 2.5, 5, 10, 15, 20, 25, 30, 40, 50, 100, 250, 500, 750 or 1000 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated.

5 For the treatment of diseases of the eye, ophthalmic preparations for ocular administration comprising 0.001-1% by weight solutions or suspensions of the compounds of Formula I in an acceptable ophthalmic formulation may be used.

Another aspect of the present invention provides pharmaceutical compositions which comprises a compound of Formula I and a pharmaceutically acceptable carrier. The term "composition", as in pharmaceutical composition, is  
10 intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or  
15 interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and pharmaceutically acceptable excipients.

Any suitable route of administration may be employed for providing a  
20 mammal, especially a human, with an effective dosage of a compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like.

The pharmaceutical compositions of the present invention comprise a  
25 compound of Formula I as an active ingredient or a pharmaceutically acceptable salt thereof, and may also contain a pharmaceutically acceptable carrier and optionally other therapeutic ingredients. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. In particular, the term  
30 "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic bases or acids and organic bases or acids.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular

(ophthalmic), pulmonary (aerosol inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulizers. The compounds may also be delivered as powders which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery systems for inhalation are metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons and dry powder inhalation (DPI) aerosol, which may be formulated as a dry powder of a compound of Formula I with or without additional excipients.

Suitable topical formulations of a compound of formula I include transdermal devices, aerosols, creams, solutions, ointments, gels, lotions, dusting powders, and the like. The topical pharmaceutical compositions containing the compounds of the present invention ordinarily include about 0.005% to 5% by weight of the active compound in admixture with a pharmaceutically acceptable vehicle. Transdermal skin patches useful for administering the compounds of the present invention include those well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course be continuous rather than intermittent throughout the dosage regimen.

In practical use, the compounds of Formula I can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for

example, powders, capsules and tablets, with the solid oral preparations being preferred over the liquid preparations. Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be  
5 coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, the compounds of Formula I may also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; 3,630,200 and 4,008,719.

10           Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules (including timed release and sustained release formulations), pills, cachets, powders, granules or tablets each containing a predetermined amount of the active ingredient, as a powder or granules or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid,  
15 an oil-in-water emulsion or a water-in-oil liquid emulsion, including elixirs, tinctures, solutions, suspensions, syrups and emulsions. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and  
20 intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. For example, a tablet may be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine, the active ingredient in a free-flowing  
25 form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Desirably, each tablet contains from 0.01 to 1,000 mg, particularly 0.01, 0.05, 0.1, 0.5, 1, 2.5, 3, 5, 6, 10, 15, 25, 50, 75, 100, 125, 150, 175, 180, 200,  
30 225, 500, 750 and 1,000 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. and each cachet or capsule contains from about 0.01 to 1,000 mg, particularly 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 3, 5, 6, 10, 15, 25, 50, 75, 100, 125, 150, 175, 180, 200, 225, 500, 750 and 1,000

milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated.

Additional suitable means of administration of the compounds of the present invention include injection, intravenous bolus or infusion, intraperitoneal, subcutaneous, intramuscular and topical, with or without occlusion.

Exemplifying the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Also exemplifying the invention is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

The dose may be administered in a single daily dose or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, based on the properties of the individual compound selected for administration, the dose may be administered less frequently, e.g., weekly, twice weekly, monthly, etc. The unit dosage will, of course, be correspondingly larger for the less frequent administration.

When administered via intranasal routes, transdermal routes, by rectal or vaginal suppositories, or through a continual intravenous solution, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The following are examples of representative pharmaceutical dosage forms for the compounds of Formula I:

Injectable Suspension (I.M.) mg/mL

Compound of Formula I	10
Methylcellulose	5.0
Tween 80	0.5
Benzyl alcohol	9.0
Benzalkonium chloride	1.0
Water for injection to a total volume of 1 mL	

<u>Tablet</u>	<u>mg/tablet</u>
Compound of Formula I	25
Microcrystalline Cellulose	415
Povidone	14.0
5 Pregelatinized Starch	43.5
Magnesium Stearate	2.5
	500

<u>Capsule</u>	<u>mg/capsule</u>
10 Compound of Formula I	25
Lactose Powder	573.5
Magnesium Stearate	1.5
	600

<u>15 Aerosol</u>	<u>Per canister</u>
Compound of Formula I	24 mg
Lecithin, NF Liq. Conc.	1.2 mg
Trichlorofluoromethane, NF	4.025 g
Dichlorodifluoromethane, NF	12.15 g

- 20 Compounds of Formula I may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of Formula I. When a
- 25 compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of Formula I is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of Formula I. Examples of other active ingredients that
- 30 may be combined with a compound of Formula I include, but are not limited to: antipsychotic agents, cognition enhancing agents, anti-migraine agents, anti-asthmatic agents, antiinflammatory agents, anxiolytics, anti-Parkinson's agents, anti-epileptics, anorectic agents, serotonin reuptake inhibitors, and other anti-obesity agents, which may be administered separately or in the same pharmaceutical compositions.

The present invention also provides a method for the treatment or prevention of a CB1 receptor modulator mediated disease, which method comprises administration to a patient in need of such treatment or at risk of developing a CB1 receptor modulator mediated disease of an amount of a CB1 receptor modulator and  
5 an amount of one or more active ingredients, such that together they give effective relief.

In a further aspect of the present invention, there is provided a pharmaceutical composition comprising a CB1 receptor modulator and one or more active ingredients, together with at least one pharmaceutically acceptable carrier or  
10 excipient.

Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and one or more active ingredients for the manufacture of a medicament for the treatment or prevention of a CB1 receptor modulator mediated disease. In a further or alternative aspect of the present  
15 invention, there is therefore provided a product comprising a CB1 receptor modulator and one or more active ingredients as a combined preparation for simultaneous, separate or sequential use in the treatment or prevention of CB1 receptor modulator mediated disease. Such a combined preparation may be, for example, in the form of a twin pack.

20 It will be appreciated that for the treatment or prevention of eating disorders, including obesity, bulimia nervosa and compulsive eating disorders, a compound of the present invention may be used in conjunction with other anorectic agents.

The present invention also provides a method for the treatment or  
25 prevention of eating disorders, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of an anorectic agent, such that together they give effective relief.

Suitable anorectic agents of use in combination with a compound of the present invention include, but are not limited to, aminorex, amphetamine, amphetamine, benzphetamine, chlorphentermine, clobenzorex, cloforex, clominorex, clortermine, cyclexedrine, dexfenfluramine, dextroamphetamine, diethylpropion,  
30 diphemethoxidine, *N*-ethylamphetamine, fenbutrazate, fenfluramine, fenisorex, fenproporex, fludorex, fluminorex, furfurylmethylamphetamine, levamphetamine, levophacetoperane, mazindol, mefenorex, metamfepramone, methamphetamine,

norpseudoephedrine, pentorex, phendimetrazine, phenmetrazine, phentermine, phenylpropanolamine, picilorex and sibutramine; and pharmaceutically acceptable salts thereof.

5 A particularly suitable class of anorectic agent are the halogenated amphetamine derivatives, including chlorphentermine, cloforex, clortermine, dexfenfluramine, fenfluramine, picilorex and sibutramine; and pharmaceutically acceptable salts thereof

Particularly preferred halogenated amphetamine derivatives of use in combination with a compound of the present invention include: fenfluramine and  
10 dexfenfluramine, and pharmaceutically acceptable salts thereof.

It will be appreciated that for the treatment or prevention of obesity, the compounds of the present invention may also be used in combination with a selective serotonin reuptake inhibitor (SSRI).

The present invention also provides a method for the treatment or  
15 prevention of obesity, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of an SSRI, such that together they give effective relief.

Suitable selective serotonin reuptake inhibitors of use in combination with a compound of the present invention include: fluoxetine, fluvoxamine,  
20 paroxetine sertraline, and imipramine, and pharmaceutically acceptable salts thereof.

It will be appreciated that for the treatment or prevention of obesity, the compounds of the present invention may also be used in combination with an opioid antagonist.

The present invention also provides a method for the treatment or  
25 prevention of obesity, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of an opioid antagonist, such that together they give effective relief.

Suitable opioid antagonists of use in combination with a compound of the present invention include: naltrexone, 3-methoxynaltrexone, naloxone and  
30 nalmeffene, and pharmaceutically acceptable salts thereof.

It will be appreciated that for the treatment or prevention of obesity, the compounds of the present invention may also be used in combination with another anti-obesity agent.

The present invention also provides a method for the treatment or prevention of obesity, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of another anti-obesity agent, such that together they give effective relief.

- 5                    Suitable anti-obesity agents of use in combination with a compound of the present invention, include, but are not limited to: 1) growth hormone secretagogues, such as those disclosed and specifically described in U.S. Patent 5,536,716; 2) growth hormone secretagogue receptor agonists/antagonists, such as NN703, hexarelin, MK-0677, SM-130686, CP-424,391, L-692,429 and L-163,255, 10 and such as those disclosed in U.S. Patent No. 6,358,951, U.S. Patent Application Nos. 2002/049196 and 2002/022637, and PCT Application Nos. WO 01/56592 and WO 02/32888; 3) melanocortin agonists, such as Melanotan II or those described in WO 99/64002 and WO 00/74679; 4) Mc4r (melanocortin 4 receptor) agonists, such as CHIR86036 (Chiron), ME-10142, and ME-10145 (Melacure), and those disclosed in 15 PCT Application Nos. WO 01/991752, WO 01/74844, WO 02/12166, WO 02/11715, and WO 02/12178; 5)  $\beta$ -3 agonists, such as AD9677/TAK677 (Dainippon/Takeda), CL-316,243, SB 418790, BRL-37344, L-796568, BMS-196085, BRL-35135A, CGP12177A, BTA-243, Trecadrine, Zeneca D7114, SR 59119A, and such as those disclosed in U.S. Patent Application Nos. 5,705,515, and US 5,451,677 and PCT 20 Patent Publications WO94/18161, WO95/29159, WO97/46556, WO98/04526 and WO98/32753, WO 01/74782, and WO 02/32897; 6) 5HT-2 agonists; 7) 5HT2C (serotonin receptor 2C) agonists, such as BVT933, DPCA37215, WAY161503, R-1065, and those disclosed in U.S. Patent No. 3,914,250, and PCT Application Nos. WO 02/36596, WO 02/48124, WO 02/10169, WO 01/66548, WO 02/44152, WO 25 02/51844, WO 02/40456, and WO 02/40457; 8) orexin antagonists, such as SB-334867-A, and those disclosed in PCT Patent Application Nos. WO 01/96302, WO 01/68609, WO 02/51232, WO 02/51838 and WO 02/090355; 9) melanin concentrating hormone antagonists; 10) melanin-concentrating hormone 1 receptor (MCH1R) antagonists, such as T-226296 (Takeda), and those disclosed in PCT Patent 30 Application Nos. WO 01/82925, WO 01/87834, WO 02/06245, WO 02/04433, WO 02/51809 and WO 02/083134, and Japanese Patent Application No. JP 13226269; 11) melanin-concentrating hormone 2 receptor (MCH2R) agonist/antagonists; 12) galanin antagonists; 13) CCK agonists; 14) CCK-A (cholecystokinin -A) agonists, such as AR-R 15849, GI 181771, JMV-180, A-71378, A-71623 and SR146131, and those



disclosed in U.S. Patent No. 5,739,106; 15) GLP-1 agonists; 16) corticotropin-releasing hormone agonists; 17) NPY 5 antagonists, such as GW-569180A, GW-594884A, GW-587081X, GW-548118X, FR226928, FR 240662, FR252384, 1229U91, GI-264879A, CGP71683A, LY-377897, PD-160170, SR-120562A, SR-120819A and JCF-104, and those disclosed in U.S. Patent Nos. 6,140,354, 6,191,160, 6,313,298, 6,337,332, 6,329,395, 6,326,375, 6,335,345, and 6,340,683, European Patent Nos. EP-01010691, and EP-01044970, and PCT Patent Publication Nos. WO 97/19682, WO 97/20820, WO 97/20821, WO 97/20822, WO 97/20823, WO 98/27063, WO 00/64880, WO 00/68197, WO 00/69849, WO 01/09120, WO 01/14376, WO 01/85714, WO 01/85730, WO 01/07409, WO 01/02379, WO 01/02379, WO 01/23388, WO 01/23389, WO 01/44201, WO 01/62737, WO 01/62738, WO 01/09120, WO 02/22592, WO 0248152, and WO 02/49648; 18) NPY 1 antagonists, such as BIBP3226, J-115814, BIBO 3304, LY-357897, CP-671906, GI-264879A, and those disclosed in U.S. Patent No. 6,001,836, and PCT Patent Publication Nos. WO 96/14307, WO 01/23387, WO 99/51600, WO 01/85690, WO 01/85098, WO 01/85173, and WO 01/89528; 19) histamine receptor-3 (H3) modulators; 20) histamine receptor-3 (H3) antagonists/inverse agonists, such as hioperamide, 3-(1H-imidazol-4-yl)propyl N-(4-pentenyl)carbamate, clobenpropit, iodophenpropit, imoproxifan, GT2394 (Gliatech), and those described and disclosed in PCT Application No. WO 02/15905, and O-[3-(1H-imidazol-4-yl)propanol]-carbamates (Kiec-Kononowicz, K. et al., *Pharmazie*, 55:349-55 (2000)), piperidine-containing histamine H3-receptor antagonists (Lazewska, D. et al., *Pharmazie*, 56:927-32 (2001), benzophenone derivatives and related compounds (Sasse, A. et al., *Arch. Pharm.(Weinheim)* 334:45-52 (2001)), substituted N-phenylcarbamates (Reidemeister, S. et al., *Pharmazie*, 55:83-6 (2000)), and proxifan derivatives (Sasse, A. et al., *J. Med. Chem.* 43:3335-43 (2000)); 21)  $\beta$ -hydroxy steroid dehydrogenase-1 inhibitors ( $\beta$ -HSD-1); 22) PDE (phosphodiesterase) inhibitors, such as theophylline, pentoxifylline, zaprinast, sildenafil, amrinone, milrinone, cilostamide, rolipram, and cilomilast; 23) phosphodiesterase-3B (PDE3B) inhibitors; 24) NE (norepinephrine) transport inhibitors, such as GW 320659, despiramine, talsupram, and nomifensine; 25) non-selective serotonin/norepinephrine transport inhibitors, such as sibutramine or fenfluramine; 26) ghrelin antagonists, such as those disclosed in PCT Application Nos. WO.01/87335, and WO 02/08250; 27) leptin, including recombinant human leptin (PEG-OB, Hoffman La Roche) and recombinant methionyl human leptin (Amgen); 28) leptin derivatives, such as those disclosed in U.S. Patent Nos.

5,552,524, 5,552,523, 5,552,522, 5,521,283, and PCT International Publication Nos. WO 96/23513, WO 96/23514, WO 96/23515, WO 96/23516, WO 96/23517, WO 96/23518, WO 96/23519, and WO 96/23520; 29) BRS3 (bombesin receptor subtype 3) agonists; 30) CNTF (Ciliary neurotrophic factors), such as GI-181771 (Glaxo-SmithKline), SR146131 (Sanofi Synthelabo), butabindide, PD170,292, and PD 149164 (Pfizer); 31) CNTF derivatives, such as axokine (Regeneron), and those disclosed in PCT Application Nos. WO 94/09134, WO 98/22128, and WO 99/43813; 32) monoamine reuptake inhibitors, such as those disclosed in PCT Application Nos. WO 01/27068, and WO 01/62341; 33) UCP-1 (uncoupling protein-1), 2, or 3  
 10 activators, such as phytanic acid, 4-[(E)-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1-propenyl]benzoic acid (TTNPB), retinoic acid, and those disclosed in PCT Patent Application No. WO 99/00123; 34) thyroid hormone  $\beta$  agonists, such as KB-2611 (KaroBioBMS), and those disclosed in PCT Application No. WO 02/15845, and Japanese Patent Application No. JP 2000256190; 35) FAS (fatty acid synthase)  
 15 inhibitors, such as Cerulenin and C75; 36) DGAT1 (diacylglycerol acyltransferase 1) inhibitors; 37) DGAT2 (diacylglycerol acyltransferase 2) inhibitors; 38) ACC2 (acetyl-CoA carboxylase-2) inhibitors; 39) glucocorticoid antagonists; 40) acyl-estrogens, such as oleoyl-estrone, disclosed in del Mar-Grasa, M. et al., Obesity Research, 9:202-9 (2001); 41) lipase inhibitors, such as orlistat (Xenical®), Triton  
 20 WR1339, RHC80267, lipstatin, tetrahydrolipstatin, teasaponin, diethylumbelliferyl phosphate, and those disclosed in PCT Application No. WO 01/77094; 42) fatty acid transporter inhibitors; 43) dicarboxylate transporter inhibitors; 44) glucose transporter inhibitors; 45) phosphate transporter inhibitors; 46) serotonin reuptake inhibitors, such as those disclosed in U.S. Patent Application No. 6,365,633, and PCT Patent  
 25 Application Nos. WO 01/27060, and WO 01/162341; 47) Metformin (Glucophage®); and/or 48) Topiramate (Topimax®).

Specific NPY5 antagonists of use in combination with a compound of the present invention are selected from the group consisting of:

- (1) 3-oxo-N-(5-phenyl-2-pyrazinyl)-spiro[isobenzofuran-1(3H),4'-piperidine]-1'-carboxamide,  
 30
- (2) 3-oxo-N-(7-trifluoromethylpyrido[3,2-b]pyridin-2-yl)spiro-[isobenzofuran-1(3H),4'-piperidine]-1'-carboxamide,
- (3) N-[5-(3-fluorophenyl)-2-pyrimidinyl]-3-oxospiro-[isobenzofuran-1(3H),4'-piperidine]-1'-carboxamide,

- (4) trans-3'-oxo-N-(5-phenyl-2-pyrimidinyl)spiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide,
- (5) trans-3'-oxo-N-[1-(3-quinolyl)-4-imidazolyl]spiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide,
- 5 (6) trans-3-oxo-N-(5-phenyl-2-pyrazinyl)spiro[4-azaiso-benzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- (7) trans-N-[5-(3-fluorophenyl)-2-pyrimidinyl]-3-oxospiro[5-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- (8) trans-N-[5-(2-fluorophenyl)-2-pyrimidinyl]-3-oxospiro[5-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- 10 (9) trans-N-[1-(3,5-difluorophenyl)-4-imidazolyl]-3-oxospiro[7-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- (10) trans-3-oxo-N-(1-phenyl-4-pyrazolyl)spiro[4-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- 15 (11) trans-N-[1-(2-fluorophenyl)-3-pyrazolyl]-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- (12) trans-3-oxo-N-(1-phenyl-3-pyrazolyl)spiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- (13) trans-3-oxo-N-(2-phenyl-1,2,3-triazol-4-yl)spiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide,
- 20 and pharmaceutically acceptable salts and esters thereof.

"Obesity" is a condition in which there is an excess of body fat. The operational definition of obesity is based on the Body Mass Index (BMI), which is calculated as body weight per height in meters squared ( $\text{kg/m}^2$ ). "Obesity" refers to a condition whereby an otherwise healthy subject has a Body Mass Index (BMI) greater than or equal to  $30 \text{ kg/m}^2$ , or a condition whereby a subject with at least one co-morbidity has a BMI greater than or equal to  $27 \text{ kg/m}^2$ . An "obese subject" is an otherwise healthy subject with a Body Mass Index (BMI) greater than or equal to  $30 \text{ kg/m}^2$  or a subject with at least one co-morbidity with a BMI greater than or equal to  $27 \text{ kg/m}^2$ . A "subject at risk for obesity" is an otherwise healthy subject with a BMI of  $25 \text{ kg/m}^2$  to less than  $30 \text{ kg/m}^2$  or a subject with at least one co-morbidity with a BMI of  $25 \text{ kg/m}^2$  to less than  $27 \text{ kg/m}^2$ .

The increased risks associated with obesity occur at a lower Body Mass Index (BMI) in Asians. In Asian countries, including Japan, "obesity" refers to a condition whereby a subject with at least one obesity-induced or obesity-related co-

morbidity that requires weight reduction or that would be improved by weight reduction, has a BMI greater than or equal to 25 kg/m<sup>2</sup>. In Asian countries, including Japan, an "obese subject" refers to a subject with at least one obesity-induced or obesity-related co-morbidity that requires weight reduction or that would be improved by weight reduction, with a BMI greater than or equal to 25 kg/m<sup>2</sup>. In Asian countries, a "subject at risk of obesity" is a subject with a BMI of greater than 23 kg/m<sup>2</sup> to less than 25 kg/m<sup>2</sup>.

As used herein, the term "obesity" is meant to encompass all of the above definitions of obesity.

Obesity-induced or obesity-related co-morbidities include, but are not limited to, diabetes, non-insulin dependent diabetes mellitus - type 2, impaired glucose tolerance, impaired fasting glucose, insulin resistance syndrome, dyslipidemia, hypertension, hyperuricacidemia, gout, coronary artery disease, myocardial infarction, angina pectoris, sleep apnea syndrome, Pickwickian syndrome, fatty liver; cerebral infarction, cerebral thrombosis, transient ischemic attack, orthopedic disorders, arthritis deformans, lumbodynia, emmeniopathy, and infertility. In particular, co-morbidities include: hypertension, hyperlipidemia, dyslipidemia, glucose intolerance, cardiovascular disease, sleep apnea, diabetes mellitus, and other obesity-related conditions.

"Treatment" (of obesity and obesity-related disorders) refers to the administration of the compounds of the present invention to reduce or maintain the body weight of an obese subject. One outcome of treatment may be reducing the body weight of an obese subject relative to that subject's body weight immediately before the administration of the compounds of the present invention. Another outcome of treatment may be preventing body weight regain of body weight previously lost as a result of diet, exercise, or pharmacotherapy. Another outcome of treatment may be decreasing the occurrence of and/or the severity of obesity-related diseases. The treatment may suitably result in a reduction in food or calorie intake by the subject, including a reduction in total food intake, or a reduction of intake of specific components of the diet such as carbohydrates or fats; and/or the inhibition of nutrient absorption; and/or the inhibition of the reduction of metabolic rate; and in weight reduction in patients in need thereof. The treatment may also result in an alteration of metabolic rate, such as an increase in metabolic rate, rather than or in addition to an inhibition of the reduction of metabolic rate; and/or in minimization of the metabolic resistance that normally results from weight loss.

“Prevention” (of obesity and obesity-related disorders) refers to the administration of the compounds of the present invention to reduce or maintain the body weight of a subject at risk of obesity. One outcome of prevention may be reducing the body weight of a subject at risk of obesity relative to that subject’s body weight immediately before the administration of the compounds of the present invention. Another outcome of prevention may be preventing body weight regain of body weight previously lost as a result of diet, exercise, or pharmacotherapy. Another outcome of prevention may be preventing obesity from occurring if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Another outcome of prevention may be decreasing the occurrence and/or severity of obesity-related disorders if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Moreover, if treatment is commenced in already obese subjects, such treatment may prevent the occurrence, progression or severity of obesity-related disorders, such as, but not limited to, arteriosclerosis, Type II diabetes, polycystic ovarian disease, cardiovascular diseases, osteoarthritis, dermatological disorders, hypertension, insulin resistance, hypercholesterolemia, hypertriglyceridemia, and cholelithiasis.

The obesity-related disorders herein are associated with, caused by, or result from obesity. Examples of obesity-related disorders include overeating and bulimia, hypertension, diabetes, elevated plasma insulin concentrations and insulin resistance, dyslipidemias, hyperlipidemia, endometrial, breast, prostate and colon cancer, osteoarthritis, obstructive sleep apnea, cholelithiasis, gallstones, heart disease, abnormal heart rhythms and arrhythmias, myocardial infarction, congestive heart failure, coronary heart disease, sudden death, stroke, polycystic ovarian disease, craniopharyngioma, the Prader-Willi Syndrome, Frohlich’s syndrome, GH-deficient subjects, normal variant short stature, Turner’s syndrome, and other pathological conditions showing reduced metabolic activity or a decrease in resting energy expenditure as a percentage of total fat-free mass, e.g, children with acute lymphoblastic leukemia. Further examples of obesity-related disorders are metabolic syndrome, also known as syndrome X, insulin resistance syndrome, sexual and reproductive dysfunction, such as infertility, hypogonadism in males and hirsutism in females, gastrointestinal motility disorders, such as obesity-related gastro-esophageal reflux, respiratory disorders, such as obesity-hypoventilation syndrome (Pickwickian syndrome), cardiovascular disorders, inflammation, such as systemic inflammation of the vasculature, arteriosclerosis, hypercholesterolemia, hyperuricaemia, lower back

pain, gallbladder disease, gout, and kidney cancer. The compounds of the present invention are also useful for reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy.

The term "diabetes," as used herein, includes both insulin-  
5 dependent diabetes mellitus (i.e., IDDM, also known as type I diabetes) and non-insulin-dependent diabetes mellitus (i.e., NIDDM, also known as Type II diabetes. Type I diabetes, or insulin-dependent diabetes, is the result of an absolute deficiency of insulin, the hormone which regulates glucose utilization. Type II  
10 diabetes, or insulin-independent diabetes (i.e., non-insulin-dependent diabetes mellitus), often occurs in the face of normal, or even elevated levels of insulin and appears to be the result of the inability of tissues to respond appropriately to insulin. Most of the Type II diabetics are also obese. The compounds of the present invention are useful for treating both Type I and Type II diabetes. The compounds are especially effective for treating Type II diabetes. The compounds  
15 of the present invention are also useful for treating and/or preventing gestational diabetes mellitus.

It will be appreciated that for the treatment or prevention of migraine, a compound of the present invention may be used in conjunction with other anti-migraine agents, such as ergotamines or 5-HT<sub>1</sub> agonists, especially sumatriptan,  
20 naratriptan, zolmatriptan or rizatriptan.

It will be appreciated that for the treatment of depression or anxiety, a compound of the present invention may be used in conjunction with other anti-depressant or anti-anxiety agents.

Suitable classes of anti-depressant agents include norepinephrine  
25 reuptake inhibitors, selective serotonin reuptake inhibitors (SSRIs), monoamine oxidase inhibitors (MAOIs), reversible inhibitors of monoamine oxidase (RIMAs), serotonin and noradrenaline reuptake inhibitors (SNRIs), corticotropin releasing factor (CRF) antagonists,  $\alpha$ -adrenoreceptor antagonists, neurokinin-1 receptor antagonists and atypical anti-depressants.

30 Suitable norepinephrine reuptake inhibitors include tertiary amine tricyclics and secondary amine tricyclics. Suitable examples of tertiary amine tricyclics include: amitriptyline, clomipramine, doxepin, imipramine and trimipramine, and pharmaceutically acceptable salts thereof. Suitable examples of secondary amine tricyclics include: amoxapine, desipramine, maprotiline,  
35 nortriptyline and protriptyline, and pharmaceutically acceptable salts thereof.

Suitable selective serotonin reuptake inhibitors include: fluoxetine, fluvoxamine, paroxetine, imipramine and sertraline, and pharmaceutically acceptable salts thereof.

5 Suitable monoamine oxidase inhibitors include: isocarboxazid, phenelzine, tranylcypromine and selegiline, and pharmaceutically acceptable salts thereof.

Suitable reversible inhibitors of monoamine oxidase include: moclobemide, and pharmaceutically acceptable salts thereof.

10 Suitable serotonin and noradrenaline reuptake inhibitors of use in the present invention include: venlafaxine, and pharmaceutically acceptable salts thereof.

Suitable CRF antagonists include those compounds described in International Patent Specification Nos. WO 94/13643, WO 94/13644, WO 94/13661, WO 94/13676 and WO 94/13677.

15 Suitable neurokinin-1 receptor antagonists may be peptidal or non-peptidal in nature, however, the use of a non-peptidal neurokinin-1 receptor antagonist is preferred. In a preferred embodiment, the neurokinin-1 receptor antagonist is a CNS-penetrant neurokinin-1 receptor antagonist. In addition, for convenience the use of an orally active neurokinin-1 receptor antagonist is preferred. To facilitate dosing, it is also preferred that the neurokinin-1 receptor antagonist is a  
20 long acting neurokinin-1 receptor antagonist. An especially preferred class of neurokinin-1 receptor antagonists of use in the present invention are those compounds which are orally active and long acting.

Neurokinin-1 receptor antagonists of use in the present invention are fully described, for example, in U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930,  
25 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0 436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275, 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0  
30 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535, 0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and 0 776 893; PCT International Patent Publication Nos. WO 90/05525, 90/05729; 91/09844, 91/18899, 92/01688, 92/06079, 92/12151, 92/15585, 92/17449, 92/20661, 92/20676, 92/21677, 92/22569, 93/00330, 93/00331, 93/01159, 93/01165,

93/01169, 93/01170, 93/06099, 93/09116, 93/10073, 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440, 94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843, 94/08997, 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767, 5 94/15903, 94/19320, 94/19323, 94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040, 95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679, 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338, 95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 96/05193, 96/05203, 96/06094, 96/07649, 96/10562, 96/16939, 96/18643, 96/20197, 10 96/21661, 96/29304, 96/29317, 96/29326, 96/29328, 96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362, 97/18206, 97/19084, 97/19942, 97/21702, and 97/49710, 98/24438-98/24441, 98/24442-98/24445, 02/16343, and 02/16344; and in British Patent Publication Nos. 2 266 529, 2 268 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689.

15 Specific neurokinin-1 receptor antagonists of use in the present invention include:

- (±)-(2R3R,2S3S)-N-([2-cyclopropoxy-5-(trifluoromethoxy)-phenyl]methyl)-2-phenylpiperidin-3-amine;
- 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3(S)-(4-fluorophenyl)-4-(3-20 (5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenyl-morpholine;
- 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenyl-morpholine;
- 25 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino)methyl-1,2,3-triazol-4-yl)methyl-3-(S)-phenylmorpholine;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-30 dimethylamino)methyl-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine;
- (3S,5R,6S)-3-[2-cyclopropoxy-5-(trifluoromethoxy)phenyl]-6-phenyl-1-oxa-7-aza-spiro[4.5]decane;
- (3R,5R,6S)-3-[2-cyclopropoxy-5-(trifluoromethoxy)phenyl]-6-phenyl-1-oxa-7-aza-spiro[4.5]decane;



2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine;

2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(4-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;

5 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;

2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(2-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;

10 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxophosphoryl-1H-1,2,4-triazolo)methyl)morpholine;

2-(S)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1-monophosphoryl-5-oxo-4H-1,2,4-triazolo)methyl)morpholine;

2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(4-N,N-dimethylaminobut-2-yn-yl)-3-(S)-(4-fluorophenyl)morpholine;

15 or a pharmaceutically acceptable salt thereof.

Suitable atypical anti-depressants include: bupropion, lithium, nefazodone, trazodone and viloxazine, and pharmaceutically acceptable salts thereof.

Suitable classes of anti-anxiety agents include benzodiazepines and 5-HT<sub>1A</sub> agonists or antagonists, especially 5-HT<sub>1A</sub> partial agonists, and corticotropin releasing factor (CRF) antagonists.

Suitable benzodiazepines include: alprazolam, chlordiazepoxide, clonazepam, chlorazepate, diazepam, halazepam, lorazepam, oxazepam and prazepam, and pharmaceutically acceptable salts thereof.

25 Suitable 5-HT<sub>1A</sub> receptor agonists or antagonists include, in particular, the 5-HT<sub>1A</sub> receptor partial agonists buspirone, flesinoxan, gepirone and ipsapirone, and pharmaceutically acceptable salts thereof.

Suitable corticotropin releasing factor (CRF) antagonists include those previously discussed herein.

30 As used herein, the term "substance abuse disorders" includes substance dependence or abuse with or without physiological dependence. The substances associated with these disorders are: alcohol, amphetamines (or amphetamine-like substances), caffeine, cannabis, cocaine, hallucinogens, inhalants, marijuana, nicotine, opioids, phencyclidine (or phencyclidine-like compounds), sedative-hypnotics or benzodiazepines, and other (or unknown) substances and  
35 combinations of all of the above.

In particular, the term "substance abuse disorders" includes drug withdrawal disorders such as alcohol withdrawal with or without perceptual disturbances; alcohol withdrawal delirium; amphetamine withdrawal; cocaine withdrawal; nicotine withdrawal; opioid withdrawal; sedative, hypnotic or anxiolytic withdrawal with or without perceptual disturbances; sedative, hypnotic or anxiolytic withdrawal delirium; and withdrawal symptoms due to other substances. It will be appreciated that reference to treatment of nicotine withdrawal includes the treatment of symptoms associated with smoking cessation.

Other "substance abuse disorders" include substance-induced anxiety disorder with onset during withdrawal; substance-induced mood disorder with onset during withdrawal; and substance-induced sleep disorder with onset during withdrawal.

It will be appreciated that a combination of a conventional antipsychotic drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of mania. Such a combination would be expected to provide for a rapid onset of action to treat a manic episode thereby enabling prescription on an "as needed basis". Furthermore, such a combination may enable a lower dose of the antipsychotic agent to be used without compromising the efficacy of the antipsychotic agent, thereby minimizing the risk of adverse side-effects. A yet further advantage of such a combination is that, due to the action of the CB1 receptor modulator, adverse side-effects caused by the antipsychotic agent such as acute dystonias, dyskinesias, akathisia and tremor may be reduced or prevented.

Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and an antipsychotic agent for the manufacture of a medicament for the treatment or prevention of mania.

The present invention also provides a method for the treatment or prevention of mania, which method comprises administration to a patient in need of such treatment or at risk of developing mania of an amount of a CB1 receptor modulator and an amount of an antipsychotic agent, such that together they give effective relief.

In a further aspect of the present invention, there is provided a pharmaceutical composition comprising a CB1 receptor modulator and an antipsychotic agent, together with at least one pharmaceutically acceptable carrier or excipient.

It will be appreciated that the CB1 receptor modulator and the antipsychotic agent may be present as a combined preparation for simultaneous, separate or sequential use for the treatment or prevention of mania. Such combined preparations may be, for example, in the form of a twin pack.

5           In a further or alternative aspect of the present invention, there is therefore provided a product comprising a CB1 receptor modulator and an antipsychotic agent as a combined preparation for simultaneous, separate or sequential use in the treatment or prevention of mania.

10           It will be appreciated that when using a combination of the present invention, the CB1 receptor modulator and the antipsychotic agent may be in the same pharmaceutically acceptable carrier and therefore administered simultaneously. They may be in separate pharmaceutical carriers such as conventional oral dosage forms which are taken simultaneously. The term "combination" also refers to the case where the compounds are provided in separate dosage forms and are administered  
15 sequentially. Therefore, by way of example, the antipsychotic agent may be administered as a tablet and then, within a reasonable period of time, the CB1 receptor modulator may be administered either as an oral dosage form such as a tablet or a fast-dissolving oral dosage form. By a "fast-dissolving oral formulation" is meant, an oral delivery form which when placed on the tongue of a patient, dissolves within  
20 about 10 seconds.

Included within the scope of the present invention is the use of CB1 receptor modulators in combination with an antipsychotic agent in the treatment or prevention of hypomania.

25           It will be appreciated that a combination of a conventional antipsychotic drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of schizophrenic disorders. Such a combination would be expected to provide for a rapid onset of action to treat schizophrenic symptoms thereby enabling prescription on an "as needed basis". Furthermore, such a combination may enable a lower dose of the CNS agent to be used without compromising the efficacy of the  
30 antipsychotic agent, thereby minimizing the risk of adverse side-effects. A yet further advantage of such a combination is that, due to the action of the CB1 receptor modulator, adverse side-effects caused by the antipsychotic agent such as acute dystonias, dyskinesias, akathisia and tremor may be reduced or prevented.

As used herein, the term "schizophrenic disorders" includes paranoid, disorganized, catatonic, undifferentiated and residual schizophrenia; schizophreniform disorder; schizoaffective disorder; delusional disorder; brief psychotic disorder; shared psychotic disorder; substance-induced psychotic disorder; and psychotic disorder not otherwise specified.

Other conditions commonly associated with schizophrenic disorders include self-injurious behavior (e.g. Lesch-Nyhan syndrome) and suicidal gestures.

Suitable antipsychotic agents of use in combination with a CB1 receptor modulator include the phenothiazine, thioxanthene, heterocyclic dibenzazepine, butyrophenone, diphenylbutylpiperidine and indolone classes of antipsychotic agent. Suitable examples of phenothiazines include chlorpromazine, mesoridazine, thioridazine, acetophenazine, fluphenazine, perphenazine and trifluoperazine. Suitable examples of thioxanthenes include chlorprothixene and thiothixene. Suitable examples of dibenzazepines include clozapine and olanzapine. An example of a butyrophenone is haloperidol. An example of a diphenylbutylpiperidine is pimozide. An example of an indolone is molindolone. Other antipsychotic agents include loxapine, sulpiride and risperidone. It will be appreciated that the antipsychotic agents when used in combination with a CB1 receptor modulator may be in the form of a pharmaceutically acceptable salt, for example, chlorpromazine hydrochloride, mesoridazine besylate, thioridazine hydrochloride, acetophenazine maleate, fluphenazine hydrochloride, flurphenazine enathate, fluphenazine decanoate, trifluoperazine hydrochloride, thiothixene hydrochloride, haloperidol decanoate, loxapine succinate and molindone hydrochloride. Perphenazine, chlorprothixene, clozapine, olanzapine, haloperidol, pimozide and risperidone are commonly used in a non-salt form.

Other classes of antipsychotic agent of use in combination with a CB1 receptor modulator include dopamine receptor antagonists, especially D2, D3 and D4 dopamine receptor antagonists, and muscarinic m1 receptor agonists. An example of a D3 dopamine receptor antagonist is the compound PNU-99194A. An example of a D4 dopamine receptor antagonist is PNU-101387. An example of a muscarinic m1 receptor agonist is xanomeline.

Another class of antipsychotic agent of use in combination with a CB1 receptor modulator is the 5-HT<sub>2A</sub> receptor antagonists, examples of which include MDL100907 and fananserin. Also of use in combination with a CB1 receptor

modulator are the serotonin dopamine antagonists (SDAs) which are believed to combine 5-HT<sub>2A</sub> and dopamine receptor antagonist activity, examples of which include olanzapine and ziperastidone.

Still further, NK-1 receptor antagonists may be favorably employed with the CB1 receptor modulators of the present invention. Preferred NK-1 receptor antagonists for use in the present invention are selected from the classes of compounds described in European Patent Specification No. 0 577 394, and International Patent Specification Nos. 95/08549, 95/18124, 95/23798, 96/05181, and 98/49710 (Application No. PCT/GB97/01630). The preparation of such compounds is fully described in the aforementioned publications.

Particularly preferred NK-1 receptor antagonists of use in the present invention include: (3S,5R,6S)-3-[2-cyclopropoxy-5-(trifluoromethoxy)phenyl]-6-phenyl-1-oxa-7-aza-spiro[4.5]decane;  
 (3R,5R,6S)-3-[2-cyclopropoxy-5-(trifluoromethoxy)phenyl]-6-phenyl-1-oxa-7-aza-spiro[4.5]decane;  
 (±)-(2R,3R,2S,3S)-N- {[2-cyclopropoxy-5-(trifluoromethoxy)phenyl]methyl}-2-phenylpiperidin-3-amine;  
 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenyl-morpholine;  
 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenyl-morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino)methyl-1,2,3-triazol-4-yl)methyl-3-(S)-phenylmorpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino)methyl-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(4-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;

- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(2-monophosphoryl-5-oxo-1H-1,2,4-triazolo)methyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxyphosphoryl-1H-1,2,4-triazolo)methyl)morpholine;  
 5 2-(S)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1-monophosphoryl-5-oxo-4H-1,2,4-triazolo)methyl)morpholine;  
 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(4-N,N-dimethylaminobut-2-yn-yl)-3-(S)-(4-fluorophenyl)morpholine;  
 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine;  
 10 or a pharmaceutically acceptable salt thereof.

It will be appreciated that a combination of a conventional anti-asthmatic drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of asthma.

- 15 Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and an anti-asthmatic agent for the manufacture of a medicament for the treatment or prevention of asthma.

- The present invention also provides a method for the treatment or prevention of asthma, which method comprises administration to a patient in need of  
 20 such treatment an amount of a compound of the present invention and an amount of an anti-asthmatic agent, such that together they give effective relief.

- Suitable anti-asthmatic agents of use in combination with a compound of the present invention include, but are not limited to: (a) VLA-4 antagonists such as natalizumab and the compounds described in US 5,510,332, WO97/03094,  
 25 WO97/02289, WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973 and WO96/31206; (b) steroids and corticosteroids such as beclomethasone, methylprednisolone, betamethasone, prednisone, dexamethasone, and hydrocortisone; (c) antihistamines (H1-histamine antagonists) such as bromopheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine,  
 30 diphenhydramine, diphenylpyraline, tripeleminamine, hydroxyzine, methdilazine, promethazine, trimeprazine, azatadine, cyproheptadine, antazoline, pheniramine pyrilamine, astemizole, terfenadine, loratadine, desloratadine, cetirizine, fexofenadine, descarboethoxyloratadine, and the like; (d) non-steroidal anti-asthmatics including  $\beta$ 2-agonists (such as terbutaline, metaproterenol, fenoterol,

isoetharine, albuterol, bitolterol, salmeterol, epinephrine, and pirbuterol), theophylline, cromolyn sodium, atropine, ipratropium bromide, leukotriene antagonists (such as zafirlukast, montelukast, pranlukast, iralukast, pobilukast, and SKB-106,203), and leukotriene biosynthesis inhibitors (such as zileuton and BAY-1005); (e) anti-cholinergic agents including muscarinic antagonists (such as ipratropium bromide and atropine); (f) antagonists of the chemokine receptors, especially CCR-1, CCR-2, and CCR-3; (g) immunosuppressants such as cyclosporin, tacrolimus, rapamycin and other FK-506 type immunosuppressants; (h) non-steroidal antiinflammatory agents (NSAIDs) such as propionic acid derivatives (alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, miroprofen, naproxen, oxaprozin, piroprofen, pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid derivatives (indomethacin, acemetacin, alclofenac, clidanaç, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac), fenamic acid derivatives (flufenamic acid, meclofenamic acid, mefenamic acid, niflumic acid and tolfenamic acid), biphenylcarboxylic acid derivatives (diflunisal and flufenisal), oxicams (isoxicam, piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid, sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone, phenylbutazone); (i) cyclooxygenase-2 (COX-2) inhibitors such as celecoxib; (j) anti-diabetic agents such as insulin, sulfonylureas, biguanides (metformin),  $\alpha$ -glucosidase inhibitors (acarbose) and glitazones (troglitazone, pioglitazone, englitazone, MCC-555, BRL49653 and the like); (k) preparations of interferon beta (interferon beta-1a, interferon beta-1b); (l) other compounds such as 5-aminosalicylic acid and prodrugs thereof, and pharmaceutically acceptable salts thereof.

It will be appreciated that a combination of a conventional anti-constipation drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of constipation.

Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and an anti-constipation agent for the manufacture of a medicament for the treatment or prevention of constipation.

The present invention also provides a method for the treatment or prevention of constipation, which method comprises administration to a patient in

need of such treatment an amount of a compound of the present invention and an amount of an anti-constipation agent, such that together they give effective relief.

It will be appreciated that a combination of a conventional anti-constipation drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of chronic intestinal pseudo-obstruction.

Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and an anti-constipation agent for the manufacture of a medicament for the treatment or prevention of chronic intestinal pseudo-obstruction.

The present invention also provides a method for the treatment or prevention of chronic intestinal pseudo-obstruction, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of an anti-constipation agent, such that together they give effective relief.

Suitable anti-constipation agents of use in combination with a compound of the present invention include, but are not limited to, osmotic agents, laxatives and detergent laxatives (or wetting agents), bulking agents, and stimulants; and pharmaceutically acceptable salts thereof.

A particularly suitable class of osmotic agents include, but are not limited to sorbitol, lactulose, polyethylene glycol, magnesium, phosphate, and sulfate; and pharmaceutically acceptable salts thereof.

A particularly suitable class of laxatives and detergent laxatives, include, but are not limited to, magnesium, and docusate sodium; and pharmaceutically acceptable salts thereof.

A particularly suitable class of bulking agents include, but are not limited to, psyllium, methylcellulose, and calcium polycarbophil; and pharmaceutically acceptable salts thereof.

A particularly suitable class of stimulants include, but are not limited to, anthroquinones, and phenolphthalein; and pharmaceutically acceptable salts thereof.

It will be appreciated that a combination of a conventional anti-cirrhosis drug with a CB1 receptor modulator may provide an enhanced effect in the treatment of cirrhosis of the liver.



Thus, according to a further aspect of the present invention there is provided the use of a CB1 receptor modulator and an anti-cirrhosis agent for the manufacture of a medicament for the treatment or prevention of cirrhosis of the liver.

5 The present invention also provides a method for the treatment or prevention of cirrhosis of the liver, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an anti-cirrhosis agent, such that together they give effective relief.

Suitable anti-cirrhosis agents of use in combination with a compound of the present invention include, but are not limited to, corticosteroids, penicillamine, 10 colchicine, interferon- $\gamma$ , 2-oxoglutarate analogs, prostaglandin analogs, and other anti-inflammatory drugs and antimetabolites such as azathioprine, methotrexate, leflunamide, indomethacin, naproxen, and 6-mercaptopurine; and pharmaceutically acceptable salts thereof.

15 The method of treatment of this invention comprises a method of modulating the CB1 receptor and treating CB1 receptor mediated diseases by administering to a patient in need of such treatment a non-toxic therapeutically effective amount of a compound of this invention that selectively antagonizes the CB1 receptor in preference to the other CB or G-protein coupled receptors.

The term "therapeutically effective amount" means the amount the 20 compound of structural formula I that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disorder being treated. The novel methods of treatment of this invention are for disorders known to those skilled in the art. The term "mammal" includes humans.


25 The weight ratio of the compound of the Formula I to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the Formula I is combined with a  $\beta$ -3 agonist the weight ratio of the compound of the Formula I to the  $\beta$ -3 agonist will generally range from about 1000:1 30 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the Formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

## Abbreviations used in the following Schemes and Examples:

	aq.:	aqueous
	API-ES:	atmospheric pressured ionization-electrospray (mass spectrum term)
	BOC:	tert-butoxycarbonyl
5	brine:	saturated sodium chloride solution
	CDCl <sub>3</sub> :	chloroform, deuterated
	Celite:	CELITE brand diatomaceous earth
	CH <sub>2</sub> Cl <sub>2</sub> :	dichloromethane
	DMAP:	4-dimethylaminopyridine
10	DMSO:	dimethylsulfoxide
	EDC:	1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride
	Et:	ethyl
	Et <sub>2</sub> O:	diethyl ether
	EtOAc:	ethyl acetate
15	g or gm:	gram
	h or hr:	hours
	HOBt:	1-hydroxybenzotriazole
	HPLC:	high pressure liquid chromatography
	HPLC-MS:	high pressure liquid chromatography-Mass Spectroscopy
20	LC-MS	liquid chromatography-mass spectrum
	MeOH:	methanol
	mg:	milligram
	MHz:	megahertz
	min:	minutes
25	mL:	milliliter
	mmol:	millimole
	MS or ms:	mass spectrum
	rt:	room temperature
	TFA:	trifluoroacetic acid
30	TLC:	thin layer chromatography

Compounds of the present invention may be prepared by procedures illustrated in the accompanying schemes.

Scheme 1.




In Scheme 1, an appropriately substituted amine **A** is reacted with an  
 5 aryl carboxylic acid **B** under standard amide bond forming conditions to afford the  
 amide **C**.

In order to illustrate the invention, the following examples are  
 included. These examples do not limit the invention. They are only meant to suggest  
 a method of reducing the invention to practice. Those skilled in the art may find other  
 10 methods of practicing the invention which are readily apparent to them. However,  
 those methods are also deemed to be within the scope of this invention.

#### General Procedures.

The LC/MS analyses were preformed using a Micromass ZMD mass  
 15 spectrometer coupled to an Agilent 1100 Series HPLC utilizing a YMC ODS-A 4.6 x  
 50 mm column, eluting at 2.5 mL/min, with a solvent gradient of 10 to 95% B over  
 4.5 min, followed by 0.5 min at 95% B: solvent A = 0.06% TFA in water; solvent B =  
 0.05% TFA in acetonitrile. <sup>1</sup>H-NMR spectra were obtained on a 400 or 500 MHz  
 Varian Spectrometer in CDCl<sub>3</sub> or CD<sub>3</sub>OD as indicated and chemical shifts are  
 20 reported as  $\delta$  using the solvent peak as reference and coupling constants are reported  
 in hertz (Hz).


#### REFERENCE EXAMPLE 1



N-[2,3-Bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride

The preparation of the two diastereomers (alpha and beta) of N-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride salt has been disclosed (Schultz, E.M, *et al. J. Med Chem.* **1967**, *10*, 717). Diastereomer  $\alpha$ : LC-MS: calculated for  $C_{16}H_{17}Cl_2N$  293, observed m/e 294 ( $M + H$ )<sup>+</sup> (retention time 2.5 min). Diastereomer  $\beta$ : LC-MS: calculated for  $C_{16}H_{17}Cl_2N$  293, observed m/e 294 ( $M + H$ )<sup>+</sup> (retention time 2.2 min).

10

REFERENCE EXAMPLE 2N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-amine hydrochloride15 (Diastereomer  $\alpha$ )Step A 3-(4-Chlorophenyl)-2-phenylpropanoic acid, methyl ester

To a solution of methyl phenylacetate (12 g, 80 mmol) and 4-chlorobenzyl bromide (16 g, 80 mmol) in 250 mL of anhydrous tetrahydrofuran at -78°C was added sodium hexamethyldisilazide (1 M in tetrahydrofuran, 80 mL, 80 mmol) (potassium hexamethyldisilazide in toluene may be used with similar results). The reaction was allowed to warm to room temperature overnight. The volatile materials were removed on a rotary evaporator, and the resulting mixture was partitioned between saturated ammonium chloride (200 mL) and ethyl acetate (200 mL). The organic layer was separated and the aqueous layer extracted with ethyl acetate (2 x 200 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.36-7.10 (m, 9H), 3.81 (dd, 1H), 3.52 (s, 3H), 3.36 (dd, 1H), 3.02 (dd, 1H).

Step B 3-(4-Chlorophenyl)-2-phenylpropanoic acid

To a mixture of methyl 3-(4-chlorophenyl)-2-phenylpropionate (Step A, 20 g, 74 mmol) in acetonitrile (100 mL) and water (100 mL) was added lithium hydroxide monohydrate (8.8 g, 0.21 mol). After stirring at room temperature for 3 days, the volatile materials were removed by concentrating on a rotary evaporator and the residue was partitioned between water (300 mL) and hexane/ether (1:1, 200 mL). The water layer was separated, acidified to pH = 2-3, and extracted with ethyl acetate (2 x 200 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.34-7.10 (m, 9H), 3.82 (dd, 1H), 3.36 (dd, 1H), 2.98 (dd, 1H).

10 Step C *N*-Methoxy-*N*-methyl-3-(4-chlorophenyl)-2-phenylpropanamide

To a solution of 3-(4-chlorophenyl)-2-phenylpropionic acid (Step B, 14 g, 55 mmol) in methylene chloride (125 mL) at 0°C were added dropwise dimethylformamide (50 μL) and oxalyl chloride (14 g, 0.11 mol). The reaction was allowed to warm to room temperature overnight and concentrated to dryness to give the crude acyl chloride, which was used without further purification. To a solution of the crude acyl chloride in methylene chloride (250 mL) was added *N*-methoxy-*N*-methylamine hydrochloride (11 g, 0.11 mol) and triethyl amine (dried over activated molecular sieves, 30 mL, 0.22 mol) at 0°C. After stirring at room temperature for 4 h, the reaction mixture was diluted with ether (500 mL) and successively washed with water, dilute aqueous sodium hydrogen sulfate and brine, dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the crude product, which was used without further purification. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.4-7.1 (m, 9H), 4.38 (br, 1H), 3.48 (s, 3H), 3.35 (dd, 1H), 3.10 (s, 3H), 2.92 (dd, 1H); LC-MS: m/e 304 (3.6 min).

25 Step D 4-(4-Chlorophenyl)-3-phenyl-2-butanone

To a solution of *N*-methoxy-*N*-methyl-3-(4-chlorophenyl)-2-phenylpropanamide (Step C, 16 g, 53 mmol, dried by azeotroping with toluene) in anhydrous tetrahydrofuran (200 mL) at 0°C was added methylmagnesium bromide (3 M in ether, 35 mL, 0.11 mol). After stirring at 0°C for 2 h, the reaction was quenched with methanol (5 mL) and 2 M hydrochloric acid (50 mL). The volatile materials were removed by concentrating on a rotary evaporator and the residue partitioned between saturated ammonium chloride (200 mL) and ether (200 mL). The organic layer was separated, and the aqueous layer was extracted with ether (2 times 200 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the title compound, which was used without further

purification.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.45-7.02 (m, 9H), 4.08 (dd, 1H), 3.34 (dd, 1H), 2.90 (dd, 1H), 2.03 (s, 3H).

Step E 4-(4-Chlorophenyl)-3-phenyl-2-butanol

To a solution of 4-(4-chlorophenyl)-3-phenyl-2-butanone (Step D, 13 g, 50 mmol) in methanol (100 mL) at  $0^\circ\text{C}$  was added sodium borohydride (3.8 g, 100 mmol). After stirring at  $0^\circ\text{C}$  for 30 min, the reaction was quenched by adding 2 M hydrochloric acid (50 mL). The volatile materials were removed by concentrating on a rotary evaporator and the residue partitioned between water (100 mL) and ethyl acetate (200 mL). The organic layer was separated and the aqueous layer extracted with ethyl acetate (2 x 200 mL). The combined organic extracts were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated to dryness to give the crude product, which was purified by flash column chromatography on silica gel eluted with 10% ethyl acetate in hexane to afford the pure faster eluting isomer and a mixture containing both the faster eluting isomer and the slower eluting isomer.

15 Faster eluting isomer:  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.25-7.00 (m, 9H), 4.00 (m, 1H), 3.15 (m, 1H), 2.97 (m, 1H), 2.85 (m, 1H), 1.10 (d, 3H).

Step F 4-(4-Chlorophenyl)-2-methanesulfonyloxy-3-phenylbutane

To a solution of 4-(4-chlorophenyl)-3-phenyl-2-butanol (Step E, faster eluting isomer, 9.0 g, 34 mmol) in ethyl acetate (100 mL) at  $0^\circ\text{C}$  was added triethyl amine (dried over activated molecular sieves, 5.8 mL, 42 mmol) and methanesulfonyl chloride (3.0 mL, 38 mmol). After stirring at  $0^\circ\text{C}$  for 30 min, the reaction was quenched by addition of saturated aqueous sodium bicarbonate (100 mL). After stirring at room temperature for 1 h, the organic layer was separated, dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give the title compound, which was used without further purification.

25  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.3-7.0 (m, 9H), 5.05 (m, 1H), 3.2-3.0 (m, 3H), 2.80 (s, 3H), 1.40 (d, 3H).

Step G 2-Azido-4-(4-chlorophenyl)-3-phenylbutane

To a solution of 4-(4-chlorophenyl)-2-methanesulfonyloxy-3-phenylbutane (Step F, 12 g, 34 mmol) in dimethylformamide (50 mL) was added sodium azide (11 g, 0.17 mol). After stirring at  $120^\circ\text{C}$  for 1 h, the reaction mixture was poured into water (200 mL), and the product was extracted with ether (2 times 100 mL). The combined organic extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to dryness, and the residue was purified on a silica gel column eluting with hexane to give the title compound.

30

Step H 2-(*N*-*tert*-Butoxycarbonyl)amino-4-(4-chlorophenyl)-3-phenylbutane


To a solution of 2-azido-4-(4-chlorophenyl)-3-phenylbutane (Step G, 7.0 g, 24 mmol) in ethyl acetate (150 mL) was added di(*tert*-butyl) dicarbonate (8.0 g, 37 mmol) and platinum dioxide (0.50 g, 2.2 mmol). The mixture was degassed and filled with hydrogen with a balloon. After stirring for 1 day, the reaction mixture was filtered through Celite, and the filtrate was concentrated to give the crude product. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.25-6.88 (m, 9H), 3.89 (m, 1H), 3.20 (m, 1H), 2.86-2.77 (m, 2H), 1.54 (s, 9H), 0.92 (d, 3H).

Step I *N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-amine hydrochloride

(Diastereomer α).

2-(*N*-*tert*-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-phenylbutane (Step H, 7.0 g, 24 mmol) was treated with a saturated solution of hydrogen chloride in ethyl acetate (100 mL) at room temperature for 30 min (4 M hydrogen chloride in dioxane may be used with similar results). The mixture was concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.35-6.98 (m, 9H), 3.62 (m, 1H), 3.20 (dd, 1H), 3.05 (m, 1H), 2.98 (dd, 1H), 1.19 (d, 3H). LC-MS: *m/e* 260 (M + H)<sup>+</sup> (2.3 min).

REFERENCE EXAMPLE 3



*N*-[3-(4-Chlorophenyl)-2(S)-phenyl-1(S)-methylpropyl]-amine hydrochloride

Step A 4-(4-Chlorophenyl)-3(S)-phenyl-2(R)-butanol

A sample of magnesium (20 g, 0.82 mol) was activated by stirring under nitrogen for 12 h, and anhydrous ether (100 mL) was added to cover the solid material. The mixture was cooled to 0°C, and 4-chlorobenzyl chloride (40 g, 0.25 mmol) in 400 mL of anhydrous ether was added dropwise. After stirring at room temperature for 1 h, a sample of the above solution (32 mL) was added to (1*R*,2*R*)-1-phenylpropylene oxide

(1.0 g, 7.5 mmol) in 100 mL of ether at 0°C via syringe. After stirring at 0°C for 2 h, the reaction was quenched by addition of saturated aqueous ammonium chloride (100 mL). The organic layer was separated and the aqueous layer extracted with ether (2 x 100 mL). The combined organic extracts were washed with brine, dried over  
 5 anhydrous magnesium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with hexane to 15% ethyl acetate in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.28-7.02 (m, 9H), 4.01 (m, 1H), 3.14 (dd, 1H), 2.97 (dd, 1H), 2.85 (m, 1H), 1.12 (d, 3H).


10

Step B N-[3-(4-chlorophenyl)-2(S)-phenyl-1(S)-methylpropyl]-amine, hydrochloride

The product of Step A (4-(4-chlorophenyl)-3(S)-phenyl-2(R)-butanol, 1.8 g, 7.0 mmol) was converted to the title compound following the steps described in  
 15 Reference Example 2, Steps F through I, except hydrogen chloride in dioxane (4 M) was used in place of hydrogen chloride in ethyl acetate. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.35-6.98 (m, 9H), 3.62 (m, 1H), 3.20 (dd, 1H), 3.05 (m, 1H), 2.98 (dd, 1H), 1.19 (d, 3H). LC-MS: m/e 260 (M + H)<sup>+</sup> (2.3 min).

20

#### REFERENCE EXAMPLE 4



N-[3-(4-chlorophenyl)-2-(3-pyridyl)-1-methylpropyl]-amine, hydrochloride (mixture of diastereomers α/β 10:1)

25 Step A 4-(4-Chlorophenyl)-3-pyridyl-2-butanone

To a solution of 3-pyridylacetone hydrochloride (Wibaud, van der V. *Recl. Trav. Chim. Pays-Bas*. **1952**, 71, 798) (10 g, 58 mmol) and 4-chlorobenzyl chloride (9.1 g, 58 mmol) in 100 mL of methylene chloride at -78°C was added cesium hydroxide




monohydrate (39 g, 0.23 mol) and tetrabutyl ammonium iodide (1 g). The reaction was allowed to warm to room temperature overnight, and the resulting mixture was partitioned between brine (100 mL) and ethyl acetate (100 mL). The organic layer was separated and the aqueous layer extracted with ethyl acetate (2 x 100 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.42 (d, 1H), 8.34 (d, 1H), 7.72 (d, 1H), 7.40 (dd, 1H), 7.18 (d, 2H), 7.06 (d, 1H), 4.23 (dd, 1H), 3.38 (dd, 1H), 2.95 (dd, 1H), 2.10 (s, 3H). LC-MS: m/e 260 (M + H)<sup>+</sup> (1.9 min).

Step B N-[3-(4-chlorophenyl)-2-(3-pyridyl)-1-methylpropyl]-amine, hydrochloride (mixture of diastereomers α/β 10:1)

The product of Step 1 (4-(4-chlorophenyl)-3-pyridyl-2-butanone) was converted to the title compound following the procedure described in Reference Example 2, Steps E through I. LC-MS: m/e 261 (M + H)<sup>+</sup> (1.2 min).

#### REFERENCE EXAMPLE 5



2-Amino-4-(4-chlorophenyl)-3-(3-fluorophenyl)butane hydrochloride salt (mixture of diastereomers α/β 5:1)

Step A Methyl 3-(4-Chlorophenyl)-2-(3-fluorophenyl)propionate

To a solution of 3-fluorophenylacetic acid (5.0 g, 32 mmol) in methanol (25 mL) and methylene chloride (25 mL) at 0°C was added trimethylsilyldiazomethane (2 M in hexane, 30 mL, 60 mmol). After stirring at room temperature for 15 min, the reaction mixture was concentrated to dryness, and the residue was azeotroped with toluene to give the crude methyl 3-fluorophenylacetate (5.6 g), which was used without further purification. The crude methyl 3-

fluorophenylacetate obtained above was converted to the title compound by reacting with 4-chlorobenzyl bromide (4.6 g, 22 mmol) and sodium hexamethyldisilazide (1 M in tetrahydrofuran, 15 mL, 15 mmol) following the procedure described in Reference Example 2, Step A. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.35-6.88 (m, 8H), 3.92 (t, 1H),

5 3.60 (s, 3H), 3.34 (dd, 1H), 3.00 (dd, 1H). LC-MS: m/e 305 (M + Na)<sup>+</sup> (3.9 min).

Step B N-Methoxy-N-methyl-3-(4-chlorophenyl)-2-(3-fluorophenyl)propanamide

To a suspension *N*-methoxy-*N*-methylamine hydrochloride (2.0 g, 21 mmol) in 50 mL of methylene chloride at 0°C was added dimethylaluminum chloride (1 M in hexane, 21 mL, 21 mmol). After stirring at room temperature for 1 h, a solution of methyl 3-(4-chlorophenyl)-2-(3-fluorophenyl)propionate (Step 1, 2.0 g, 10 mmol) in methylene chloride (10 mL) was added, and the resulting mixture was stirred overnight. The reaction mixture was quenched by addition of methanol (5 mL), and the resulting mixture was concentrated with silica gel (50 g). The material was loaded onto a silica gel column, which was eluted with 10% ethyl acetate in hexane to 2% ammonia in methanol (2 M) in 10% ethyl acetate/hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.35-6.90 (m, 8H), 4.39 (br, 1H), 3.41 (s, 3H), 3.38-3.30 (m, 1H), 3.08 (s, 3H), 2.92 (dd, 1H). LC-MS: m/e 322 (M + H)<sup>+</sup> (3.6 min).

Step C 4-(4-Chlorophenyl)-3-(3-fluorophenyl)-2-butanol

20 The product of Step 2 (*N*-methoxy-*N*-methyl-3-(4-chlorophenyl)-2-phenylpropionamide) (0.74 g, 2.3 mmol) was converted to the title compound as a 5:1 mixture of diastereomers following the procedure described in Reference Example 2, Steps D through E. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.22-6.78 (m, 8H), 3.98 (m, 1H), 3.11 (dd, 1H), 2.94 (dd, 1H), 2.85 (m, 1H), 1.08 (d, 3H).

25 Step D 2-Azido-4-(4-chlorophenyl)-3-(3-fluorophenyl)butane


To a mixture of 4-(4-chlorophenyl)-2-(3-fluorophenyl)-2-butanol (Step 3, 0.65 g, 2.3 mmol), triphenylphosphine (1.2 g, 4.7 mmol), imidazole (0.32 g, 4.7 mmol) and zinc azide dipyridine complex (Viaud, M.C.; Rollin, P. *Synthesis* **1990**, 130) (0.72 g, 2.3 mmol) in 10 mL of methylene chloride was added diethylazodicarboxylate (0.73 mL, 4.7 mmol) at 0°C. After stirring at room temperature for 30 min, the resulting mixture was concentrated with silica gel (20 g) and loaded onto a silica gel column, which was eluted with 2% ether in hexane to 2% ammonia in methanol (2 M) in 2% ether/hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.25-6.85 (m, 8H), 3.76 (m, 1H), 3.33 (m, 1H), 2.92 (m, 2H), 1.15 (d, 3H).

Step E      2-Amino-4-(4-Chlorophenyl)-3-(3-fluorophenyl)butane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  5:1)

The product of Step D (2-azido-4-(4-chlorophenyl)-3-(3-fluorophenyl)butane) (0.49 g, 1.6 mmol) was converted to the title compound following the steps described in Reference Example 2, Steps H - I.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.32-6.90 (m, 7H), 3.61 (m, 1H), 3.20 (dd, 1H), 3.11 (m, 1H), 2.92 (dd, 1H), 1.19 (d, 3H). LC-MS: m/e 278 ( $\text{M} + \text{H}$ )<sup>+</sup> (2.4 min).


The amines of Reference Examples 6-7 were prepared according to the procedures described in Reference Example 5 substituting the appropriate pyridyl derivative for 3-fluorophenylacetic acid in Step A:

REFERENCE EXAMPLE 6



2-Amino-4-(4-chlorophenyl)-3-(2-pyridyl)butane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1). LC-MS: m/e 261 ( $\text{M} + \text{H}$ )<sup>+</sup> (1.6 min).

REFERENCE EXAMPLE 7




2-Amino-4-(4-chlorophenyl)-3-(4-pyridyl)butane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1)

Trimethylaluminum was used in place of dimethylaluminum chloride at Step B of Reference Example 5. LC-MS: m/e 261 (M + H)<sup>+</sup>.

5

#### REFERENCE EXAMPLE 8




10 2-Amino-4-(5-chloro-2-pyridyl)-3-phenylbutane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1)

5-Chloro-2-chloromethylpyridine (Weidmann, K. et al. *J. Med. Chem.* 1992, 35, 438) and phenylacetone were used in place of 4-chlorobenzyl bromide and 3-pyridylacetone in Step A of Reference Example 4. LC-MS: m/e 261 (M + H)<sup>+</sup>.

15

#### REFERENCE EXAMPLE 9



1-Methylcyclobutanecarboxylic Acid

Step A Ethyl 1-Methylcyclobutanecarboxylic Acid

20 To a solution of ethyl cyclobutanecarboxylate (1.0 g, 7.8 mmol) and methyl iodide (2.4 mL, 39 mmol) in 20 mL of anhydrous tetrahydrofuran at -78°C was added potassium hexamethyldisilazide (0.5 M in toluene, 23 mL, 12 mmol), and the reaction was allowed to warm to room temperature overnight. After quenching with saturated ammonium chloride (10 mL), the resulting mixture was partitioned between water


(100 mL) and ethyl acetate (100 mL). The organic layer was separated, washed with water and brine, dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the title compound, which was used without further purification.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  4.12 (q, 2H), 2.5-1.8 (m, 6H), 1.48 (s, 3H), 1.25 (t, 3H).

5    Step B        1-Methylcyclobutanecarboxylic Acid

To a solution of ethyl 1-methylcyclobutanecarboxylate (Step A, 1.0 g, 7.0 mmol) in water (10 mL) and dioxane (10 mL) was added lithium hydroxide monohydrate (2 g, 48 mmol), and the mixture was heated at  $100^\circ\text{C}$  for 2 days. After cooling to room temperature, the reaction was quenched with 2 M hydrochloric acid to pH = 2, and the  
10    resulting mixture was partitioned between water (100 mL) and ethyl acetate (100 mL). The organic layer was separated, washed with water and brine, dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the title compound, which was used without further purification.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  2.5-1.8 (m, 6H), 1.39 (s, 3H).

15


REFERENCE EXAMPLE 10



1-Ethylcyclobutanecarboxylic Acid

The title compound was prepared following the same procedure as described for  
20    Reference Example 9 substituting methyl iodide with ethyl iodide.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  2.44-2.34 (m, 2H), 1.96-1.84 (m, 4H), 1.81 (q, 2H), 0.84 (t, 3H).

REFERENCE EXAMPLE 11




25    1-Propylcyclobutanecarboxylic Acid

The title compound was prepared following the same procedure as described for Reference Example 9 substituting methyl iodide with propyl iodide.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  2.46-2.34 (m, 2H), 1.96-1.84 (m, 4H), 1.78-1.70 (m, 2H), 1.30-1.20 (m, 2H), 0.92 (t, 3H).

5

#### REFERENCE EXAMPLE 12




#### 1-Isopropylcyclobutanecarboxylic Acid

The title compound was prepared following the same procedure as described for Reference Example 9 substituting methyl iodide with isopropyl iodide at Step A and potassium hydroxide in dimethylsulfoxide and water for lithium hydroxide in dioxane and water.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  2.7-1.7 (m, 7 H), 0.93 (d, 6H).

10

#### REFERENCE EXAMPLE 13




15

#### 1-Benzylcyclobutanecarboxylic Acid

The title compound was prepared following the same procedure as described for Reference Example 9 substituting methyl iodide with benzyl bromide.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.26-7.14 (m, 5H), 3.09 (s, 2H), 2.42-2.34 (m, 2H), 2.11-2.04 (m, 2H), 1.95-1.83 (m, 2H).

20


#### REFERENCE EXAMPLE 14



#### 1-Phenylcyclobutanecarboxylic Acid

A mixture of 1-phenylcyclobutanecarbonitrile (5.0 g, 32 mmol) in water (50 mL) and concentrated hydrochloric acid (50 mL) was heated at 100°C for 3 h. After cooling to room temperature, the product was extracted with ethyl acetate (2x50 mL), and the ethyl acetate solution was back extracted with 2 M aqueous sodium hydroxide (2x50 mL). The aqueous extracts were neutralized with concentrated hydrochloric acid (pH = 2), and the product was extracted with ethyl acetate (2 x 50 mL). The organic extracts were dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the title compound as a white solid (0.66 g). <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.40-7.18 (m, 5H), 2.86-2.75 (m, 2H), 2.54-2.43 (m, 2H), 2.08-1.98 (m, 1H), 1.90-1.80 (m, 1H).

#### REFERENCE EXAMPLE 15



#### 1-tert-Butoxycarbonyl-3-ethylazetidine-3-carboxylic Acid

##### Step A Methyl 1-tert-Butoxycarbonylazetidine-3-carboxylate

To a solution of 1-tert-butoxycarbonylazetidine-3-carboxylic acid (0.90 g, 4.5 mmol) in methanol (10 mL) and methylene chloride (10 mL) at 0°C was added trimethylsilyldiazomethane (2 M in hexane, 4 mL, 7.0 mmol) until a yellow color persisted. The reaction was stirred at room temperature for 10 min, and was concentrated to dryness to give the title compound, which was used without further


purification.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  4.15 (d, 2H), 3.76 (s, 3H), 3.72 (d, 2H), 1.94 (q, 2H), 1.42 (s, 9H), 0.88 (t, 3H). LC-MS:  $m/e$  266 ( $\text{M} + \text{Na}$ ) $^+$ .

Step B 1-tert-Butoxycarbonyl-3-ethylazetidine-3-carboxylic Acid

The title compound was prepared following the procedure described for Reference

- 5 Example 10 substituting cyclobutanecarboxylate with methyl 1-tert-butoxycarbonylazetidine-3-carboxylate.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  4.12 (d, 2H), 3.70 (d, 2H), 1.82 (q, 2H), 1.42 (s, 9H), 0.90 (t, 3H).

REFERENCE EXAMPLE 16




10

1-tert-Butoxycarbonyl-2-ethylazetidine-2-carboxylic Acid

The title compound was prepared following the procedure described for Reference Example 15 substituting 1-tert-butoxycarbonylazetidine-3-carboxylic acid with 1-tert-butoxycarbonylazetidine-2-carboxylic acid at Step A.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):

- 15  $\delta$  3.98-3.90 (m, 1H), 3.80-3.62 (m, 1H), 2.36-1.82 (m, 4H), 1.42 (s, 9H), 1.02 (t, 3H).

REFERENCE EXAMPLE 17



N-([2-(3-Bromophenyl)-3-(4-chlorophenyl)-1,2-dimethylpropyl]amine, hydrochloride (Diastereomer  $\alpha$  and  $\beta$ )

20

Step A: 1-(3-bromophenyl)acetone



To a solution of N-methoxy-N-methylacetamide (10 g, 0.10 mol) in 200 mL of ether at 0°C was added 3-bromobenzylmagnesium bromide (0.25 M, 200 mL, 50 mmol). After stirring at 0°C for 2 h, the reaction mixture was partitioned between hexane and saturated aqueous ammonium chloride. The organic layer was separated, washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated to dryness to give the title compound, which was used without further purification. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.5-7.1 (m, 4H), 3.78 (s, 2H), 2.19 (s, 3H).

**Step B:**        3-(3-Bromophenyl)-2-butanone.

To a solution of 3-bromophenylacetone (4.7 g, 22 mmol) in acetonitrile (100 mL) was added methyl iodide (1.4 mL, 22 mmol) and cesium carbonate (14 g, 44 mmol). After stirring at room temperature for 17 h, the reaction mixture was poured into ether (100 mL) and water (100 mL). The organic layer was separated and the aqueous layer extracted with ether. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.45-7.40 (m, 2H), 7.3-7.2 (m, 2H), 3.87 (q, 1H), 2.06 (s, 3H), 1.34 (d, 3H).

**Step C:**        3-(3-Bromophenyl)-4-(4-chlorophenyl)-3-methyl-2-butanone.

To a solution of 3-(3-bromophenyl)-2-butanone (2.0 g, 8.8 mmol) in methylene chloride (100 mL) was added 4-chlorobenzyl chloride (1.4 g, 8.8 mmol), tetrabutylammonium iodide (0.16 g, 0.44 mmol) and cesium hydroxide monohydrate (5.9 g, 35 mmol). After stirring at room temperature for 3.5 h, the reaction mixture was poured into ethyl acetate (100 mL) and water (100 mL). The organic layer was separated and the aqueous layer extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over magnesium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.5-7.1 (m, 4 H), 7.08 (d, 2H), 6.68 (d, 2H), 3.16 (ABq, 2H), 1.98 (s, 3H), 1.42 (s, 3H).

**Step D:**        3-(3-Bromophenyl)-4-(4-chlorophenyl)-3-methyl-2-butanol.

To a solution of 3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methyl-2-butanone (1.6 g, 4.6 mmol) in methanol (50 mL) was added sodium borohydride (0.26 g, 6.8 mmol). After stirring at room temperature for 10 min, the reaction was quenched by addition of saturated aqueous ammonium chloride (25 mL). The precipitate was filtered off and washed with ethyl acetate (25 mL). The organic layer of the filtrate was separated, washed with brine, dried over anhydrous sodium sulfate, filtered and

concentrated to dryness. The residue was purified by flash column chromatography on silica gel eluted with 5% ethyl acetate in hexane to afford the title compound as two separate diastereomers. Faster eluting diastereomer (Diastereomer  $\alpha$ )  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.63 (s, 1H), 7.42-7.18 (m, 3H), 7.05 (d, 2H), 6.80 (d, 2H),

5 3.92 (q, 1H), 3.19 (d, 1H), 2.86 (d, 1H), 1.13 (s, 3H), 1.02 (d, 3H). Slower eluting diastereomer (Diastereomer  $\beta$ )  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ): 7.40-7.18 (m, 4H), 7.04 (d, 2H), 6.64 (d, 2H), 4.12 (q, 1H), 3.04 (ABq, 2H), 1.17 (s, 3H), 0.84 (d, 3H).

Step E: 2-Azido-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane.

To a solution of 3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methyl-2-butanol (faster eluting diastereomer, 0.90 g, 2.5 mmol) in ethyl acetate (80 mL) at  $0^\circ\text{C}$  was added triethyl amine (dried over activated molecular sieves, 0.42 mL, 3.1 mmol) and methanesulfonyl chloride (0.22 mL, 2.8 mmol). After stirring at  $0^\circ\text{C}$  for 2 h, the reaction was quenched by addition of saturated aqueous sodium bicarbonate (10 mL). After stirring at room temperature for 0.5 h, the organic layer was separated, washed  
15 with brine, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness to give the crude sulfonate, which was used without further purification.

Thus, a mixture of the sulfonate and sodium azide (0.83 g, 0.13 mol) in dimethylformamide (5 mL) was heated at  $120^\circ\text{C}$  for 4 h. The reaction mixture was cooled to room temperature and was poured into water (40 mL), and the product was  
20 extracted with ether (2 x 20 mL). The combined organic extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to dryness, and the residue was purified on a silica gel column eluting with hexane to give the title compound (Diastereomer  $\alpha$ ).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.43-7.20 (m, 4H), 7.04 (d, 2H), 6.64 (d, 2H), 4.10 (q, 1H), 3.10 (d, 1H), 3.00 (d, 1H), 1.10 (s, 3H), 1.02  
25 (d, 3H).

The slower eluting diastereomer was converted to the other diastereomer (Diastereomer  $\beta$ ) of the title compound following the same procedure as described for the faster eluting diastereomer.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.60-7.20 (m, 4H), 7.07 (d, 2H), 6.80 (d, 2H), 3.90 (q, 1H), 3.17 (d, 1H), 2.92 (d, 1H), 1.22  
30 (d, 3H), 1.20 (s, 3H).

Step F: 2-(*N*-tert-Butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane

To a solution of 2-azido-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane (Diastereomer  $\alpha$ , 0.26 g, 0.68 mmol) in ethyl acetate (5 mL) was added di(*tert*-

butyl)dicarbonate (0.18 g, 0.82 mmol) and platinum dioxide (0.025 g). The mixture was degassed and charged with hydrogen with a balloon. After stirring for 1 day, the reaction mixture was filtered through CELITE, diatomaceous earth, and the filtrate was concentrated to give diastereomer  $\alpha$  of the title compound.

- 5 Diastereomer  $\beta$  of 2-azido-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane was converted to the Diastereomer  $\beta$  of the title compound following the same procedure as described for Diastereomer  $\alpha$ .


Step G: N-[3-(4-Chlorophenyl)-2-(3-bromophenyl)-1,2-dimethylpropyl]-amine hydrochloride (Diastereomer  $\alpha$  and  $\beta$ ).

- 10 2-(*N-tert*-Butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane (Diastereomer  $\alpha$ , 0.35 g, 0.76 mmol) was treated with 4 M hydrogen chloride in dioxane (5 mL) at room temperature for 2 h. The mixture was concentrated to dryness to give Diastereomer  $\alpha$  of the title compound. LC-MS:  $m/e$  352 ( $M + H$ )<sup>+</sup> (3.0 min).

- 15 Diastereomer  $\beta$  of 2-azido-3-(3-bromophenyl)-4-(4-chlorophenyl)-3-methylbutane was converted to Diastereomer  $\beta$  of the title compound following the same procedure as described for Diastereomer  $\alpha$ . LC-MS:  $m/e$  352 ( $M + H$ )<sup>+</sup> (3.0 min).

20

#### REFERENCE EXAMPLE 18



N-([3-(4-Chlorophenyl)-2-phenyl-2-cyano-1-methyl]propyl)amine, hydrochloride

Step A: 4-(4-Chlorophenyl)-3-cyano-3-phenyl-2-butanone.

- 25 To a solution of  $\alpha$ -acetylphenylacetonitrile (1.0 g, 6.3 mmol) in acetonitrile (25 mL) was added 4-chlorobenzyl bromide (1.3 g, 6.3 mmol) and cesium carbonate (8.2 g, 25 mmol). After stirring at room temperature for 2 h, the reaction mixture was poured into ethyl acetate (100 mL) and water (100 mL). The organic layer was separated, washed with brine, dried over magnesium sulfate, filtered, and concentrated to

dryness, and the residue was purified on a silica gel column eluting with 1 to 5% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.5-6.9 (m, 9H), 3.56 (d, 1H), 3.37 (d, 1H), 2.22 (s, 3H). LC-MS: m/e 306 (M + Na)<sup>+</sup> (3.0 min).

5    Step B:        N-[4-(4-Chlorophenyl)-3-cyano-3-phenyl-2-butyldene]-2-methylpropane-(S)-sulfinamide.

To a solution of 4-(4-chlorophenyl)-3-cyano-3-phenyl-2-butanone (1.9 g, 6.7 mmol) and (S)-2-methylsulfinamide (0.74 g, 6.1 mmol) in tetrahydrofuran (25 mL) was added titanium tetraethoxide (4.0 mL, 18 mmol). After stirring at 60°C for 6 h and  
10    75°C for 18 h, the reaction mixture was poured into a well-stirred brine solution (50 mL). The resulting mixture was filtered through CELITE diatomaceous earth and washed with ethyl acetate (20 mL), and the filtrate was extracted with ethyl acetate (2 x 50 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column  
15    chromatography on silica gel eluted with 10 to 20% ethyl acetate in hexane to give the title compound as a 1:1 mixture of diastereomers. LC-MS: m/e 387 (M + H)<sup>+</sup> (3.6 min).


Step C:        N-([3-(4-Chlorophenyl)-2-cyano-2-phenyl-1-methyl]propyl)-2-methylpropane-(S)-sulfinamide

20    To a solution of N-[4-(4-chlorophenyl)-3-cyano-3-phenyl-2-butyldene]-2-methylpropane-(S)-sulfinamide (0.50 g, 1.3 mmol) in methanol (25 mL) at 0°C was added sodium borohydride (0.075 g, 1.9 mmol). After stirring for 15 min, the reaction was quenched by addition of saturated aqueous ammonium chloride (25 mL). The organic layer was separated and the aqueous layer was extracted with ethyl  
25    acetate. The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated to dryness to give the title compound. LC-MS: m/e 389 (M + H)<sup>+</sup> (3.4 min).

Step D:        N-([3-(4-Chlorophenyl)-2-cyano-2-phenyl-1-methyl]propyl)amine hydrochloride salt

30    N-([3-(4-Chlorophenyl)-2-cyano-2-phenyl-1-methyl]propyl)-2-methylpropane-(S)-sulfinamide (0.55 g, 1.4 mmol) in methanol (20 mL) was added 4 M hydrogen chloride in dioxane (25 mL). After stirring for 30 min, the mixture was concentrated to dryness to give the title compound as a mixture of diastereomers (α and β). LC-MS: m/e 285 (M + H)<sup>+</sup> (Major diastereomer: 2.0; Minor diastereomer: 2.1 min).

35

REFERENCE EXAMPLE 19

N-([3-(4-Chlorophenyl)-2-(3-bromophenyl)-2-hydroxy]propyl)amine hydrochloride

Step A: 1-Bromo-3-([(N-tert-butoxycarbonyl)amino]acetyl)benzene

- 5 To a solution of 1-bromo-3-iodobenzene (8.8 mL, 69 mmol) in 200 mL of ether at -78°C was added tert-butyllithium (1.7 M in pentane, 40 mL, 69 mmol). After stirring at -78°C for 30 min, a solution of N-(tert-butoxycarbonyl)glycine N'-methoxy-N'-methylamide (5.0 g, 23 mmol) in 100 mL of tetrahydrofuran was added. After stirring at -78°C for 2 h, the reaction was allowed to warm up to 0°C, and was
- 10 quenched with dilute aqueous ammonium chloride (200 mL). The organic layer was separated, washed with brine, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 5-10% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.12 (s, 1H), 7.97 (d, 1H), 7.80 (d, 1H), 7.43 (t, 1H), 4.50 (s, 2H), 1.42 (s, 9H).
- 15


Step B: 3-(4-Chlorophenyl)-2-(3-bromophenyl)-1-[(N-butoxycarbonyl)amino]-2-hydroxy]propane

- To a solution of 1-bromo-3-([(N-tert-butoxycarbonyl)amino]acetyl)benzene (0.65 g, 2.1 mmol) in 25 mL of ether at -78°C was added 4-chlorobenzylmagnesium chloride (0.25 M in ether, 21 mL, 5.2 mmol). The reaction was allowed to warm up to -10°C over 3.5 h and was quenched at -10°C with saturated aqueous ammonium chloride (50 mL). The organic layer was separated, washed with water, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness. The residue was purified by flash column chromatography on silica gel eluted with 5-10% ethyl acetate in hexane
- 20 to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.5-7.1 (m, 4H), 7.10 (d, 2H), 6.92 (d, 2H), 3.55 (d, 2H), 3.40 (d, 2H), 3.02 (ABq, 2H), 1.38 (s, 9H).
- 25

Step C: N-([3-(4-Chlorophenyl)-2-(3-bromophenyl)-2-hydroxy]propyl)amine hydrochloride

To a solution of 3-(4-chlorophenyl)-2-(3-bromophenyl)-1-[(N-butoxycarbonyl)amino-2-hydroxy]propane (0.38 g, 0.86 mmol) in ethyl acetate (10 mL) was added 4 M hydrogen chloride in dioxane (20 mL). After stirring for 1 h, the mixture was concentrated to dryness to give the title compound. LC-MS: m/e 340 (M + H)<sup>+</sup> (2.8 min).

### REFERENCE EXAMPLE 20



N-([3-(4-Chlorophenyl)-2-(3-bromophenyl)-2-fluoro-1(S)-methyl]propyl)amine hydrochloride

Step A: 3-(3-Bromophenyl)-2(S)-[(N-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-hydroxy] butane

The title compound was prepared following the same procedure described for Reference Example 19, Step A and B substituting N-(tert-butoxycarbonyl)glycine N'-methoxy-N'-methylamide with N-(tert-butoxycarbonyl)-L-alanine N'-methoxy-N'-methylamide. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.5-7.0 (m, 6H), 6.82 (d, 2H), 4.11 (m, 1H), 3.07 (ABq, 2H), 1.50 (s, 9H), 0.87 (d, 3H).

Step B: 3-(3-Bromophenyl)-2(S)-[(N-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-fluoro] butane

To a solution of 3-(3-bromophenyl)-2(S)-[(N-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-hydroxy] butane (2.0 g, 4.4 mmol) in 15 mL of methylene chloride at -78°C was added (dimethylamino)sulfur trifluoride (1.1 mL, 8.8 mmol), and the reaction was allowed to warm up to room temperature over 2.5 h. The reaction was quenched by carefully transferring to a well-stirred saturated aqueous sodium bicarbonate (50 mL). The mixture was extracted with ether (2 x 50 mL), and the combined extracts were dried over anhydrous magnesium sulfate, filtered and concentrated to dryness. The residue was purified on a silica gel column eluting with 4-20% ethyl acetate in hexane to give the title compound as one major diastereomer


and some contamination of the corresponding dehydration product.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.4-7.1 (m, 4H), 7.06 (d, 2H), 6.85 (d, 2H), 4.19 (m, 1H), 3.43 (dd, 1H), 3.10 (dd, 1H), 1.50 (s, 9H), 0.93 (d, 3H).

Step C: N-([3-(4-Chlorophenyl)-2-(3-bromophenyl)-2-fluoro-1(S)-methyl]propyl)amine hydrochloride

To a solution of 3-(3-bromophenyl)-2(S)-[(N-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-fluoro] butane (0.16 g, 0.35 mmol) in ethyl acetate (1 mL) was added 4 M hydrogen chloride in dioxane (4 mL). After stirring for 2 h, the mixture was concentrated to dryness to give the title compound. LC-MS:  $m/e$  356 ( $M + H$ ) $^+$  (3.1 min).

The amines of Reference Examples 21-28 were prepared by the same procedures described in Reference Example 1:

#### REFERENCE EXAMPLE 21



2-Amino-3,4-diphenylbutane hydrochloride salt


Diastereomer  $\alpha$ :

LC-MS: calculated for  $\text{C}_{16}\text{H}_{19}\text{N}$  225, observed  $m/e$  226 ( $M + H$ ) $^+$  (2.0 min).

Diastereomer  $\beta$ :

LC-MS: calculated for  $\text{C}_{16}\text{H}_{19}\text{N}$  225, observed  $m/e$  226 ( $M + H$ ) $^+$  (1.9 min).

#### REFERENCE EXAMPLE 22



3-Amino-1,2-diphenylpentane hydrochloride salt


Diastereomer  $\alpha$ :

LC-MS: calculated for  $C_{17}H_{21}N$  239, observed  $m/e$  240 ( $M + H$ )<sup>+</sup> (2.1 min).

5 Diastereomer  $\beta$ :

LC-MS: calculated for  $C_{17}H_{21}N$  239, observed  $m/e$  240 ( $M + H$ )<sup>+</sup> (2.0 min).

REFERENCE EXAMPLE 23



10

1-Amino-1,2,3-triphenylpropane *p*-toluenesulfonate salt

Diastereomer  $\alpha$ :


LC-MS: calculated for  $C_{21}H_{21}N$  287, observed  $m/e$  288 ( $M + H$ )<sup>+</sup> (2.3 min).

Diastereomer  $\beta$ :

15 LC-MS: calculated for  $C_{21}H_{21}N$  287, observed  $m/e$  288 ( $M + H$ )<sup>+</sup> (2.3 min).

REFERENCE EXAMPLE 24





2-Amino-4-(4-chlorophenyl)-3-phenylbutane hydrochloride salt


Diastereomer  $\alpha$ :

LC-MS: calculated for  $C_{16}H_{18}ClN$  259, observed  $m/e$  260 ( $M + H$ )<sup>+</sup> (2.3 min).

5 Diastereomer  $\beta$ :

LC-MS: calculated for  $C_{16}H_{18}ClN$  259, observed  $m/e$  260 ( $M + H$ )<sup>+</sup> (2.2 min).

REFERENCE EXAMPLE 25



10

2-Amino-3-(4-chlorophenyl)-4-phenylbutane hydrochloride salt


Diastereomer  $\alpha$ :

LC-MS: calculated for  $C_{16}H_{18}ClN$  259, observed  $m/e$  260 ( $M + H$ )<sup>+</sup> (2.3 min).

Diastereomer  $\beta$ :

15 LC-MS: calculated for  $C_{16}H_{18}ClN$  259, observed  $m/e$  260 ( $M + H$ )<sup>+</sup> (2.1 min).

REFERENCE EXAMPLE 26



2-Amino-4-(4-methoxycarbonylphenyl)-3-phenylbutane hydrochloride salt


Diastereomer  $\alpha$ :

LC-MS: calculated for  $C_{18}H_{21}NO_2$  283, observed m/e 284 ( $M + H$ )<sup>+</sup> (2.0 min).

5 Diastereomer  $\beta$ :

LC-MS: calculated for  $C_{18}H_{21}NO_2$  283, observed m/e 284 ( $M + H$ )<sup>+</sup> (1.9 min).

REFERENCE EXAMPLE 27




10

2-Amino-3-(2-Chlorophenyl)-4-phenylbutane (mixture of diastereomers  $\alpha/\beta$  1:2)

LC-MS: calculated for  $C_{16}H_{18}ClN$  259, observed m/e 260 ( $M + H$ )<sup>+</sup> (1.9/2.0 min).

REFERENCE EXAMPLE 28

15



2-Amino-3-(4-methoxyphenyl)-4-phenylbutane (mixture of diastereomers  $\alpha/\beta$  2:5)


LC-MS:  $m/e$  256 ( $M + H$ )<sup>+</sup> (1.7 min).

5

The amines of Reference Examples 29-30 were prepared according to the procedures described in Reference Example 32:

REFERENCE EXAMPLE 29

10




2-Amino-4-(4-chlorophenyl)-3-(2-fluorophenyl)butane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1)

LC-MS:  $m/e$  278 ( $M + H$ )<sup>+</sup> (2.3 min).

15

REFERENCE EXAMPLE 30




2-Amino-4-(4-chlorophenyl)-3-(4-fluorophenyl)butane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1)

LC-MS:  $m/e$  278 ( $M + H$ )<sup>+</sup> (2.5 min).

5

#### REFERENCE EXAMPLE 31



2-Amino-4-(4-cyanophenyl)-3-phenylbutane hydrochloride salt (mixture of diastereomers  $\alpha/\beta$  10:1)

10

Step A 4-(4-Cyanophenyl)-3-phenyl-2-butanone.

To a solution of phenylacetone (1.2 g, 9.0 mmol) and 4-cyanobenzyl chloride (1.4 g, 9.0 mmol) in 20 mL  $CH_2Cl_2$  at  $-78^\circ C$  was added cesium hydroxide monohydrate (4.5 g, 27 mmol) and tetrabutyl ammonium iodide (20 mg). The reaction was allowed to warm to room temperature over 6 h, and the resulting mixture partitioned between brine (100 mL) and EtOAc (100 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 100 mL). The combined organic extracts were dried over  $MgSO_4$ , filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 20-50% EtOAc in hexane to give the title compound.  $^1H$  NMR (500 MHz,  $CD_3OD$ ):  $\delta$  7.52 (d, 2H),

20


7.34-7.16 (m; 7H), 4.12 (dd, 1H), 3.41 (dd, 1H), 3.00 (dd, 1H). LC-MS: m/e 250 (M + H)<sup>+</sup> (3.2 min).

Step B      2-Amino-4-(3-cyanophenyl)-3-phenylbutane hydrochloride salt  
                  (mixture of diastereomers  $\alpha/\beta$  10:1).

- 5    The product of Step A (4-(4-cyanophenyl)-3-phenyl-2-butanone) (1.0 g, 4.0 mmol) was converted to the title compound following the procedure described in Reference Example 2, Steps E-I. LC-MS: m/e 251 (M + H)<sup>+</sup> (1.9 min).

10

### REFERENCE EXAMPLE 32



2-Amino-4-(2,4-dichlorophenyl)-3-(4-chlorophenyl)butane hydrochloride salt (3 isomers)

- 15    Step A      Methyl 3-(2,4-Dichlorophenyl)-2-(4-chlorophenyl)propionate.  
          A sample of 4-chlorophenylacetic acid (4.2 g, 25 mmol) was converted to the title compound (6.5 g) following the procedure in Reference Example 5, Step A substituting 4-chlorophenylacetic acid for 3-fluorophenylacetic acid and 2,4-dichlorobenzyl bromide for 4-chlorobenzyl. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.40  
 20    (d, 1H), 7.32-7.22 (m, 4 H), 7.15 (dd, 1H), 7.08 (d, 1H), 4.00 (t, 1H), 3.62 (s, 3H), 3.44 (dd, 1H), 3.12 (dd, 1H).

Step B      3-(2,4-Dichlorophenyl)-2-(4-chlorophenyl)propanol.

- To a solution of methyl 3-(2,4-dichlorophenyl)-2-(4-chlorophenyl) propionate (6.4 g, 8.6 mmol) in 50 mL ether at -40°C was added lithium aluminum hydride (1.4 g, 37  
 25    mmol), and the reaction was allowed to warm to room temperature over 2 h. The reaction was quenched by addition of MeOH (3 mL) dropwise at -10°C, and the mixture was partitioned between 100 mL saturated ammonium chloride and EtOAc (100 mL). The organic layer was separated and the aqueous

layer extracted with EtOAc (2 x 100 mL). The combined organic extracts were dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to dryness to give the title compound, which was used without further purification. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.4-6.9 (m, 7H), 3.72 (m, 2H), 3.24 (dd, 1H), 3.16 (m, 1H), 2.85 (dd,

5 1H).

Step C 3-(2,4-Dichlorophenyl)-2-(4-chlorophenyl)propanal.

To a solution of 3-(2,4-dichlorophenyl)-2-(4-chlorophenyl)propanol (Step B, 0.89 g, 2.8 mmol) in 20 mL CH<sub>2</sub>Cl<sub>2</sub> was added crushed activated molecular sieves (4 g).

After stirring at room temperature for 10 min, pyridinium chlorochromate (0.90 g, 4.2 mmol) was added. After stirring at room temperature for 1 h, CELITE diatomaceous earth (4 g) was added followed by 100 mL ether. The resulting mixture was filtered through a silica gel pad, which was washed with ether (2 x 50 mL). The filtrate was concentrated to dryness and azeotroped with toluene to give the title compound, which was used without further purification.

15 Step D N-[3-(2,4-Dichlorophenyl)-2-(4-chlorophenyl)propylidene]-2-methylpropanesulfonamide.

To a solution of 3-(2,4-dichlorophenyl)-2-(4-chlorophenyl)propanal (Step C, 0.90 g, 2.8 mmol) in 6 mL THF was added (R)-(+)-2-methyl-2-propane-sulfonamide (0.5 gm, 4.1 mmol) followed by the addition of titanium tetraethoxide (1.5 mL, 8.0 mmol).

20 After stirring at room temperature overnight, the reaction mixture was added to a well-stirred brine solution (50 mL). The resulting mixture was filtered through CELITE diatomaceous earth and washed with EtOAc (20 mL), and the filtrate was extracted with EtOAc (2 x 50 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 10% ether in hexane to give the title compound as a 1:1 mixture of diastereomers. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.11 (m, 1H), 7.41 (m, 1H), 7.35-7.31 (m, 4 H), 7.16-7.06 (m, 2H), 4.26 (m, 1H), 3.78-3.58 (m, 1H), 3.22-3.14 (m, 1H), 1.13/1.12 (s, 9H).


25 Step E N-[3-(2,4-Dichlorophenyl)-2-(4-chlorophenyl)-1-methylpropyl]-2-methylpropanesulfonamide (3 isomers).

To a solution of N-[3-(2,4-dichlorophenyl)-2-(4-chlorophenyl)-1-methylpropylidene]-2-methylpropanesulfonamide (Step D, 0.51 g, 1.3 mmol) in 6 mL CH<sub>2</sub>Cl<sub>2</sub> at -60°C was added methylmagnesium bromide (3 M in ether, 0.90 mL, 2.7 mmol). After stirring at -60°C for 6 h, the reaction was allowed to warm to room temperature overnight.

35 The resulting mixture was partitioned between saturated aqueous ammonium chloride

- (50 mL) and EtOAc (50 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 50 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 30 to 50% EtOAc in hexane to give the title compound as one pure faster eluting enantiomer and a 1:1 mixture of slower co-eluting diastereomers. The addition of the methyl Grignard reagent was apparently stereoselective for one of the sulfinamide diastereomers.
- Faster eluting isomer:  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.30 (d, 1H), 7.22 (d, 2H), 7.12 (d, 2H), 7.03 (dd, 1H), 6.94 (d, 1H), 3.62 (m, 1H), 3.56 (dd, 1H), 2.97 (dd, 1H), 1.23 (s, 9H), 1.04 (d, 3H). LC-MS:  $m/e$  432 ( $\text{M} + \text{H}$ ) $^+$  (4.2 min).
- Slower eluting isomers (1:1):  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.33/7.30 (d, 1H), 7.21/7.18 (d, 2H), 7.06/7.04 (d, 2H), 6.99/6.97 (dd, 1H), 6.79/6.75 (d, 1H), 3.70-3.55 (m, 1H), 3.07/2.97 (m, 1H), 2.90/2.80 (dd, 1H), 1.32/0.95 (s, 9H), 1.49/1.10 (d, 3H).
- Step F      2-Amino-4-(2,4-dichlorophenyl)-3-(4-chlorophenyl)butane  
hydrochloride (3 isomers).
- To a solution of *N*-[3-(2,4-dichlorophenyl)-2-(4-chlorophenyl)-1-methylpropyl]-2-methylpropanesulfinamide (Step F, faster eluting isomer, 50 mg, 0.11 mmol) in 5 mL MeOH was added hydrogen chloride in dioxane (4 M, 2 mL). After stirring at room temperature for 10 min, the reaction mixture was concentrated to dryness to give the title compound as one pure isomer.
- Isomer 1:  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.35 (d, 1H), 7.29 (d, 2H), 7.15 (d, 2H), 7.06 (dd, 1H), 6.91 (d, 1H), 3.68 (m, 1H), 3.36 (dd, 1H), 3.06 (dd, 1H), 1.18 (d, 3H). LC-MS:  $m/e$  328 ( $\text{M} + \text{H}$ ) $^+$  (2.8 min).
- The two slower co-eluting isomers were treated in the same fashion to give two other isomers of the title compound. Isomer 2 and 3 (1:1): LC-MS:  $m/e$  328 ( $\text{M} + \text{H}$ ) $^+$  (2.7/2.8 min).

### REFERENCE EXAMPLE 33



2-Amino-4-(4-chloro-2-fluorophenyl)-3-(4-chlorophenyl)butane hydrochloride salt  
(Isomers, 1, 2 and 3).

The title compound was prepared according to the procedures of


- 5 Reference Example 40 substituting 2,5-dichlorobenzyl bromide with 4-chloro-2-fluorobenzyl bromide.

Isomer 1: LC-MS:  $m/e$  312 ( $M + H$ )<sup>+</sup> (2.6 min).

Isomer 2 and 3 (1:1): LC-MS:  $m/e$  312 ( $M + H$ )<sup>+</sup> (2.5/2.6 min).

10

REFERENCE EXAMPLE 34



2-(4-Chlorophenyl)-2-(4-chlorophenyl)ethanol hydrochloride salt.

Step A 2-(4-Chlorophenyl)-2-(4-chlorophenyl)ethanol.

- 15 To a suspension of 2-(4-chlorophenyl)-2-(4-chlorophenyl)acetic acid (Newman et al *J. Amer. Chem. Soc.* **1947**, *69*, 718) (1.0 g, 3.4 mmol) in 10 mL THF at 0°C was added borane (1 M in THF, 6.8 mL, 6.8 mmol). After stirring at room temperature for 2 h, the reaction was quenched by addition of 2 M hydrochloric acid (10 mL). The volatile materials were removed on a rotary evaporator, and the resulting mixture was
- 20 partitioned between brine (20 mL) and EtOAc (30 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 20 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to



dryness to give the title compound, which was used without further purification. LC-MS:  $m/e$  283 ( $M + H$ )<sup>+</sup> (3.4 min).

Step B 2-(4-Chlorophenoxy)-2-(4-chlorophenyl)ethyl Azide


- 5 2-(4-Chlorophenoxy)-2-(4-chlorophenyl)ethanol (Step A, 0.45 g, 2.4 mmol) was converted to the title compound (0.29 g) following the procedure described in Reference Example 5, Step D. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.41 (d, 2H), 7.37 (d, 2H), 7.18 (d, 2H), 6.86 (d, 2H), 5.42 (dd, 1H), 3.69 (dd, 1H), 3.45 (dd, 1H). LC-MS:  $m/e$  308 ( $M + H$ )<sup>+</sup> (4.3 min).

Step C 2-(4-Chlorophenoxy)-2-(4-chlorophenyl)ethylamine.

- 10 To a solution of 2-(4-chlorophenoxy)-2-(4-chlorophenyl)ethyl azide (Step B, 0.23 g, 0.75 mmol) in 4 mL THF at -20°C was added trimethylphosphine (0.18 mL, 1.8 mmol), and the reaction was allowed to warm to room temperature over 2 h. Lithium hydroxide monohydrate (61 mg, 1.5 mmol) was added followed by 2 mL water. After stirring at room temperature for 30 min, the reaction was quenched by addition of 2 M
- 15 hydrochloric acid (final pH = 2). The volatile materials were removed on a rotary evaporator, and the resulting mixture was partitioned between brine (20 mL), 5 N aqueous sodium hydroxide (20 mL), ether (20 mL) and toluene (20 mL). The organic layer was separated and the aqueous layer extracted with ether (40 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to
- 20 dryness to give the title compound (0.43 g), which was contaminated with trimethylphosphine oxide and was used without further purification. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.46-7.40 (m, 4H), 7.20 (d, 2H), 6.91 (d, 2H), 5.53 (m, 2H), 3.36 (m, 2H). LC-MS:  $m/e$  282 ( $M + H$ )<sup>+</sup> (2.5 min).

25

REFERENCE EXAMPLE 35



2,2-Bis(4-chlorophenyl)ethylamine hydrochloride saltStep A      Methyl 3,3-Bis(4-chlorophenyl)propenoate

A mixture of di(4-chlorophenyl)ketone (7.5 g, 30 mmol) and methyl (triphenylphosphoranylidene)acetate (10 g, 30 mmol) in 20 mL toluene was heated at 130°C while allowing the solvent to slowly evaporate overnight. The resulting mixture was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and toluene (20 mL) and was concentrated with 30 g silica gel. The material was loaded onto a silica gel column, which was eluted with 6:3:1 hexane/CH<sub>2</sub>Cl<sub>2</sub>/ether to give the title compound.

Step B      Methyl 3,3-Bis(4-chlorophenyl)propionate

A suspension of methyl 3,3-bis(4-chlorophenyl)propenoate (Step A, 3.0 g, 14 mmol) and platinum dioxide (0.30 g) in MeOH (20 mL) and 2 M aqueous hydrochloric acid (1 mL) was degassed and filled with hydrogen with a balloon. After stirring at room temperature for 2 h, the reaction mixture was filtered through CELITE diatomaceous earth, and the filtrate was concentrated to dryness. The residue was dissolved in 50 mL ether and was concentrated with 20 g silica gel. The material was loaded onto a silica gel column, which was eluted with 10% ether in hexane to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.29-7.22 (m, 4H), 4.50 (t, 1H), 3.56 (s, 3H), 3.07 (d, 2H). LC-MS: m/e 309 (M + H)<sup>+</sup> (4.1 min).

Step C      3,3-Bis(4-chlorophenyl)propionic Acid

A mixture of methyl 3,3-bis(4-chlorophenyl)propionate (Step B, 0.78 g, 3.9 mmol), lithium hydroxide monohydrate (0.33 g, 7.8 mmol) in 1:1:1 MeOH/ THF/water (15 mL) was stirred at room temperature overnight. The resulting mixture was partitioned between 2 M aqueous hydrochloric acid (50 mL) and ether (50 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 50 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.29-7.23 (m, 4H), 4.49 (t, 1H), 3.02 (d, 2H).

Step D      N-[2,2-Bis(4-chlorophenyl)ethyl]allylcarbamate.


To a solution of 3,3-bis(4-chlorophenyl)propionic acid (Step C, 0.32 g, 1.1 mmol) and triethyl amine (0.60 mL, 4.3 mmol) in 4 mL THF at 0 °C was added ethyl chloroformate (0.31 mL, 3.3 mmol). After stirring at room temperature for 30 min, the reaction was cooled to 0°C, and was added sodium azide (0.35 g, 5.4 mmol) in 2 mL water. After stirring at room temperature for 1 h, the reaction mixture was partitioned between brine (20 mL) and EtOAc (20 mL). The organic layer was

separated and the aqueous layer extracted with EtOAc (2 x 20 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was dissolved in allylic alcohol (1 mL) and toluene (1 mL). After stirring at 80°C overnight, the reaction mixture was concentrated to dryness, and the residue was purified by flash column chromatography on silica gel column eluted with 20% EtOAc in hexane to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.30-7.21 (m, 4H), 5.84 (m, 1H), 5.17 (dd, 1H), 5.10 (dd, 1H), 4.46 (d, 2H), 4.22 (t, 1H), 3.68 (d, 2H). LC-MS: m/e 350 (M + H)<sup>+</sup> (3.9 min).

Step E 2,2-Bis(4-chlorophenyl)ethylamine hydrochloride salt.

- 10 To a solution of *N*-[2,2-bis(4-chlorophenyl)ethyl]allylcarbamate (Step D, 0.26 g, 0.73 mmol) in 1.5 mL THF at 0°C was added tetrakis (triphenylphosphine)palladium (85 mg, 0.073 mmol) and triphenylsilane (0.18 mL, 1.1 mmol). After stirring at 0°C for 1 h, the reaction mixture was partitioned between ether (20 mL) and 2 M hydrochloric acid (20 mL). The aqueous layer was separated, and was added 5 N aqueous sodium
- 15 hydroxide (final pH > 12). The product was extracted with ether (3 x 30 mL), and the combined extracts were dried over sodium hydroxide, and filtered through CELITE, diatomaceous earth. After addition of 4 M hydrogen chloride in dioxane (2 mL), the filtrate was concentrated to dryness to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.40-7.34 (m, 4H), 4.28 (m, 1H), 3.62 (d, 2H). LC-MS: m/e 266 (M +
- 20 H)<sup>+</sup> (2.3 min).

REFERENCE EXAMPLE 36



- 25 2-Amino-3-(4-chlorophenylthio)-3-(4-chlorophenyl)propane hydrochloride salt (two diastereomers)

Step A Methyl 2-(4-Chlorophenylthio)-2-(4-chlorophenyl)acetate.

To a solution of 2-(4-chlorophenylthio)-2-(4-chlorophenyl)acetic acid (Nicolaescu et al *Rev. Roum. Chim.* **1979**, 24, 137) (1.0 g, 3.0 mmol) in MeOH (10 mL) and CH<sub>2</sub>Cl<sub>2</sub>


(10 mL) at 0°C was added trimethylsilyldiazomethane (2 M in hexane) until a yellow color persisted. Concentration afforded the title compound, which was used without further purification.

Step B      2-Amino-3-(4-chlorophenylthio)-3-(4-chlorophenyl)propane  
 5              hydrochloride salt (two diastereomers)

The product of Step A (methyl 2-(4-chlorophenylthio)-2-(4-chlorophenyl)acetate) (1.1 g, 3.0 mmol) was converted to the title compound following the procedures described in Reference Example 5, Steps B-E.

LC-MS:  $m/e$  312 ( $M + H$ )<sup>+</sup> (2.7 min).

### REFERENCE EXAMPLE 37



2-Amino-3,4-bis(4-chlorophenyl)-2-methylbutane hydrochloride salt

15    Step A      Methyl 2,3-Bis(4-chlorophenyl)propionate.

The title compound was prepared following the procedure described in Reference Example 2, Step A, substituting methyl phenylacetate with methyl 4-chlorophenylacetate. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.30-7.22 (m, 4H), 7.19 (d, 2H), 7.09 (d, 2H), 3.90 (t, 1H), 3.58 (s, 3H), 3.32 (dd, 1H), 2.98 (dd, 1H).

20    Step B      3,4-Bis(4-chlorophenyl)-2-methyl-2-butanol.

To a solution of methyl 2,3-bis(4-chlorophenyl)propionate (2.6 g, 8.4 mmol) in ether (20 mL) was added methylmagnesium bromide (3 M in ether, 8.4 mL, 25 mmol) at -10°C, and the reaction was allowed to warm to room temperature over 2 h. The reaction mixture was poured into saturated aqueous ammonium chloride (100 mL), and the product was extracted with EtOAc (3 x 100mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to dryness to give the title compound, which was used without further purification. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.17 (ABq, 4H), 7.06 (d, 2H), 6.93 (d, 2H), 3.32 (dd, 1H), 2.94 (dd, 1H), 2.84 (dd, 1H), 1.20 (s, 3H), 1.16 (s, 3H).


Step C      *N*-[2,3-Bis(4-chlorophenyl)-1,1-dimethylpropyl]chloroacetamide.

To a solution of 3,4-bis(4-chlorophenyl)-2-methyl-2-butanol (Step B, 1.4 g, 4.5 mmol) and chloroacetonitrile (0.57 mL, 9.1 mmol) in acetic acid (0.7 mL) at  $-10^{\circ}\text{C}$  was added concentrated sulfuric acid (0.31 mL, 14 mmol). After stirring at  $-10^{\circ}\text{C}$  for 15 min and room temperature for 2 h, the reaction mixture was poured onto ice (20 g), and the product was extracted with EtOAc (3 x 20 mL). The combined extracts were washed with brine/saturated aqueous sodium bicarbonate, dried over anhydrous  $\text{MgSO}_4$ , filtered, and concentrated to dryness to give the title compound.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.19 (ABq, 4H), 7.06 (d, 2H), 6.95 (d, 2H), 3.93 (ABq, 2H), 3.89 (dd, 1H), 3.10 (dd, 1H), 2.99 (dd, 1H), 1.43 (s, 3H), 1.25 (s, 3H). LC-MS:  $m/e$  384 ( $\text{M} + \text{H}$ ) $^+$  (3.9 min).

Step D      2-Amino-3,4-bis(4-chlorophenyl)-2-methylbutane hydrochloride

To a solution of *N*-[2,3-bis(4-chlorophenyl)-1,1-dimethylpropyl] chloroacetamide (Step C, 1.3 g, 3.8 mmol) in ethanol (10 mL) and acetic acid (2 mL) was added thiourea (0.34 g, 4.5 mmol). The reaction was stirred at  $80^{\circ}\text{C}$  overnight to give a white precipitate. The precipitate was removed by filtration and washed with ethanol (10 mL), and the filtrate was diluted with dilute aqueous sodium hydroxide and extracted with hexane (2 x 50 mL). The combined extracts were dried over sodium hydroxide, filtered, and concentrated to dryness, and the residue was taken up by hydrogen chloride in dioxane (4 M, 5 mL) and concentrated to dryness to give the title compound.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ): (free amine)  $\delta$  7.22-7.14 (m, 4H), 7.06 (d, 2H), 6.96 (d, 2H), 3.22 (dd, 1H), 2.95 (dd, 1H), 2.86 (dd, 1H), 1.16 (s, 3H), 1.10 (s, 3H).

REFERENCE EXAMPLE 38



2-Amino-5-methyl-3-phenylhexane hydrochloride salt

Step A      4-Methyl-2-phenylpentanoic acid


- A solution of 0.25 g (1.84 mmol) of phenylacetic acid in 3.6 mL dry THF was cooled in ice bath and 4 mL 1M lithium bis(trimethylsilyl)amide was added. After 15 min, 0.23 mL (2.02 mmol) of isobutyl iodide was added and the cold bath was removed. After stirring the reaction overnight, it was quenched with water and extracted once with EtOAc. The aqueous layer was acidified with 1.2 N HCl and extracted with EtOAc. The EtOAc solution was washed with brine, dried and concentrated to furnish the title compound which was used in the next step without purification.  $^1\text{H}$  NMR: (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.92 (d, 6H), 1.51 (m, 1H), 1.72 (m, 1H), 1.98 (m, 1H), 3.67 (m, 1H), 7.0-7.4 (m, 5H).
- 10 Step B N-Methoxy-N-methyl-4-methyl-2-phenylpentanamide  
To a solution of 0.234 g (1.22 mmol) of 4-methyl-2-phenylpentanoic acid in 6 mL  $\text{CH}_2\text{Cl}_2$  and 2 drops of DMF, 0.12 mL (1.34 mmol) of oxalyl chloride was added. The solution was stirred for 1 h and concentrated. The residue was dissolved in 1 mL  $\text{CH}_2\text{Cl}_2$  and added to a mixture of 0.142 g N,O-dimethylhydroxylamine hydrochloride in 4 mL  $\text{CH}_2\text{Cl}_2$  and 4 mL saturated  $\text{NaHCO}_3$ . After stirring for 4 h, the layers were separated and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$ . The combined  $\text{CH}_2\text{Cl}_2$  layer was washed with brine, dried and concentrated to give the title compound which was used in the next step without purification.  $^1\text{H}$  NMR: (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.94 and 0.96 (2d, 6H), 1.5 (m, 1H), 1.67 (m, 1H), 2.0 (m, 1H), 3.19 (s, 3H), 3.54 (s, 3H), 4.18 (br, 1H), 7.2-7.4 (m, 5H).
- 20 Step C 5-Methyl-3-phenyl-2-hexanone  
To a solution of 75 mg (0.317 mmol) N-methoxy-N-methyl-4-methyl-2-phenylpentanamide in 1 mL dry THF, 0.45 mL 1.4 M methylmagnesium bromide was added. The reaction was stirred for 1 h, quenched with 1.2 N HCl and extracted with EtOAc. The EtOAc solution was washed with brine, dried and concentrated leaving the title compound.  $^1\text{H}$  NMR: (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.95 (2d, 6H), 1.42 (m, 1H), 1.67 (m, 1H), 1.9 (m, 1H), 2.06 (s, 3H), 3.73 (m, 1H), 7.0-7.4 (m, 5H).
- 25 Step D 5-Methyl-3-phenyl-2-hexanol  
A solution of 66 mg (0.345 mmol) of 5-methyl-3-phenyl-2-hexanone in 1 mL MeOH was treated with 16 mg sodium borohydride. After 1.5 h, the reaction was quenched with 1.2 N HCl and concentrated. The residue was partitioned between EtOAc and water. The organic layer was washed with brine, dried and concentrated to yield the crude title compound which was used without purification.  $^1\text{H}$  NMR: (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.88 (2d, 6H), 1.0-1.8 (m, 4H), 1.2 (d, 3H), 2.64 (m, 1H), 3.9 (m, 1H), 7.2-7.4 (m, 5H).
- 35

Step E 2-Azido-5-methyl-3-phenylhexane.

To a solution of 60 mg 5-methyl-3-phenyl-2-hexanol in 2 mL CH<sub>2</sub>Cl<sub>2</sub>, 0.163 g (0.62 mmol) of triphenylphosphine and 96 mg (0.31 mmol) of zinc azide pyridine were added. The reaction mixture was cooled in an ice bath and 98 mL (0.62 mmol) of DEAD was added. The cold bath was removed and the solution was stirred for 3 h. The reaction mixture was filtered through a pad of CELITE diatomaceous earth and the pad was rinsed with CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated and the residue was purified by prep-TLC using 20% EtOAc-hexane to isolate the title compound. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 0.88 (2d, 6H), 1.12 (d, 3H), 1.31 (m, 1H), 1.72 (m, 2H), 2.68 (m, 1H), 3.53 (m, 1H), 7.2-7.4 (m, 5H).

Step F 2-Amino-5-methyl-3-phenylhexane.


To a solution of 32 mg 2-azido-5-methyl-3-phenylhexane in 1 mL MeOH and 2 drops of 1.2 N HCl, 4 mg PtO<sub>2</sub> was added and the solution was stirred under H<sub>2</sub> atmosphere for 2 h. The reaction was filtered through a pad of CELITE diatomaceous earth and the pad was rinsed with MeOH. The combined filtrate was concentrated to give the desired product. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 0.86 (m, 6H), 0.99 (d, 3H), 1.25 (m, 1H), 1.54 (m, 1H), 1.77 (m, 1H), 2.73 (m, 1H), 3.19 (m, 1H), 7.2-7.4 (m, 5H).

REFERENCE EXAMPLE 39N-[3-(4-Chlorophenyl)-2-(3,5-difluorophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer α)

The title compounds was prepared following the procedures described for Reference Example 2 substituting methyl phenylacetate with methyl 3,5-difluorophenylacetate (prepared from 3,5-difluorophenylacetic acid and trimethylsilyldiazomethane) at Step

A and sodium borohydride in MeOH with lithium tri(*sec*-butylborohydride in THF at Step E. LC-MS:  $m/e$  296 ( $M + H$ )<sup>+</sup> (2.39 min).

5


REFERENCE EXAMPLE 40

N-[2-(3-Bromophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

10

The title compounds was prepared following the procedures described for Reference Example 2 substituting methyl phenylacetate with methyl 3-bromophenylacetate (prepared from 3-bromophenylacetic acid and trimethylsilyldiazomethane) at Step A and sodium borohydride in MeOH with lithium tri(*sec*-butylborohydride in THF at Step E. LC-MS:  $m/e$  338 ( $M + H$ )<sup>+</sup> (2.5 min).

15

REFERENCE EXAMPLE 41

N-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )




Step A      2-(*N*-*tert*-Butoxycarbonyl)amino-4-(4-chlorophenyl)-3-(3-cyanophenyl)butane

To a solution of 2-(*N*-*tert*-butoxycarbonyl)amino-3-bromophenyl-4-(4-chlorophenyl)butane (1.0 g, 2.3 mmol) in 5 mL DMF was added zinc cyanide (0.16 g, 1.4 mmol), tris(dibenzylidene-acetone)dipalladium chloroform complex (3.0 mg, 2.8  $\mu$ mol), 1,1'-bis(diphenylphosphino)ferrocene (5.0 mg, 9.0  $\mu$ mol) and water (0.1 mL). After heating at 120°C for 6 h under nitrogen, another batch of zinc cyanide (0.16 g, 1.4 mmol), tris(dibenzylideneacetone)dipalladium chloroform complex (5.0 mg, 4.8  $\mu$ mol), 1,1'-bis(diphenylphosphino)ferrocene (5.0 mg, 9.0  $\mu$ mol) and water (0.05 mL) was added, and heating was continued for another 18 h. After cooling to room temperature, the resulting mixture was partitioned between water (50 mL) and ether (50 mL). The organic layer was separated and the aqueous layer extracted with ether (2 x 50 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated, and the residue was purified by flash column chromatography on silica gel eluted with 20% EtOAc in hexane to afford the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.6-7.3 (m, 4H), 7.10 (d, 2H), 6.92 (d, 2H), 3.88 (m, 1H), 3.20 (m, 1H), 2.97 (m, 1H), 1.82 (m, 1H), 1.45 (s, 9H), 0.94 (d, 3H). LC-MS: *m/e* 385 (M + H)<sup>+</sup> (3.9 min).

Step B      *N*-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 2, Step I. LC-MS: *m/e* 285 (M + H)<sup>+</sup> (2.2 min).

REFERENCE EXAMPLE 42



*N*-[2-(3-Chlorophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride

(Diastereomer  $\alpha$ )

Step A      2-(*N*-*tert*-Butoxycarbonyl)amino-4-(4-chlorophenyl)-3-(3-trimethylstannylphenyl)butane

To a solution of 2-(*N*-*tert*-butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)butane (intermediate of Reference Example 40, 1.5 g, 3.4 mmol) in 15 mL anhydrous dioxane was added hexamethylditin (1.6 g, 4.8 mmol), triphenylphosphine (18 mg, 0.068 mmol), lithium chloride (0.16 g, 3.8 mmol) and tetrakis(triphenyl-phosphine)palladium (0.20 g, 0.17 mmol). After heating at 95°C for 7.5 h under nitrogen, the reaction mixture was cooled to room temperature, diluted with EtOAc (100 mL), washed with 10% aqueous potassium fluoride and brine, dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated to dryness. The residue was purified by flash column chromatography on silica gel eluted with 20% EtOAc in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.3-7.2 (m, 2H), 7.07 (d, J=8.5 Hz, 2H), 7.06-6.99 (m, 2H), 6.86 (d, J=8.5 Hz, 2H), 3.93 (m, 1H), 3.18 (m, 1H), 2.76 (m, 2H), 1.51 (s, 9H), 0.94 (d, J=7.0 Hz, 3H), 0.21 (s, 9H).


Step B      2-(*N*-*tert*-Butoxycarbonyl)amino-3-(3-chlorophenyl)-4-(4-chlorophenyl)butane

To a solution of 2-(*N*-*tert*-butoxycarbonyl)amino-4-(4-chlorophenyl)-3-(3-trimethylstannylphenyl)butane (0.55 g, 1.0 mmol) in 5 mL CH<sub>2</sub>Cl<sub>2</sub> at 0°C was added *tert*-butoxychloride (freshly prepared, 0.20 mL, 1.1 mmol). The reaction was allowed to warm to room temperature over 2 h, and the resulting mixture was concentrated with 2 g silica gel. The residue was purified by flash column chromatography on silica gel eluted with 10% ether in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.25-7.15(m, 2H), 7.11 (d, J=8.5 Hz, 2H), 7.09 (m, 1H), 6.99 (d, J=7.5 Hz, 1H), 6.92 (d, J=8.5 Hz, 2H), 3.88 (m, 1H), 3.19 (dd, J=13.0, 3.5 Hz, 1H), 2.90-2.75 (m, 2H), 1.50 (s, 9H), 0.94 (d, J=6.5 Hz).

Step C      *N*-[2-(3-Chlorophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 2, Step I. LC-MS: m/e 294 (M + H)<sup>+</sup> (2.82 min).

#### REFERENCE EXAMPLE 43




N-[2-(3-Bromophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride and N-[3-(4-Chlorophenyl)-2-(3-iodophenyl)-1-methylpropyl]amine hydrochloride (1:1 mixture) (Diastereomer  $\alpha$ )

- 5 Step A 2-(N-tert-Butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)-butane and 2-(N-tert-Butoxycarbonyl)amino-4-(4-chlorophenyl)-3-(3-iodophenyl)butane

To a solution of 2-(N-tert-butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)butane (intermediate of Reference Example 40, 2.6 g, 5.9 mmol) in 7 mL anhydrous THF at 0°C was added methylmagnesium chloride (3 M in THF, 3.9 mL, 12 mmol). After 30 min, the reaction mixture was cooled to -78°C, and was added tert-butyllithium (1.7 M, 10 mL, 17 mmol). After stirring at -78°C for 2 h, the reaction was allowed to warm to 0°C, and half of the resulting mixture was added to a suspension of iodine (5.0 g, mmol) in 10 mL THF at -40°C. The reaction mixture was allowed to warm to room temperature over 2 h, and was partitioned between ether (100 mL) and saturated aqueous ammonium chloride (100 mL). The organic layer was separated and the aqueous layer extracted with ether (2 x 50 mL). The combined extracts were washed with dilute aqueous sodium thiosulfate (2x) and brine, dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated to dryness. The residue was purified by flash column chromatography on silica gel eluted with 10% EtOAc in hexane to afford the title compounds as a 1:1 mixture.

- 25 Step B N-[2-(3-Bromophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride and N-[3-(4-chlorophenyl)-2-(3-iodophenyl)-1-methylpropyl]amine hydrochloride (1:1 mixture) (Diastereomer  $\alpha$ )

The title compound was prepared following procedure described for Reference Example 2, Step I. LC-MS: m/e 338/386/ (M + H)<sup>+</sup> (2.6 min).

REFERENCE EXAMPLE 445 2-Amino-4-(4-chlorophenyl)-3-cyclobutylmethoxybutane.Step A Methyl 2-diazo-3-(4-chlorophenyl)propanoate.

DL-4-Chlorophenylalanine methyl ester (5.0 g, 23.36 mmol) was dissolved in 120 mL chloroform and placed into an oven-dried 3-neck flask equipped with a condenser and an addition funnel. Glacial acetic acid (0.267 mL, 4.672 mmol) was added. Finally,  
 10 isoamylnitrite (3.8 mL, 28 mmol) was added dropwise while slowly bringing the reaction to reflux (73°C). The reaction was refluxed for 30 minutes and then cooled to 0°C. The reaction mixture was washed with cold 1 N sulfuric acid solution, cold water, cold saturated aqueous sodium bicarbonate solution, and then cold water again. The organic extracts were dried over MgSO<sub>4</sub>, filtered and concentrated under reduced  
 15 pressure. The crude mixture was purified by flash chromatography (Biotage 40M cartridge, gradient elution using hexane and EtOAc (100:1 to 50:1) to provide a yellow oil, homogeneous by TLC, R<sub>f</sub>=0.48 (4:1 hexanes:EtOAc). 500 MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 3.65 (s, 2H); 3.83 (s, 3H); 7.22 (d, J=8.5 Hz, 2H), 7.34 (d, J=8.5, 2H).

Step B Methyl 3-(4-chlorophenyl)-2-cyclobutylmethoxypropanoate.

20 To a solution of 500 mg (2.23 mmol) of methyl-2-diazo-3-(4-chlorophenyl)propanoate (obtained from Step A) and 1.05 mL (5 eq; 11.1 mmol) of cyclobutanemethanol in 5 mL benzene in a pressure tube was added 10 mg (1 mole %) of Rh<sub>2</sub>(OAc)<sub>4</sub> catalyst. The tube was sealed and heated to 90°C for 1.5 h. The solvents were evaporated under reduced pressure and the crude material was taken up  
 25 in CH<sub>2</sub>Cl<sub>2</sub> and purified by flash chromatography via gradient elution using mixtures of hexane and EtOAc (100:1 to 50:1). This provided the title compound as a clear oil. TLC R<sub>f</sub>=0.53 (4:1 hexanes:EtOAc). 500 MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.68 (m, 2H);

1.85 (m, 1H); 1.88 (m, 1H); 2.01 (m, 2H); 2.53 (sep, 1H); 2.98 (m, 2H); 3.24 (dd, 1H); 3.58 (dd, 1H); 3.76 (s, 3H); 3.98 (dd, 1H); 7.20 (d, 2H); 7.28 (d, 2H).

Step C      4-(4-Chlorophenyl)-3-cyclobutylmethoxybutan-2-one.

At 0°C, under anhydrous conditions, to a stirred suspension of N,O-dimethylhydroxylaminehydrochloride (732 mg, 7.50 mmol) in 60 mL CH<sub>2</sub>Cl<sub>2</sub> was added dimethylaluminum chloride (7.5 mL, 1M solution in hexanes). The solution was allowed to warm to room temperature over a period of one hour. At that point a solution of methyl 2-cyclobutylmethoxy-3-(4-chlorophenyl) propanoate (531 mg, 1.88 mmol, obtained from Step B) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL) was added dropwise. The reaction was allowed to stir overnight at room temperature when TLC indicated completion of reaction. The reaction was worked up by the addition of pH=8 phosphate buffer (25 mL, approx. 3 mL/mmol of Me<sub>2</sub>AlCl) and allowed to stir at room temperature for 30 minutes, diluted with chloroform (75 mL), and the phases were separated. The organic layer was washed with water and dried over MgSO<sub>4</sub>. The solvents were evaporated under reduced pressure and the crude product was purified by flash chromatography (gradient elution using hexane and EtOAc, 20:1 to 5:1) to give the Weinreb amide as a clear oil. This purified material (424 mg, 1.36 mmol) was dissolved in 10 mL THF, injected into an oven dried flask, and cooled to 0°C under nitrogen. Methyl magnesium bromide (1.4 mL 3M solution in ether) was added to the solution dropwise. The reaction was allowed to warm to room temperature. After 4 h the TLC indicated a complete reaction. The reaction was quenched with enough 10% citric acid to bring the pH of the solution to approximately 3. The aqueous layer was extract with ether. The combined organics were washed with water and then dried over MgSO<sub>4</sub>. The solvents were evaporated under reduced pressure and the crude material was purified by flash chromatography (hexane:EtOAc, 100:1 to 50:1), resulting in 250 mg the title compound as a clear oil. TLC R<sub>f</sub>=0.55 (4:1 hexanes:EtOAc). 500 MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.71 (m, 2H); 1.84 (m, 1H); 1.91 (m, 1H); 2.01 (m, 2H); 2.17 (s, 3H); 2.53 (sep, 1H); 2.90 (m, 2H); 3.28 (dd, 1H); 3.43 (dd, 1H); 3.81 (dd, 1H).


Step D      2-Amino-4-(4-chlorophenyl)-3-cyclobutylmethoxybutane.

A solution of 3-cyclobutylmethoxy-4-(4-chlorophenyl)butan-2-one (247 mg, 0.925 mmol, obtained from Step C) in 0.5 mL CH<sub>2</sub>Cl<sub>2</sub> was added to a stirred suspension of NH<sub>4</sub>OAc (715 mg, 9.25 mmol) and NaBH<sub>3</sub>CN (35 mg, 0.555 mmol) at room temperature and allowed to stir overnight. The reaction was quenched by the addition of 2.2 mL conc. HCl allowed to stir for 30 minutes. The solvents were evaporated

under reduced pressure and the residue was partitioned between ether and water. The aqueous layer was washed two more times with ether. The combined organics were dried over  $\text{Na}_2\text{SO}_4$ . The crude product mixture obtained after filtration and removal of volatiles was purified by flash chromatography, eluting using mixtures of mixtures of  $\text{CH}_2\text{Cl}_2$  and MeOH (100%  $\text{CH}_2\text{Cl}_2$ , to 5% MeOH in  $\text{CH}_2\text{Cl}_2$ ) to provide the title compound as a yellow oil, homogeneous by TLC  $R_f=0.12$  (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ).  
5 500 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.16 (t, 3H); 1.67 (m, 2H); 1.85 (m, 3H); 2.01 (m, 2H); 2.48 (m, 1H); 2.74 (m, 2H); 2.90 (dd, 1H); 3.15 (d quint, 2H); 3.37 (m, 2H).

10 2-Amino-4-(4-chlorophenyl)-3-methoxy-butane, 2-amino-4-(4-chlorophenyl)-3-ethoxy-butane, 2-amino-4-(4-chlorophenyl)-3-n-propyloxy-butane, 2-amino-4-(4-chlorophenyl)-3-n-pentyloxy-butane, and 2-amino-4-(4-chlorophenyl)-3-cyclopentylmethoxy-butane were prepared according to the procedures described in Reference Example 44 substituting an appropriate alcohol for cyclobutylmethanol in  
15 Step B.

#### REFERENCE EXAMPLE 45



#### 2-Amino-4-(4-chlorophenyl)-3-(1-pyrrolidinyl)-butane hydrochloride.

##### Step A Ethyl 3-(4-chlorophenyl)-2-pyrrolidin-N-yl-propanoate.

20 While stirring rapidly, to a mixture of DL-4-chlorophenylalanine methyl ester hydrochloride (2.5 g, 10 mmole), 40 mL ethanol and sodium carbonate (3.18 g, 30 mmole) was added dropwise a solution of 1,4-dibromobutane (2.16 g, 10 mmol) dissolved in 20 mL ethanol. The mixture was refluxed overnight. The volatiles were removed under reduced pressure, and the residue was partitioned between water and  
25 EtOAc. The aqueous layer was re-extracted with EtOAc thrice. The organic layers were combined and washed with water and brine and dried over anhydrous  $\text{MgSO}_4$ . The crude product obtained after filtration and removal of volatiles was purified via flash chromatography using mixtures of  $\text{CH}_2\text{Cl}_2$  and MeOH to provide the titled

compound as an oil, homogeneous by TLC,  $R_f = 0.55$  in 95:5  $\text{CH}_2\text{Cl}_2$ : MeOH.

LC/MS  $m/e = 282.1$  ( $M+1$ ). 400 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.12(t,  $J = 7.2$  Hz, 3H), 1.72 (m, 4H), 2.67 (m, 1H), 2.76(m, 1H), 3.05 (m, 4H), 3.43 (m, 1H), 4.05 (m, 2H), 7.13 (d,  $J = 8.2$  Hz, 2H), 7.24 (d,  $J = 8.2$  Hz, 2H)

5    Step B    4-(4-Chlorophenyl)-3-(1-pyrrolidinyl)-butan-2-one.

The title compound was prepared according to the procedure of Reference Example 2, Step C except that ethyl 3-(4-chlorophenyl)-2-(1-pyrrolidinyl)-propanoate (from Step A) was the ester used (two steps). TLC  $R_f = 0.7$  (95:5  $\text{CH}_2\text{Cl}_2$ : MeOH). LC/MS  $m/e = 252$  ( $M+1$ ). 500 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.86(br s, 4H), 2.03 (s, 3H), 2.66 (m, 2H), 2.78 (m, 2H), 2.98 (dd,  $J=2.9, 10.3$  Hz, 1H), 3.08 (m, 1H), 3.43 (m, 1H), 7.12 (d,  $J = 8.3$  Hz, 2H), 7.26 (d,  $J = 8.3$  Hz, 2H)

Step C    4-(4-Chlorophenyl)-3-pyrrolidin-N-yl-butan-2-one oxime.


To a solution of 4-(4-chlorophenyl)-3-pyrrolidin-N-yl-butan-2-one (200 mg, 0.79 mmol, from Step B) dissolved in ethanol (2 mL), was added pyridine (63 mg, 0.79 mmol), and hydroxylamine hydrochloride (78 mg, 1.12 mmol). The mixture was refluxed for 24h when LC/MS indicated disappearance of all starting material. The mixture was cooled to room temperature, concentrated under reduced pressure, treated with 33% aqueous potassium carbonate, and extracted with chloroform 5 times. The organic layers were combined and filtered over glass wool and dried over potassium carbonate. The filtrate obtained after passing through sintered glass was concentrated to give the oxime, homogeneous by TLC,  $R_f = 0.3$  in 95:5  $\text{CH}_2\text{Cl}_2$ : MeOH. LC/MS  $m/e = 267$  ( $M+1$ ). 500 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.73(m, 4H), 1.76 (s, 3H), 2.40 (m, 2H), 2.60 (m, 2H), 2.72 (dd,  $J=2.7, 10.8$  Hz, 1H), 2.94 (dd,  $J=4.3, 8.8$  Hz, 1H), 3.03 (dd,  $J = 4.4, 13.3$ Hz, 1H), 3.8 (s, 1H), 6.96 (d,  $J = 8.3$  Hz, 2H), 7.11 (d,  $J = 8.3$  Hz, 2H)

Step D    2-Amino-4-(4-chlorophenyl)-3-pyrrolidin-N-yl-butane hydrochloride.

At room temperature, to a solution of 4-(4-chlorophenyl)-3-pyrrolidin-N-yl-butan-2-one oxime (173 mg, 0.648 mmol, from Step C) in 1.8 mL anhydrous THF was added dropwise a 1M solution of lithium aluminum hydride in THF (0.778 mmole). The mixture was refluxed for 20 h. The reaction was quenched by addition of saturated aqueous sodium sulfate (0.1 mL), and stirred overnight. This mixture was filtered over a pad of CELITE diatomaceous earth, and the filtrate was concentrated to dryness. The mass spectrum of this material looked very messy, so the HCl salt was prepared (by addition of a HCl(g) in ether solution) in attempt to clean up the mess.

By NMR, the reductive amination provided a ~1:1 mixture of the two diastereomeric pairs of amines. This HCl salt was rather sticky and difficult to work with and therefore was used in the ensuing coupling experiment without further purification. LC/MS  $m/e = 253$  ( $M+1$ ). 500 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  1.56, 1.59 (2 d,  $J = 7.2$  Hz, 3H), 2.03 (m, 6H), 2.08 (m, 2H), 3.20-4.00 (m, 3H), 7.43 (m, 4H)

#### REFERENCE EXAMPLE 46



#### 10 Benzyl 3-amino-2-(4-chlorobenzyl)butyrate.

##### Step A Benzyl 2-(4-chlorobenzyl)-3-ketobutyrate.

Benzyl acetoacetate (1.92 g, 10 mmole) and 4-chlorobenzylbromide (2.05 g, 10 mmole) were dissolved in 40mL anhydrous THF and cooled to  $-10^\circ\text{C}$ . To this mixture was added dropwise slowly a solution of solution of sodium hexamethyl  
 15 disilazide (0.5M solution in THF). Monoalkylation occurred almost exclusively of bisalkylation between  $-10$  and  $5^\circ\text{C}$ . After quenching with water, the organics were extracted with EtOAc three times. The combined organic layer was washed with brine and dried over anhydrous  $\text{MgSO}_4$ . The crude product obtained after filtration and removal of volatiles was purified via flash chromatography using gradient elution  
 20 (mixtures of hexane and EtOAc) to provide of the title compound as a clear yellow liquid, homogeneous by TLC,  $R_f=0.4$  in 4:1 hexane:EtOAc. By NMR, this compound, this compound exists in a ~4:1 ratio of the keto:enol forms. 400 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.08, 2.18 (2 s, 3H), 3.15 (m, 2H), 3.80 (t,  $J = 7.5$  Hz, 0.8 H), 5.14, 5.17 (2 s, 2H), 7.05-7.39 (m, 9H).


##### 25 Step B Benzyl 3-amino-2-(4-chlorobenzyl)butyrate.

Benzyl 2-(4-chlorobenzyl)-3-ketobutyrate (317 mg, 1 mmole, obtained from Step A) was added to a cooled mixture of 7M ammonia in MeOH (2.42 mL) and glacial acetic



acid (1.6 mL). To this solution, at  $\sim 10^{\circ}\text{C}$ , was added sodium cyanoborohydride (101 mg, 1.75 mmol) in small portions. This mixture was stirred at room temperature for 40 h. The excess sodium cyanoborohydride was destroyed by the addition of 6M HCl (to pH 1). The residue obtained after removal of volatiles was taken up in a minimal amount of water and extracted with ether. The aqueous layer was basified to pH 10 using solid KOH. This layer was then saturated with sodium chloride and then extracted with EtOAc. Further analyses of the ether and the EtOAc layers suggest that the desired product resides the EtOAc layer. This material was used in the ensuing coupling reaction without further purification. Proton NMR spectrum show that the two pairs of diastereomers are obtained in  $\sim 1:1$  ratio, homogeneous by TLC,  $R_f = 0.4$  in 95:5  $\text{CH}_2\text{Cl}_2$  : MeOH. LC/MS  $m/e = 318$  ( $M+1$ ). 400 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.27, 1.29 (2 d,  $J=7\text{Hz}$ , 3H), 2.85 (m, 1H), 3.03 (m, 1H), 3.15 (m, 1H), 3.55 (m, 1H), 4.85 (br, 2H), 5.00-5.18 (m, 2H), 7.0-7.2 (m, 9H).

#### REFERENCE EXAMPLE 47



#### 2-Amino-4-(4-chlorophenyl)-3-cyclopentylbutane.

##### Step A      Methyl 3-(4-chlorophenyl)-2-cyclopentylpropanoate.

A mixture of methyl cyclopentylacetate (3.52 g, 25 mmol) and 4-chlorobenzyl bromide (4.75 g, 23 mmol) was dissolved in 100 mL THF in an oven-dried flask. The solution was cooled to  $-40^{\circ}\text{C}$  and 23 mL 1M NaHMDS solution in hexanes was added slowly over an hour while maintaining the temperature at  $-40^{\circ}\text{C}$ . The solution was then stirred for an additional 3 h at  $-40^{\circ}\text{C}$ . The reaction was quenched at  $-40^{\circ}\text{C}$  with enough 10% citric acid solution to bring the pH to  $\sim 3.5$ . The aqueous layer was extracted with ether three times. The combined organics were washed with water and dried over  $\text{MgSO}_4$ . The solvents were evaporated under reduced pressure and the crude material was purified by flash chromatography [Biotage 40 M, gradient elution using mixtures of hexane and EtOAc (from 0 – 1% EtOAc)]. This provided a light brown oil, which is a 3:1 ratio of the title compound : methyl cyclopentylacetate

based on the methyl ester peak integrations. TLC of the desired product:  $R_f=0.34$  in 20:1 hexane:EtOAc. The complete separation of the title compound from the starting material was not practical in this case, as they had overlapping  $R_f$ 's on the TLC.

Therefore, this mixture was carried on to the next step.

5    Step B        3-(4-Chlorophenyl)-2-cyclopentylpropanoic acid.

The mixture of methyl esters from Step A (3.41 g, 14.48 mmol of methyl 3-(4-chlorophenyl)-2-cyclopentylpropanoate--assuming 3:1 mixture obtained in Step A.) was dissolved in 10 mL DMSO and 4 mL distilled water. Then powdered KOH (3.25 g, 57.92 mmol) was added and the solution was stirred overnight at room temperature.

10    The next day the pH was brought to 2 with 2 N HCl. The aqueous layer was extracted 3 times with ether. The combined organic extracts were dried over anhydrous sodium sulfate. Filtration and evaporation of volatiles provided the mixture of acids as an oil. 500 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.28 (m, 2H), 1.64 (m, 6H), 2.06 (m, 1H), 2.47 (m, 1H), 2.86 (t, 2H).

15    Step C        3-(4-Chlorophenyl)-2-cyclopentyl -N, O-dimethyl-propanamide.

The mixture of acids obtained in Step B (3.21 g, 14.48 mmol of the desired acid--based on assumption of 3:1 mixture from Step B) was dissolved in 75 mL  $\text{CH}_2\text{Cl}_2$ .

While being stirred rigorously, *N,O*-dimethylhydroxylamine hydrochloride (1.56 g, 15.95 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (3.06 g, 16.0 mmol),  
20    diisopropylethylamine (5.56 mL, 31.90 mmol), and a catalytic amount of 4-(dimethylaminopyridine) were added sequentially. Stirring was continued overnight at room temperature. The next day the reaction mixture was diluted with EtOAc, treated with water, and the phases were separated. The aqueous layer was re-extracted with EtOAc twice. The combined organic layers were washed with water  
25    three times and then with saturated brine. The organic layer was dried over  $\text{MgSO}_4$ , filtered, and the solvents were removed under reduced pressure. The crude material was purified by flash chromatography [Biotage 40 M column, gradient elution using mixtures of hexanes and EtOAc (100:1 to 20:1)] to provide the title compound cleanly as an oil. TLC  $R_f=0.31$  (4:1 hexanes:EtOAc). LC/MS  $m/e$  295.9 ( $M+1$ ). 500 MHz  
30     $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.27(m, 2H), 1.64 (m, 6H), 1.97 (m, 1H), 2.13 (q, 1H), 2.81 (d, 1H), 2.97 (d, 1H), 3.07 (s, 3H), 3.17 (s, 3H). LC/MS  $m/e$  295.9 ( $M+1$ ).

Step D        4-(4-Chlorophenyl)-3-cyclopentylbutan-2-one.

3-(4-Chlorophenyl)-2-cyclopentyl -N, O-dimethyl-propanamide (514 mg, 1.737 mmol, obtained from Step C) was dissolved in 15 mL anhydrous THF and injected

into an oven dried flask under nitrogen. The solution was cooled to 0°C and CH<sub>3</sub>MgBr (1 M in ether) was added dropwise. The ice bath was removed and the reaction was allowed to warm to room temperature and stirred for a total of 4h. TLC indicated a nearly complete reaction. The reaction was quenched with enough 10 %  
 5 citric acid to bring the pH of the solution to 3. The aqueous layer was extracted 3 times with ether and the extracts were dried over anhydrous MgSO<sub>4</sub>. The solution was filtered and the solvents were removed under reduced pressure. The crude material was purified by flash chromatography (30 mL silica; 100:1 to 50:1 hexanes: EtOAc) to provide 351 mg the title compound as an oil. TLC R<sub>f</sub>=0.49 (4:1 hexanes:  
 10 EtOAc). 500 MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.23 (m, 3H), 1.58 (m, 1H), 1.71 (m, 3H), 1.91 (s, 3H), 1.93 (m, 1H), 2.05 (m, 1H), 2.68 (m, 1H), 2.84 (m, 2H).


Step E 2-Amino-4-(4-chlorophenyl)-3-cyclopentylbutane.

The title compound was prepared according to the procedure of Reference Example 2, Step D, except that 4-(4-chlorophenyl)-3-cyclopentylbutan-2-one (obtained from  
 15 Step D) was used as the starting material. LC/MS m/e 251.9 (M+1); 500 MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.93 (m, 1H), 1.29 (q, 3H), 1.29 (m, 2H), 1.61 (m, 4H), 1.87 (m, 3H), 2.62 (m, 1H), 2.80 (m, 1H), 3.26 and 3.48 (m, 1H).

20 2-Amino-4-(4-chlorophenyl)-3-ethyl-butane and 2-amino-4-(4-chlorophenyl)-3-isopropyl-butane were also prepared according to the procedures described in Reference Example 47 substituting the appropriate ester for methyl cyclopentylacetate in Step A.

REFERENCE EXAMPLE 48

25



2-Amino-3-(1-(1,2,3-triazolyl))-4-(4-chlorophenyl)butane:

Step A Benzyl 2-(1-(1,2,3-triazolyl))acetate:

A mixture of 1,2,3-triazole (2.07 g, 30 mmol), benzyl bromoacetate (6.9 g, 30 mmol),  
 30 and diisopylethylamine (5.1 mL, 30 mmol) in 40 mL CH<sub>2</sub>Cl<sub>2</sub> was stirred

overnight at room temperature. This mixture was then diluted with ether until no further precipitate formed. The solid was filtered and washed with ether. The filtrate was concentrated and the residue was purified on silica gel using 10% hexane in CH<sub>2</sub>Cl<sub>2</sub> to give the title compound's isomer, benzyl 2-(2-(1,2,3-triazolyl))acetate as

5 amorphous solid. Further elution with a solvent mixture containing equal amounts of ether and CH<sub>2</sub>Cl<sub>2</sub> gave the title compound as amorphous solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 2.251(s, 2H), 7.267-7.390(m, 5H), 7.723(s, 1H), 7.785(s, 1H)

Step B 2-(1-(1,2,3-triazolyl))acetic acid:

Palladium hydroxide (20% on carbon, 800 mg) was added to a solution of benzyl 2-(1-(1,2,3-triazolyl))acetate (Step A, 8.68 g, 39.9 mmol) in 150 mL MeOH and the mixture was hydrogenated overnight on a Parr shaker under an atmosphere of hydrogen at room temperature and 45 psi. The catalyst was filtered through a bed of CELITE diatomaceous earth and washed with MeOH. The filtrate was concentrated to give a solid, which was dried in vacuo at 50°C for 36 h resulting in the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 5.3 (s, 2H), 7.75 (s, 1H), 8.016 (s, 1H).

Step C N-Methoxy-N-methyl-2-(1-(1,2,3-triazolyl))acetamide:

Oxalyl chloride (0.95 mL, 11 mmol) was added dropwise to a suspension of 2-(1-(1,2,3-triazolyl))acetic acid (Step B, 1.27 g, 10 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> containing 0.05 mL DMF. Vigorous effervescence was observed. This mixture was stirred at room temperature for 4 h and cooled to -78°C. A solution of N,O-dimethylhydroxylamine hydrochloride (1.2 g, 13 mmol) and diisopropylethyl amine (6.0 mL, 35 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> was added slowly over 3 min. The mixture was then allowed to warm to room temperature and stirred overnight. The reaction mixture was then diluted with ether until no additional precipitate appeared. The solid was filtered and washed with ether. The filtrate was concentrated and the residue was purified on silica gel using EtOAc as solvent to provide the title compound as amorphous solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 3.252 (s, 3H), 3.812 (s, 3H), 5.379 (s, 2H), 7.753 & 7.761 (s's, 2H).

Step D N-Methoxy-N-methyl-3-(4-chlorophenyl)-2-(1-(1,2,3-triazolyl))propionamide

30 Lithium hexamethyldisilazide (1 molar in THF, 8.4 mL, 8.4 mmol) was added dropwise to a solution of N-methoxy-N-methyl-2-(1-(1,2,3-triazolyl))acetamide (Step C, 1.19 g, 7 mmol) in 15 mL THF at -78°C. After additional 30 min stirring, a solution of 4-chlorobenzyl bromide (1.65 g, 8 mmol) in 5 mL THF was added dropwise. The mixture was allowed to warm to room temperature and stirred 5.5 h.

This mixture was purified on silica gel using 40% EtOAc in hexane to give the title compound.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.186 (s, 3H), 3.234-3.267 (m, 1H), 3.453-3.506 (m, 1H), 3.582 (s, 3H), 6.145-6.188 (m, 1H), 7.048-7.279 (m, 4H), 7.726 (s, 1H), 7.954 (s, 1H).

5    Step E    2-Azido-3-(1-(1,2,3-triazolyl))-4-(4-chlorophenyl)butane:

The product of Step D, N-methoxy-N-methyl-3-(4-chlorophenyl)-2-(1-(1,2,3-triazolyl)propionamide was converted to the title compound following the procedures described in Reference Example 2, Step D-E and Reference Example 5, Step D.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.219-1.246 (d's 3H), 3.253-4.754 (m, 4H), 6.866-7.299

10 (d's, 4H), 7.313, 7.618, 7.63, & 7.706 (s's, 2H).


Step F    2-Amino-3-(1-(1,2,3-triazolyl))-4-(4-chlorophenyl)butane:

Platinum oxide (14 mg) was added to a solution of 2-azido-3-(1-(1,2,3-triazolyl))-4-(4-chlorophenyl)butane (Step E, 138 mg, 0.5 mmol) in 4 mL MeOH. This mixture was hydrogenated in an atmosphere of hydrogen using a hydrogen filled balloon for 3

15 h at room temperature. The catalyst was filtered through a bed of CELITE diatomaceous earth and washed with MeOH. The filtrate was concentrated to give the title compound as oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.085-1.174 (d's 3H), 3.220-3.361 (m, 2H), 3.517-3.563 (m, 1H), 4.379-4.431 (m, 1H), 6.679-7.179 (d's, 4H), 7.297, 7.40, 7.592 & 7.607 (s's, 2H).

20

REFERENCE EXAMPLE 49




2-Amino-3-(1-(1,2,4-triazolyl))-4-(4-chlorophenyl)butane:

25    The title compound was prepared according to the procedures described in Reference Example 48 substituting 1,2,4-triazole for 1,2,3-triazole in Step A. The azide was separated by column chromatography on silica gel eluted with 20% hexane in EtOAc.

REFERENCE EXAMPLE 50

30



*N*-[3-(4-Chlorophenyl)-2-(3-methylphenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )


Step A      2-(*N*-*tert*-Butoxycarbonyl)amino-4-(4-chlorophenyl)-3-(3-  
5                    methylphenyl)butane

A mixture of 2-(*N*-*tert*-butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)butane (intermediate of Reference Example 40, 0.50 g, 1.1 mmol), tetramethyltin (0.41 g, 2.3 mmol), triphenylphosphine (0.12 g, 0.46 mmol), lithium chloride (0.38 g, 9.1 mmol) and dichlorobis(triphenylphosphine)palladium (0.12 g, 0.17 mmol) in 20 mL anhydrous DMF was heated at 100°C under nitrogen for 18 h. The reaction mixture was cooled to room temperature, and was partitioned between water (100 mL) and ether (100 mL). The organic layer was separated and the aqueous layer was extracted with ether (100 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated to dryness, and the residue was purified  
15 by flash column chromatography on silica gel eluted with 10% EtOAc in hexane to afford the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.2-6.8 (m, 8H), 3.84 (m, 1H), 3.16 (m, 1H), 2.80-2.68 (m, 2H), 2.24 (s, 3H), 1.45 (s, 9H), 0.86 (d, 3H). LC-MS: *m/e* 396 (M + Na)<sup>+</sup> (4.4 min).

Step B      *N*-[3-(4-Chlorophenyl)-2-(3-methylphenyl)-1-methylpropyl]amine  
20                    hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 2, Step I. LC-MS: *m/e* 274 (M + H)<sup>+</sup> (2.5 min).

#### REFERENCE EXAMPLE 51




N-[3-(4-Chlorophenyl)-2-(3-trifluoromethylphenyl)-1-methylpropyl]amine  
hydrochloride

(Diastereomer  $\alpha$ )

- 5                    The title compound was prepared following the procedure described in Reference Example 5 substituting fluorophenylacetic acid with 3-trifluoromethylphenylacetic acid at Step A. LC-MS:  $m/e$  328 ( $M + H$ )<sup>+</sup> (2.6 min).

REFERENCE EXAMPLE 52



10

N-[3-(5-Chloro-2-pyridyl)-2(S)-phenyl-1(S)-methylpropyl]amine hydrochloride  
 (Diastereomer  $\alpha$ )

Step A            5-Chloro-2-methylpyridine

- A mixture of 2,5-dichloropyridine (15 g, 0.10 mol), tetramethyltin (15 mL, 0.11 mol),  
 15            and dichlorobis(triphenylphosphine)palladium (2.0 g, 2.8 mmol) in 200 mL  
 anhydrous DMF was heated at 110°C under nitrogen for 72 h. The reaction mixture  
 was cooled to room temperature, and was poured into a saturated solution of  
 potassium fluoride (200 mL). The resulting mixture was partitioned between water  
 (500 mL) and ether (500 mL). The organic layer was separated and the aqueous layer  
 20            was extracted with ether (200 mL). The combined extracts were dried over anhydrous

MgSO<sub>4</sub>, filtered and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 2 to 10% ether in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.41 (d, 1H), 7.75 (dd, 1H), 7.30 (d, 1H), 2.53 (s, 3H).

5    Step B        4-(5-Chloro-2-pyridyl)-3(S)-phenyl-2(R)-butanol.

To a solution of 5-chloro-2-methylpyridine (Step A, 1.1 g, 8.7 mmol) in 15 mL anhydrous ether was added phenyl lithium (1.8 M in cyclohexane/ether, 7.2 mL, 13 mmol) at 0°C, and the reaction was stirred at room temperature for 30 min. The resulting mixture was cooled back to 0°C, and was added (1*R*,2*R*)-1-phenylpropylene oxide (2.3 g, 17 mmol), and the reaction was allowed to warm to room temperature overnight. The reaction mixture was partitioned between EtOAc (100 mL) and water (100 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 100 mL). The combined organic extracts were dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 10 to 40% EtOAc in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.28 (d, 1H), 7.59 (dd, 1H), 7.25-7.12 (m, 5H), 7.05 (d, 1H), 4.03 (m, 1H), 3.29 (dd, 1H), 3.19 (dd, 1H), 3.12 (m, 1H), 1.12 (d, 3H).

20    Step C        2(S)-Azido-4-(5-chloro-2-pyridyl)-3(S)-phenylbutane

To a mixture of 4-(5-chloro-2-pyridyl)-3-phenyl-2-butanol (Step B, 0.24 g, 0.92 mmol), triphenylphosphine (1.5 g, 1.4 mmol) and diphenylphosphoryl azide (0.30 mL, 1.4 mmol) in 5 mL anhydrous THF was added diethylazodicarboxylate (0.24 mL, 1.4 mmol). After stirring at room temperature overnight, the resulting mixture was concentrated with silica gel (10 g) and the residue was loaded onto a silica gel column. Elution with 5 to 15% EtOAc in hexane afforded the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.35 (d, 1H), 7.52 (dd, 1H), 7.25-7.05 (m, 5H), 6.95 (d, 1H), 3.81 (m, 1H), 3.48 (m, 1H), 3.15-3.05 (m, 2H), 1.14 (d, 3H).


25    Step D        N-[3-(5-Chloro-2-pyridyl)-2(S)-phenyl-1(S)-methylpropyl]amine, hydrochloride

30    The product of Step C (0.20 g, 0.70 mmol) was converted to the title compound following the procedure described in Reference Example 2, Steps H-I, except hydrogen chloride in dioxane (4 M) was used in place of hydrogen chloride in EtOAc. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.75 (d, 1H), 8.19 (dd, 1H), 7.55 (d, 1H), 7.4-7.2



(m, 5H), 3.78 (m, 1H), 3.62 (dd, 1H), 3.48 (m, 1H), 3.43 (dd, 1H), 1.22 (d, 3H). LC-MS:  $m/e$  261 ( $M + H$ )<sup>+</sup> (2.2 min).

### REFERENCE EXAMPLE 53



5

N-[2-(3-Bromophenyl)-3-(5-chloro-2-pyridyl)-1-methylpropyl]amine hydrochloride (Diastereomer α)

Step A 3-Bromophenylacetone

To a solution of N-methoxy-N-methylacetamide (10 g, 100 mmol) in 100 mL anhydrous ether at 0°C was added 3-bromobenzylmagnesium bromide (0.25 M in ether, 200 mL, 50 mmol). The reaction was allowed to warm to room temperature overnight and was quenched by the addition of saturated ammonium chloride (100 mL). The organic layer was separated and the aqueous layer was extracted with hexane (100 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated to dryness to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.45-7.40 (m, 2H), 7.26 (t, 1H), 7.19 (d, 1H), 2.20 (s, 3H).

15

Step B 3-(3-Bromophenyl)-4-(5-chloro-2-pyridyl)-2-butanone

A suspension of 5-chloro-2-methylpyridine (Reference Example 52, Step A, 6.4 g, 50 mmol) and N-bromosuccinimide (12.5 g, 70 mmol) in 100 mL carbon tetrachloride was heated to gentle reflux (bath temperature 90°C), and 2,2'-azobisisobutyronitrile (0.74 g) was added in several portions over 30 min. After stirring at this temperature for 5 h, the reaction mixture was concentrated. The resulting slurry was diluted with EtOAc (100 mL) and was washed with water (100 mL), saturated aqueous sodium bicarbonate/saturated aqueous sodium thiosulfate, and brine. The organic solution was dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 2 to 15% ether in CH<sub>2</sub>Cl<sub>2</sub>/hexane (1:1) to afford 2-bromomethyl-5-chloropyridine (6.0 g, 60%), which was used immediately

25

for the ensuing reaction. Thus, to a vigorously stirred solution of 2-bromomethyl-5-chloropyridine (6.0 g, 29 mmol) and 3-bromophenyl acetone (Step A, 6.0 g, 28 mmol) and tetrabutylammonium iodide (20 mg) in 30 mL CH<sub>2</sub>Cl<sub>2</sub> at -78°C was added cesium hydroxide monohydrate (10 g, 60 mmol), and the reaction was allowed to slowly warm to room temperature overnight. The reaction mixture was partitioned between EtOAc (100 mL) and water (100 mL). The organic layer was separated and the aqueous layer extracted with EtOAc (2 x 100 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography on silica gel eluted with 5 to 40% EtOAc in hexane to afford the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.44 (d, 1H), 7.66 (dd, 1H), 7.46-7.41 (m, 2H), 7.24 (t, 1H), 7.22 (d, 1H), 7.15 (d, 1H), 4.42 (dd, 1H), 3.54 (dd, 1H), 3.07 (dd, 1H), 2.12 (s, 3H). LC-MS: m/e 338 (M + H)<sup>+</sup> (3.0 min).


Step C      3-(3-Bromophenyl)-4-(5-chloro-2-pyridyl)-2-butanol

To a solution of 3-(3-bromophenyl)-4-(5-chloro-2-pyridyl)-2-butanone (Step B, 6.7 g, 20 mmol) in 50 mL anhydrous THF at -78°C was added lithium tri(sec-butyl)borohydride (1.0 M in THF, 30 mL, 30 mmol), and the reaction was allowed to warm to room temperature overnight. The reaction was cooled to 0°C, and was carefully added 2 M hydrochloric acid (50 mL), and the resulting mixture was partitioned between hexane (200 mL) and water (200 mL). The aqueous layer was separated and the organic layer extracted with 2 M hydrochloric acid (2 x 100 mL). The combined aqueous extracts were neutralized with 5 N aqueous sodium hydroxide (pH > 12), and was extracted with EtOAc (2x200 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to afford the title compound.

Step D      N-[2-(3-Bromophenyl)-3-(5-chloro-2-pyridyl)-1-methylpropyl]amine, hydrochloride

The product of Step C (5.9 g, 17 mmol) was converted to the title compound following the procedure described in Reference Example 52, Steps C-D. LC-MS: m/e 338 (M + H)<sup>+</sup> (2.3 min).

REFERENCE EXAMPLE 54




N-[3-(5-Chloro-2-pyridyl)-2-(3-chlorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described in Reference

- 5 Example 42 substituting 2-(*N-tert*-butoxycarbonyl)amino-3-bromophenyl-4-(4-chlorophenyl)butane with 2-(*N-tert*-butoxycarbonyl)amino-3-bromophenyl-4-(5-chloro-2-pyridyl)butane (intermediate of Reference Example 53; Step D) at Step A. LC-MS:  $m/e$  295 ( $M + H$ )<sup>+</sup> (2.0 min).

10

REFERENCE EXAMPLE 55



N-[2-(5-Bromo-2-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

Step A 5-Bromo-3-pyridylacetone

- 15 A mixture of 3,5-dibromopyridine (50 g, 0.21 mol), isopropenyl acetate (26 mL, 0.23 mmol), tris(dibenzylideneacetone)dipalladium (1.0 g, 1.1 mmol) and 2-(diphenylphosphino)-2'-(*N,N*-dimethylamino)biphenyl (1.6 g, 4.2 mmol) in 400 mL toluene was heated at 100°C under nitrogen for 2 h. The reaction mixture was cooled to room temperature, and was concentrated to about 100 mL. The resulting mixture  
20 was loaded onto a silica gel column, which was eluted with 0 to 60% EtOAc in

hexane to afford the title compound.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  8.54 (br s, 1H), 8.33 (br s, 1H), 7.88 (br s, 1H), 3.90 (s, 2H), 2.25 (s, 3H).

Step B 3-(5-Bromo-3-pyridyl)-4-(4-chlorophenyl)-2-butanol

The title compound was prepared following the procedure described in Reference


- 5 Example 53, Step B-C, substituting 2-bromomethyl-5-chloropyridine with 4-chlorobenzyl chloride and 3-bromophenylacetone with 5-bromo-3-pyridylacetone (Step A).  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  8.43 (d, 1H), 8.24 (d, 1H), 7.98 (dd, 1H), 7.17 (d, 2H), 7.07 (d, 2H), 4.04 (m, 1H), 3.16 (dd, 1H), 3.0-2.9 (m, 2H), 1.04 (d, 3H).

Step C N-[2-(5-Bromo-3-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]amine

10 hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 31, Step B. LC-MS:  $m/e$  339 ( $\text{M} + \text{H}$ ) $^+$  (2.5 min).

#### REFERENCE EXAMPLE 56




15

N-[2-(5-Bromo-3-pyridyl)-3-(4-fluorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 55 substituting 4-chlorobenzyl chloride with 4-fluorobenzyl chloride at Step

- 20 B. LC-MS:  $m/e$  323 ( $\text{M} + \text{H}$ ) $^+$  (2.3 min).

#### REFERENCE EXAMPLE 57



N-[3-(4-Chlorophenyl)-2-(5-cyano-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

Step A      5-Cyano-3-pyridylacetone


- 5 The title compound was prepared following the procedure described for Reference Example 55 substituting 3,5-dibromopyridine with 5-bromonicotinonitrile (5-bromo-3-cyanopyridine) at Step A.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  8.89 (d, 1H), 8.60 (d, 1H), 8.02 (t, 1H), 3.98 (s, 2H), 2.24 (s, 3H).

Step B      N-[3-(4-Chlorophenyl)-2-(5-cyano-2-pyridyl)-1-methylpropyl]amine  
10 hydrochloride (Diastereomer  $\alpha/\beta$  5:1)

The title compound was prepared following the procedure described for Reference Example 4 substituting 3-pyridylacetone with 5-cyano-3-pyridylacetone (Step A).  
LC-MS:  $m/e$  286 ( $M + H$ ) $^+$  (1.9 min).

15

REFERENCE EXAMPLE 58




N-[2-(5-Cyano-3-pyridyl)-3-(4-fluorophenyl)-1-methylpropyl]amine hydrochloride

(Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 66 substituting 4-chlorobenzyl chloride with 4-fluorobenzyl chloride at Step B. LC-MS:  $m/e$  270 ( $M + H$ )<sup>+</sup> (2.2 min).

5

REFERENCE EXAMPLE 59




N-[2-(5-Cyano-3-pyridyl)-3-(3,4-difluorophenyl)-1-methylpropyl]amine

10 hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 58 substituting 4-fluorobenzyl chloride with 3,4-difluorobenzyl chloride at Step B. LC-MS:  $m/e$  288 ( $M + H$ )<sup>+</sup> (2.3 min).

15


REFERENCE EXAMPLE 60



N-[3-(3-Chlorophenyl)-2-(5-cyano-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 58 substituting 4-fluorobenzyl chloride with 3-chlorobenzyl chloride at Step B. LC-MS: m/e 286 (M + H)<sup>+</sup> (2.4 min).

REFERENCE EXAMPLE 61



N-[3-(4-Chlorophenyl)-2-(5-chloro-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )


Step A 5-Chloro-3-pyridylacetone

The title compound was prepared following the procedure described for Reference Example 50 substituting 3,5-dibromopyridine with 3,5-dichloropyridine and 2-(diphenylphosphino)-2'-(N,N-dimethylamino)biphenyl with 2-(di-t-butylphosphino)biphenyl at Step A. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  8.42 (d, 1H), 8.27 (d, 1H), 7.73 (dd, 1H), 3.90 (s, 2H), 2.25 (s, 3H).

Step B N-[3-(4-Chlorophenyl)-2-(5-chloro-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 55, Step B-C substituting 5-bromo-3-pyridylacetone with 5-chloro-3-pyridylacetone at Step B. LC-MS: m/e 295 (M + H)<sup>+</sup> (1.9 min).


REFERENCE EXAMPLE 62



N-[2-(5-Chloro-3-pyridyl)-3-(4-fluorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

- 5 The title compound was prepared following the procedure described for Reference Example 70 substituting 4-chlorobenzyl chloride with 4-fluorobenzyl chloride at Step B. LC-MS:  $m/e$  279 ( $M + H$ )<sup>+</sup> (2.3 min).

#### REFERENCE EXAMPLE 63




- 10 2-Amino-3-(5-chloro-3-pyridyl)-5-methylhexane, Hydrochloride Salt (Diastereomer  $\alpha/\beta$  6:1)

The title compound was prepared following the procedure described for Reference Example 61 substituting 4-chlorobenzyl chloride with 1-iodo-2-methylpropane at Step B. LC-MS:  $m/e$  227 ( $M + H$ )<sup>+</sup> (2.2 min).

15

#### REFERENCE EXAMPLE 64






N-[2-(5-Chloro-3-pyridyl)-3-cyclobutyl-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha/\beta$  6:1)

- The title compound was prepared following the procedure described  
5 for Reference Example 61 substituting 4-chlorobenzyl chloride with  
(bromomethyl)cyclobutane at Step B. LC-MS:  $m/e$  239 ( $M + H$ )<sup>+</sup> (2.3 min).

#### REFERENCE EXAMPLE 65



- 10 N-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )


Step A 3-Cyanophenylacetone

- The title compound was prepared following the procedure described for Reference  
Example 55 substituting 3,5-dibromopyridine with 3-bromobenzonitrile and 2-  
15 (diphenylphosphino)-2'-(N,N-dimethylamino)biphenyl with 2-  
(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl at Step A. <sup>1</sup>H NMR (500  
MHz, CD<sub>3</sub>OD):  $\delta$  7.6 (m, 1H), 7.56 (br s, 1H), 7.50-7.48 (m, 2H), 3.88 (s, 2H), 2.21  
(s, 3H).

- Step B N-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]amine  
20 hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 55 substituting 5-bromo-3-pyridylacetone with 3-canophenylacetone at Step B. LC-MS:  $m/e$  285 ( $M + H$ )<sup>+</sup> (2.2 min).

5

REFERENCE EXAMPLE 66

N-[3-(4-Chlorophenyl)-2-(5-fluoro-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

10 Step A 5-fluoro-3-pyridylacetone


The title compound was prepared following the procedure described for Reference Example 55 substituting 3,5-dibromopyridine with 3-fluoro-5-trifluoromethanesulfonyloxy pyridine (prepared from 3-fluoro-5-hydroxypyridine and triflic anhydride) and 2-(diphenylphosphino)-2'-(N,N-dimethylamino)biphenyl with 2-(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl at Step A. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  8.34 (d, 1H), 8.22 (br s, 1H), 7.50 (ddd, 1H), 3.93 (s, 2H), 2.25 (s, 3H).

15

Step B N-[3-(4-Chlorophenyl)-2-(5-chloro-3-pyridyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

20 The title compound was prepared following the procedure described for Reference Example 55, Step B-C substituting 5-bromo-3-pyridylacetone with 5-fluoro-3-pyridylacetone at Step B. LC-MS:  $m/e$  279 ( $M + H$ )<sup>+</sup> (2.4 min).

REFERENCE EXAMPLE 67




N-[3-(4-Chlorophenyl)-2-(5-methyl-3-pyridyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

- The title compound was prepared following the procedure described for Reference Example 50 substituting 2-(*N-tert*-butoxycarbonyl)amino-3-(3-bromophenyl)-4-(4-chlorophenyl)butane with 2-(*N-tert*-butoxycarbonyl)amino-3-(5-bromo-3-pyridyl)-4-(4-chlorophenyl)butane (intermediate of Reference Example 64, Step B) at Step A. LC-MS:  $m/e$  275 ( $M + H$ )<sup>+</sup> (1.3 min).

10

REFERENCE EXAMPLE 68



N-[2-(3-Bromo-5-fluorophenyl)-3-(4-Chlorophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

- 15 Step A 3-Bromo-5-fluorophenylacetone


The title compound was prepared following the procedure described for Reference Example 55 substituting 3,5-dibromopyridine with 1,3-dibromo-5-fluorobenzene and 2-(diphenylphosphino)-2'-(*N,N*-dimethylamino)biphenyl with 1,1'-bis(diphenylphosphino)ferrocene at Step A. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.23

- 20 (d, 1H), 7.22 (s, 1H), 6.96 (d, 1H), 3.81 (s, 2H), 2.20 (s, 3H).

Step B *N*-[2-(3-Bromo-5-fluorophenyl)-3-(4-chlorophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 55, Step B substituting 5-bromo-3-pyridylacetone with 3-bromo-5-fluorophenylacetone (Step A). LC-MS:  $m/e$  356 ( $M + H$ )<sup>+</sup> (2.9 min).

#### REFERENCE EXAMPLE 69




10 *N*-[2-(3-Bromo-5-fluorophenyl)-3-(4-fluorophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the procedure described for Reference Example 68 substituting 4-chlorobenzyl chloride with 4-fluorobenzyl chloride at Step B. LC-MS:  $m/e$  340 ( $M + H$ )<sup>+</sup> (2.8 min).

15

#### REFERENCE EXAMPLE 70



20 2-Amino-3-indolin-N-yl-4-(4-chlorophenyl)butane

Step A. Ethyl 3-(4-chlorophenyl)-2-indolin-N-ylpropanoate.

In an oven-dried flask under an atmosphere of nitrogen, 1.1 g LiOH·H<sub>2</sub>O (26.25 mmol) in DMF (20 mL) was added to a stirring suspension of 4 angstrom molecular sieves. After 30 minutes of stirring at room temperature 2.8 mL (25 mmol) indoline was added dropwise. After one hour at room temperature 2.9 mL (26.25 mmol) Ethyl bromoacetate was added dropwise. After 1.5 h the solid material was filtered and the residue was washed with copious amounts of EtOAc. The organics were washed 3 times with water and the organic material was dried over MgSO<sub>4</sub>. The solvents were evaporated under reduced pressure. The crude material was then dissolved in 75 mL anhydrous THF, charged into an oven dried round bottom under an atmosphere of nitrogen, cooled to -78°C, and then treated with 26.25 mL a 1M solution of NaHMDS. The solution was allowed to stir for 30 minutes at -78°C after which the enolate was quenched with 5.4 g (26.25 mmol) of parachlorobenzyl bromide (solution in 25 mL anhydrous THF). The reaction was allowed to warm to room temperature overnight. The next day the reaction was quenched with water. The aqueous layer was extracted with 3 large portions of EtOAc. The combined organics were dried over MgSO<sub>4</sub>. The solvents were removed under reduced pressure and the residue was purified by flash chromatography which yielded the title compound as a yellow oil. LC/MS m/e=331 (M+1). TLC R<sub>f</sub>=0.22 (20:1 hexanes : EtOAc). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 1.11 (t, J=3.55 Hz, 3H), 2.96 (m, 2H), 3.06 (m, 1H), 3.25 (m, 1H), 3.60 (t, 2H), 4.07 (m, 2H), 4.36 (t, J=3.75 Hz, 1H).

Step B. N,O-dimethyl-3-(4-chlorophenyl)-2-indolin-N-ylpropanamide.

In an oven-dried flask under an atmosphere of nitrogen, 11.75 mL 1 M solution of (CH<sub>3</sub>)<sub>2</sub>AlCl in CH<sub>2</sub>Cl<sub>2</sub> was added via addition funnel to a stirring suspension of 1.15 g (11.75 mmol) N,O-dimethylhydroxylamine hydrochloride at 0°C. After warming to room temperature a solution of 970 mg (2.94 mmol) of Ethyl 3-(4-chlorophenyl)-2-indolinypropanoate in 10 mL was added via addition funnel. After stirring at room temperature for 5 h, 35 mL pH=8 phosphate buffer solution was added and the resulting solution was stirred vigorously for 30 minutes. The phases were separated and the aqueous layer was extracted 2 times with chloroform. The combined organics were washed with water and then dried over MgSO<sub>4</sub>. A brown oil was collected. The crude material was carried on to the next step. ). TLC R<sub>f</sub>=0.12 (10:1 hexanes : EtOAc). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 2.83 (m, 1H), 2.97(m, 2H), 3.13 (s, 3H), 3.34 (m, 1H), 3.45 (s, 3H), 3.61 (m, 2H), 4.87 (b, 1H), 6.54

(d, 1H), 6.66 (t, J=7.1 Hz, 1H), 7.07 (t, J=7.1 Hz, 2H), 7.18 (d, J=8.5 Hz, 2H), 7.24 (d, J=8.5 Hz, 2H)

Step C. 4-(4-chlorophenyl)-3-indolin-N-ylbutan-2-one.

In an oven dried flask under an atmosphere of nitrogen, 2.8 mL 1 M solution of  
5 CH<sub>3</sub>MgBr in THF was added dropwise to a stirring solution of N,O-dimethyl-3-(4-chlorophenyl)-2-indolinylpropanamide (965 mg) in 25 mL anhydrous THF. The solution was stirred for 4 h while being allowed to warm to room temperature. Then approximately 20 mL water were added. The solution was extracted three times with 50 mL ether. The combined extracts were dried over MgSO<sub>4</sub>. The solvents were  
10 removed under reduced pressure yielding a brown oil which was carried on to the next step without purification. LC/MS m/e=301 (M+1). TLC R<sub>f</sub>=0.5 (4:1 hexanes:EtOAc). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 2.14 (s, 3H), 2.81 (dd, J=14.6, 6.6 Hz, 1H), 2.97 (t, J=8.5 Hz, 2H), 3.26 (m, 2H), 3.5 (m, 1H), 4.21 (dd, J=6.6, 6.6 Hz), 6.39 (d, J=8 Hz, 1H), 6.66 (dd, J=7, 7 Hz, 1H), 7.07 (m, 2H), 7.13 (d, J=8.5 Hz), 7.22 (d, J=8.3 Hz).

15 Step D. 4-(4-chlorophenyl)-3-indolin-N-ylbutan-2-one methoxime.

A solution of 472 mg (1.573 mmol) of the product of Step C and 263 mg (3.147 mmol) of methoxylamine hydrochloride in anhydrous ethanol was treated with 255 μL (3.147 mmol) of pyridine. The solution was stirred for 2 h at room temperature. Solvent was removed under reduced pressure and the residue was partitioned between  
20 water and ether. The water was extracted with ether again. The extracts were then combined and dried over MgSO<sub>4</sub>, filtered and concentrate to obtain crude material. obtained. Both the E and Z isomers were carried onto the next step. LC/MS m/e=330 (M+1). TLC R<sub>f</sub>=.77 and .65 (4:1 hexanes:EtOAc). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 1.78 (2s, 1H), 2.88 (dd, J=6.2, 13.8 Hz, 1H), 2.95 (m, 2H), 3.30 (m, 2H), 3.45 (m, 1H), 3.75 and 3.89 (2s, 3H), 4.21 (dd, J=6.9, 7.8 Hz, 1H), 6.28 and 6.47 (2d, J=8.1, 1H), 6.61 (m, 1H), 7.02 (m, 2H), 7.22 (m, 4H).


Step E. 2-Amino-3-indolin-N-yl-4(4-chloro)phenylbutane

In an oven-dried flask equipped with a water condenser under an atmosphere of nitrogen, a solution of 301 mg (0.914 mmol) 4-(4-chlorophenyl)-3-indolinylbutan-2-one methoxime in 1.5 mL anhydrous THF was treated with 3.7 mL (3.7 mmol) of 1M  
30 BH<sub>3</sub>·THF at room temperature. The solution was then heated to 75°C for 2 days. The solution was then cooled to 0°C and treated with chips of ice until bubbling subsided. 500 μL of 20% KOH were then added and the solution was heated at 45°C for 2h. The solution was then cooled to room temperature and extracted with ether  
35 3x. The combined extracts were dried over MgSO<sub>4</sub>, filtered, and concentrated to

afford crude amine which was used in the next experiment without further purification. LC/MS  $m/e=302$  ( $M+1$ ).  $^1H$  NMR (500 MHz,  $CDCl_3$ ):  $\delta$  1.13, 1.14 (2d,  $J=6.5$  Hz, 1H), 1.55-1.60 (m, 2H), 2.80-3.10 (m, 4H), 3.30-3.60 (m, 2H), 6.348 and 6.38 (2d,  $J=7.9$  Hz, 1H), 6.50-6.78 (m, 2H), 6.95-7.24 (m, 5H)

5

#### REFERENCE EXAMPLE 71




#### 2-Amino-3-indol-N-yl-4(4-chloro)phenylbutane

10 This compound was prepared in an analogous manner to Reference Example 70 except that during Step A, sodium hydride was used as the base instead of the lithium hydroxide monohydrate/molecular sieves combination. LC/MS: calculated for  $C_{18}H_{19}ClN_2$  299, observed  $m/e$  300 ( $M + H$ )<sup>+</sup> (2.4 min).


15

#### REFERENCE EXAMPLE 72



#### 2-Amino-3-(N-methyl, N-phenyl)amino-4(4-chloro)phenylbutane


20 This compound was prepared in an analogous manner to Reference Example 70. LC/MS: calculated for  $C_{17}H_{21}ClN_2$  289, observed  $m/e$  290 ( $M + H$ )<sup>+</sup> (2.4 min).

REFERENCE EXAMPLE 732-Amino-3-(7-azaindol-N-yl)-4(4-chloro)phenylbutane

5                      This compound was prepared in an analogous manner to Reference Example 70. LC/MS: calculated for  $C_{17}H_{18}ClN_3$  300, observed  $m/e$  301 ( $M + H$ )<sup>+</sup> (2.7 min).

REFERENCE EXAMPLE 74


10

2-Amino-3-(benzisoxazol-3-yl)-4(4-chloro)phenylbutane

15                      This compound was prepared in an analogous manner to Reference Example 70 except starting with ethyl (benzisoxazol-3-yl)acetate. LC/MS: calculated for  $C_{17}H_{17}ClN_2O$  300, observed  $m/e$  301 ( $M + H$ )<sup>+</sup> (2.2 min).

REFERENCE EXAMPLE 75





4-(4-Methylphenyl)-3-phenylbutan-2-amine (mixture of 4 isomers)

Step A 1-Phenylacetone

To a solution of N-methyl-N-methoxyacetamide (9.9 mL, 97 mmol) in ether (300 mL) at 0°C was added benzylmagnesium chloride (97 mL a 1M solution in ether). The cloudy, white reaction mixture was warmed to room temperature for 2 h and then quenched by careful addition of 1N hydrochloric acid (100 mL). The organic phase was separated, washed with brine, dried over MgSO<sub>4</sub> and concentrated. The crude material was purified by column chromatography on silica gel eluting from 0-10% EtOAc/hexane to give the title compound. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.36 (t, J = 7.1 Hz, 2H), 7.30 (t, J = 7.3 Hz, 1H), 7.24 (d, J = 7.3 Hz, 2H), 3.72 (s, 2H), 2.18 (s, 3H). LC-MS: m/e 135 (M + H)<sup>+</sup> (1.95 min).

Step B 4-(4-Methylphenyl)-3-phenylbutan-2-one

1-Phenylacetone (200 mg, 1.49 mmol) was mixed with powdered potassium hydroxide (167 mg, 2.98 mmol) and tetra-n-butylammonium bromide (1 mol %, 5 mg) in a flask without solvent. This mixture was stirred at room temperature for 90 min. before the addition of 1-(chloromethyl)-4-methylbenzene (198 µL, 1.49 mmol). The reaction mixture was then stirred overnight before diluting with water and CH<sub>2</sub>Cl<sub>2</sub>. The aqueous layer was separated and neutralized to pH 7 with 2N hydrochloric acid and extracted again into CH<sub>2</sub>Cl<sub>2</sub>. The combined organic washes were dried with MgSO<sub>4</sub> and concentrated. The crude material was purified by column chromatography on silica gel eluting from 0-10% EtOAc/hexane to give the title compound. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.35 (t, J = 7.0 Hz, 2H), 7.29 (t, J = 7.4 Hz, 1H), 7.23 (d, J = 7.1 Hz, 2H), 7.05 (d, 7.8 Hz, 2H), 6.98 (d, J = 7.8 Hz, 2H), 3.94 (t, J = 7.3 Hz, 1H), 3.43 (dd, J = 13.9, 7.5 Hz, 1H), 2.91 (dd, J = 14, 7.1 Hz, 1H), 2.32 (s, 3H), 2.08 (s, 3H). LC-MS: m/e 239 (M + H)<sup>+</sup> (3.61 min).

Step C 4-(4-Methylphenyl)-3-phenylbutan-2-amine

To a solution of the 4-(4-methylphenyl)-3-phenylbutan-2-one (308 mg, 1.29 mmol) in 7M ammonia in MeOH (5 mL) and acetic acid (3 mL) was added sodium

- cyanoborohydride (130 mg, 2.06 mmol) and the reaction stirred at room temperature overnight. The reaction was quenched by pouring into 2M sodium carbonate solution and extracted into EtOAc. The aqueous layer was salted and re-extracted. The combined organic extracts were dried over MgSO<sub>4</sub> and concentrated to give the title compound as a mixture of 4 isomers which was used without further purification. LC-MS: m/e 240 (M + H)<sup>+</sup> (2.22 min).

REFERENCE EXAMPLE 76




- 10 4-(4-Methoxyphenyl)-3-phenylbutan-2-amine

Prepared using the procedures described in Example 75, Steps A through C using 1-(chloromethyl)-4-methoxybenzene as the alkylating agent in Step B. LC-MS: m/e 256 (M + H)<sup>+</sup> (1.90 and 2.03 min).

15


REFERENCE EXAMPLE 77



3-[2-Amino-1-(4-fluorobenzyl)propyl]benzonitrile

- 20 Prepared using the procedures described in Example 65 using 3-(2-oxopropyl)benzonitrile and 1-(chloromethyl)-4-fluorobenzene as the reactants in Step B. LC-MS: m/e 269 (M + H)<sup>+</sup> (2.87 min).

REFERENCE EXAMPLE 78




N-[2-Phenyl-3-(4-fluorophenyl)-1-methylpropyl]amine hydrochloride  
(Diastereomer  $\alpha$ )

- 5 The title compound was obtained by the method described in Reference Example 38, substituting 4-fluorobenzyl bromide for isobutyl iodide. LC-MS,  $R_t = 2.2$  min,  $m/e = 244$ .

REFERENCE EXAMPLE 79

10



2-(2,3-Dihydro-1H-indol-1-yl)-1,4-dimethylpentylamine

Step A Ethyl (2-(2,3-dihydro-1H-indol-1-yl)-4-methylpentanoate

- A solution of 0.53 g (3.3 mmol) of ethyl (S)-2-hydroxyisocaproate in 8 mL dry  
15  $\text{CH}_2\text{Cl}_2$  was cooled in a  $-78^\circ\text{C}$  bath and 0.73 mL (4.34 mmol) of triflic anhydride  
and 0.6 mL (5.36 mmol) of 2,6 lutidine were added. After 15 min 2 mL (11.5 mmol)  
of diisopropylethylamine was added and stirred for 10 min. To this solution 0.36 mL  
(3.21 mmol) of 2,3-dihydroindoline was added and stirred overnight as it slowly  
warmed to room temperature. The reaction was quenched with saturated  $\text{NaHCO}_3$   
20 solution and extracted with ether. The combined organic layer was washed with  
water, brine, dried and concentrated. The residue was purified on a flash column  
using a gradient of 5-10% EtOAc/hexane to isolate the title compound.  $^1\text{H}$  NMR:  
(500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.99 (d, 3H), 1.03 (d, 3H), 1.22 (t, 3H), 1.81 (m, 3H), 3.04 (m,  
2H), 3.57 (m, 1H), 3.66 (m, 1H), 4.14 (q, 2H), 4.24 (t, 1H), 6.4-7.1 (m, 4H).

- 25 Step B 3-(2,3-Dihydro-1H-indol-1-yl)-5-methylhexan-2-one

To a solution of 0.54 g (2.07 mmol) of ethyl (2-(2,3-dihydro-1H-indol-1-yl)-4-methylpentanoate in 10 mL CH<sub>2</sub>Cl<sub>2</sub>, 1.98 g (10 mmol) of N,O-dimethylhydroxylamine hydrochloride and 1.4 mL triethylamine were added. The mixture was cooled in an ice bath and 10 mL (10 mmol) 1 M diethylaluminum chloride in toluene was added. The reaction was stirred overnight as it warmed to room temperature then carefully quenched by pouring into 1.2 N HCl. The solution was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with brine, dried and concentrated leaving amide which was used without purification. This amide was dissolved in 5 mL THF and 2.5 mL (3.5 mmol) of 1.4 M methylmagnesium bromide was added. After 1 h, the solution was quenched with 1.2 N HCl and extracted with EtOAc. The EtOAc layer was washed with brine, dried and concentrated. The residue was chromatographed using a gradient of 5-10% EtOAc-hexane to isolate the title compound. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 0.96 (d, 3H), 0.99 (d, 3H), 1.7 (m, 3H), 2.17 (s, 3H), 3.06 (m, 2H), 3.04 (q, 1H), 3.52 (m, 1H), 4.11 (m, 1H) 6.4-7.1 (m, 4H).


Step C      2-(2,3-Dihydro-1-H-indol-1-yl)-1,4-dimethylpentylamine

To a solution of 0.185 g (0.8 mmol) of 3-(2,3-dihydro-1H-indol-1-yl)-5-methylhexan-2-one in 2 mL ethanol, 0.135 g O-methylhydroxylamine hydrochloride and 0.13 mL (1.6 mmol) of pyridine were added. After stirring for 2 h, the solution was concentrated and the residue was partitioned between water and EtOAc. The organic layer was washed with brine, dried and concentrated to give 0.2 g O-methyloxime as a mixture of isomers. This mixture was dissolved in 2 mL THF and 1.5 mL 1 M BH<sub>3</sub> in THF was added. After gas evolution ceased, the reaction was heated in a 50 °C bath. After 2 h another 1.5 mL 1 M BH<sub>3</sub> in THF was added and heating was continued overnight. The reaction mixture was cooled and quenched with MeOH and concentrated. The residue was dissolved in 6 mL CH<sub>2</sub>Cl<sub>2</sub> and 2 mL 1 N NaOH was added. After stirring for 15 min the layers were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was washed with water, brine dried and concentrated to isolate title compound as a mixture of diastereomers which was used without purification. LC-MS, R<sub>t</sub> = 2.24 min, m/e = 233.

The following amines were synthesized by the method of Reference

Example 70.

REFERENCE EXAMPLE 80




3-Cyclobutyl-2-(3,4-dihydroquinoline-1(2H)-yl)-1-methylpropylamine

LC-MS,  $R_t = 2.8$  min,  $m/e = 259$ .

5

REFERENCE EXAMPLE 81




2-(3,4-Dihydroquinoline-1(2H)-yl)-1,4-dimethylpentylamine

LC-MS,  $R_t = 2.74$  min,  $m/e = 248$ .

10

REFERENCE EXAMPLE 82



2-(1H-1,2,3-Benzotriazol-1-yl)-3-(4-chlorophenyl)-1-methylpropylamine

15 Step A 2-(1H-1,2,3-Benzotriazol-1-yl)-N-methoxy-N-methylacetamide

A mixture of 1.77 g (10 mmol) of 2-(1H-1,2,3-benzotriazol-1-yl)acetic acid, 1.07 g (11 mmoles) of N,O-dimethylhydroxylamine hydrochloride, 5.8 g (11 mmol) of PyBOP, and 3.4 mL (24.2 mmol) of diisopropylethylamine in 50 mL  $\text{CH}_2\text{Cl}_2$  was

stirred overnight at RT. This mixture was partitioned between EtOAc and water. The organic layer was washed with brine and dried over anhydrous MgSO<sub>4</sub>. Solvent removal afforded a crude product which was purified on silica gel using 60% EtOAc in hexane as solvent to give 2.01 g the desired amide as a solid. <sup>1</sup>H NMR: (CDCl<sub>3</sub>):

5    δ 3.26 (s, 3H), 3.84 (s, 3H), 5.63 (s, 2H), 7.35-8.2 (m, 4H).

Step B        2-(1H-1,2,3-Benzotriazol-1-yl)-3-(4-chlorophenyl)-N-methoxy-N-methyl-propanamide.

To a solution of 2.0 g (9 mmol) of 2-(1H-1,2,3-benzotriazol-1-yl)-N-methoxy-N-methylacetamide in 15 mL anhydrous THF at -78 °C, 10 mL (10 mmol) of 1M  
10    lithium bis(trimethylsilyl)amide was added dropwise. After stirring for 25 min, a solution of 2.06 g (10 mmol) of 4-chlorobenzyl bromide in 2 mL anhydrous THF was added. The resulting reaction mixture was allowed to warm to RT and stirred for 6 h. This reaction was quenched, diluted with 75 mL EtOAc and washed 3 times with 10 mL each of brine. After drying the organic phase solvent removal afforded a crude  
15    product which was purified on silica gel using 40% EtOAc in hexane as solvent to afford the desired product as a solid. <sup>1</sup>H NMR: (CDCl<sub>3</sub>): δ 3.2 (s, 3H), 3.34 (s, 3H), 3.52 (m, 1H), 3.7 (m, 1H), 6.32 (t, 1H), 6.9-8.2 (m, 8H).

Step C        2-(1H-1,2,3-Benzotriazol-1-yl)-3-(4-chlorophenyl)-butan-2-one.


To a solution of 1.73 g (5 mmol) of 2-(1H-1,2,3-benzotriazol-1-yl)-3-(4-  
20    chlorophenyl)-N-methoxy-N-methyl-propanamide in 10 mL anhydrous THF at 0 °C, 4 mL (10 mmol) of 2.5M methyl magnesium bromide in ether was added. The reaction mixture was stirred for 4 h as it warmed to RT. The reaction was quenched by adding 10 mL 1N HCl and the resulting mixture was partitioned between EtOAc and water. The organic phase was washed with brine and dried over anhydrous  
25    MgSO<sub>4</sub>. Solvent removal gave a crude ketone, which was purified on silica gel using 40% EtOAc in hexane to provide the desired ketone.

Step D        2-(1H-1,2,3-Benzotriazol-1-yl)-3-(4-chlorophenyl)-1-methyl propylamine

To a solution of 1.18 g (4 mmol) of 2-(1H-1,2,3-benzotriazol-1-yl)-3-(4-  
30    chlorophenyl)-butan-2-one in 8.5 mL (60 mmol) of 7N ammonia in MeOH at 0 °C, 4 mL (964 mmol) of glacial acetic acid was added followed by 410 mg (6.5 mmol) of sodium cyanoborohydride. The reaction mixture was allowed to warm to RT and stirred overnight. The reaction was partitioned between EtOAc and saturated NaHCO<sub>3</sub> solution. The organic phase was dried over anhydrous MgSO<sub>4</sub>. The  
35    solvent was removed in vacuo and the residue was purified on silica gel using a

mixture of 5% 2N methanolic ammonia solution and 95% CH<sub>2</sub>Cl<sub>2</sub> to give the desired amine as a mixture of diastereomers. LC-MS, R<sub>t</sub> = 2.0 min, m/e = 301.


### REFERENCE EXAMPLE 83



#### 3-(4-Chlorophenyl)-2-(thiophene-3-yl)-1-methylpropylamine

The title amine was prepared by the method described in Reference Example 82, substituting thiophene-3-acetic acid for 2-(1H-1,2,3-benzotriazol-1-yl)acetic acid in Step A. LC-MS, R<sub>t</sub> = 2.19 min, m/e = 266.

### REFERENCE EXAMPLE 84



#### 3-(4-Chlorophenyl)-2-(thiophene-2-yl)-1-methylpropylamine


Step A 3-(4-Chlorophenyl)-2-(thiophen-2-yl)-butan-2-one.

The title compound was obtained from 2-thiopheneacetic acid according to the procedure described in Reference Example 38, Steps A-C.

Step B 3-(4-Chlorophenyl)-2-(thiophene-2-yl)-1-methylpropylamine

This amine was synthesized by the method of Reference Example 82, Step D. LC-MS, R<sub>t</sub> = 2.18 min, m/e = 266.

### REFERENCE EXAMPLE 85




3-(4-Chlorophenyl)-1-methyl-2-(1-methyl-1H-indol-3-yl)propylamine

The title compound was prepared according to the method described in Reference Example 84. LC-MS:  $R_t = 2.5$  min,  $m/e = 313$ .

5

REFERENCE EXAMPLE 86



3-(4-Chlorophenyl)-1-methyl-2-(1H-indazol-1-yl)propylamine

Step A 3-(4-Chlorophenyl)-2-(1H-indazol-1-yl)-butan-2-one.


10 The title compound was obtained from indazol-1-yl-acetic acid by following the procedure of Reference Example 2, Steps A-D.

Step B 3-(4-Chlorophenyl)-1-methyl-2-(1H-indazol-1-yl)propylamine.

The title amine was prepared according to the procedure of Reference Example 88, Step C. LC-MS:  $R_t = 2.24$  min,  $m/e = 300$ .

15

REFERENCE EXAMPLE 87





3-(4-Chlorophenyl)- 1-methyl-2-(1-methyl-1H-indol-4-yl)propylamineStep A 4-Chloro-1-methylindole

In a 100 mL flask, 0.3 g (7.5 mmol) sodium hydride was washed twice with dry hexane. The solid was suspended in 15 mL dry THF and 1g (6.6 mmol) 4-chloroindole was drop wise added. After 15 min, 0.5 mL (7.9 mmol) methyl iodide was added and the solution was stirred overnight. The reaction was quenched with 1.2 N HCl and partitioned between ether and water. The organic layer was washed with brine, dried and concentrated keeping the bath temperature below 30 °C. The residue was purified on a flash column using a gradient of 5-10% EtOAc/hexane to isolate the desired product. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 3.84 (s, 3H), 6.63 (d, 1H), 7-7.3 (m, 4H).

Step B 1-(1-Methyl-1H-indol-4-yl)acetone

To a solution of 0.852 g (5.14 mmol) of 4-chloro-1-methylindole in 15 mL dry toluene, 0.85 mL (7.73 mmol) isopropenyl acetate and 2.3 mL (8 mmol) tributyltin methoxide were added. The solution was heated to 100 °C. After 15 min, 0.24 g (0.61 mmol) 2-dicyclohexylphosphino-2'-(N,N-dimethylamino) biphenyl and 0.14 g (0.153 mmol) tris (dibenzylideneacetone)dipalladium were added and heating was continued. After 2 h the solution was cooled, filtered through a pad of CELITE diatomaceous earth and the filtrate was concentrated to ca. 5 mL. This solution was purified on a silica column using a gradient of 5-20% EtOAc/hexane to obtain the title compound. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 2.14 (s, 3H), 3.84 (s, 3H), 3.97 (s, 2H), 6.51 (d, 1H), 7-7.3 (m, 4H).


Step C 4-(4-Chlorophenyl)-3-(1-methyl-1H-indol-4-yl)-butan-2-one

To a suspension of 135 mg (3.38 mmol) of sodium hydride in 8 mL dry THF, a solution of 605 mg (3.23 mmol) 1-(1-methyl-1H-indol-4-yl)acetone in 2 mL THF was added. The mixture was stirred for 45 min during which time the sodium hydride dissolved and a yellow orange solution resulted. The reaction was cooled in ice bath and 660 mg (3.24 mmol) 4-chlorobenzyl bromide in 1 mL THF was added. The cold bath was removed and the solution was stirred for 1.5 h. The reaction was quenched with 1.2 N HCl and extracted with EtOAc. The organic layer was washed with brine, dried and concentrated. The residue was chromatographed using a gradient of 10-20% EtOAc/hexane to isolate the desired product. <sup>1</sup>H NMR: (500 MHz, CDCl<sub>3</sub>): δ 2.03 (s, 3H), 3.07 (m, 1H), 3.58 (m, 1H), 3.84 (s, 3H), 4.23 (t, 1H), 6.52 (d, 1H), 6.9-7.3 (m, 8H).

Step D 3-(4-Chlorophenyl)- 1-methyl-2-(1-methyl-1H-indol-4-yl)propylamine

The title compound was prepared from 4-(4-chlorophenyl)-3-(1-methyl-1H-indol-4-yl)-butan-2-one by following the procedure of Reference Example 79, Step C. LC-MS,  $R_t = 2.4$  min,  $m/e = 313$ .

5

REFERENCE EXAMPLE 883-(4-Chlorophenyl)-1-methyl-2-(pyridazin-3-yl)propylamine

Step A 4-(4-Chlorophenyl)-3-(pyridazin-3-yl)-butan-2-one

10 This compound was synthesized from 3-iodopyridazine by the procedure of Reference Example 87, Steps B-C.


Step B N-2,4-Dimethoxybenzyl-N(3-(4-chlorophenyl)-1-methyl-2-(pyridazin-3-yl)propyl)amine

15 A solution of 300 mg (1.15 mmol) 4-(4-chlorophenyl)-3-(pyridazin-3-yl)-butan-2-one in 4 mL dichloroethane was treated with 234 mg (1.15 mmol) 2,4-dimethoxybenzyl amine hydrochloride, 0.16 mL (1.15 mmol) triethylamine and 488 mg (2.3 mmol) sodium triacetoxyborohydride. After stirring the reaction overnight, it was partitioned between water and  $\text{CH}_2\text{Cl}_2$ . The organic layer was washed with brine, dried and concentrated and the residue was purified on a flash column using 3% MeOH-  
20  $\text{CH}_2\text{Cl}_2$  to isolate the desired amine.

Step C 3-(4-Chlorophenyl)-1-methyl-2-(pyridazin-3-yl)propylamine

25 A solution of 300 mg N-2,4-dimethoxybenzyl-N(3-(4-chlorophenyl)-1-methyl-2-(pyridazin-3-yl)propyl)amine in 5 mL trifluoroacetic acid was heated in a 70 °C bath over night followed by 6 h in a 100 °C bath. The reaction was cooled, concentrated and the residue was diluted with EtOAc. This solution was quenched (to pH 10) with 1N NaOH and the layers were separated. The organic layer was washed with brine, dried and concentrated. The residue was purified on a prepTLC using 10% MeOH/ $\text{CH}_2\text{Cl}_2$  with 1%  $\text{NH}_4\text{OH}$  to isolate the title compound (mixture of diastereomers), starting material was also recovered. LC-MS,  $R_t = 1.63$  min,  $m/e = 262$ .


30

REFERENCE EXAMPLE 893-(4-Chlorophenyl)-1-methyl-2-(pyrimidin-5-yl)propylamineStep A 4-(4-Chlorophenyl)-3-(pyrimidin-5-yl)-butan-2-one

- 5 The title compound was obtained from 5-bromopyrimidine following the method of Reference Example 87, Steps B-C except that 2-(di-*t*-butylphosphino)biphenyl was used in place of dicyclohexylphosphino-2'-(*N,N*-dimethylamino)biphenyl in Step B.

Step B 3-(4-Chlorophenyl)-1-methyl-2-(pyrimidin-5-yl)propylamine

- The title compound was prepared by the procedure described in Reference Example 10 30, Steps E-I. LC-MS, *R*<sub>t</sub> = 1.57 min, *m/e* = 262.

REFERENCE EXAMPLE 902-(3-Cyanophenyl)-3-cyclobutyl-1-methylpropylamine

- 15 Step A 1-(3-Cyanophenyl)acetone

The title compound was prepared from 3-bromobenzonitrile and isopropenyl acetate by the procedure of Reference Example 88, Step B.

Step B 3-(3-Cyanophenyl)-4-cyclobutyl-butan-2-one

- To a solution of 1.45 g (9.07 mmol) of 1-(3-cyanophenyl)acetone in 18 mL  
20 acetonitrile, 1.1 mL (9.5 mmol) cyclobutyl bromide and 5.91 g (18.1 mmol) cesium carbonate were added. After heating the solution in a 60 °C bath overnight, it was cooled and filtered. The filtrate was partitioned between water and EtOAc and the aqueous layer was extracted with EtOAc. The combined organic layer was washed with brine, dried and concentrated. The residue was purified on a flash column using

a gradient of 5-10% EtOAc/hexane to isolate the title compound.  $^1\text{H}$  NMR: (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.5-2.2 (m, 9H), 2.13 (s, 3H), 3.64 (m, 1H), 7.4-7.7 (m, 4H).

Step C 2-(3-Cyanophenyl)-3-cyclobutyl-1-methylpropylamine


This amine was prepared by following the method of Reference Example 2, Steps E-I.

5 LC-MS,  $R_t = 2.48$  min,  $m/e = 229$ .

The compounds of Reference Examples 91-93 were obtained by procedures described in Reference Example 90.

10

REFERENCE EXAMPLE 91




2-(3-Cyanophenyl)-3-cyclopropyl-1-methylpropylamine

LC-MS,  $R_t = 1.8$  min,  $m/e = 215$ .

15

REFERENCE EXAMPLE 92




2-(3-Cyanophenyl)-3-cyclopentyl-1-methylpropylamine

LC-MS,  $R_t = 2.7$  min,  $m/e = 243$ .

20

REFERENCE EXAMPLE 93




2-(3-Cyanophenyl)-3-cyclohexyl-1-methylpropylamine

LC-MS, Rt = 2.8 min, m/e = 257.

5

#### REFERENCE EXAMPLE 94



2-(3-Cyanophenyl)-3-(1-tert-butyloxycarbonyl-piperidin-4-yl)-1-methylpropylamine


Step A      3-(3-Cyanophenyl)-4-(1-tert-butyloxycarbonyl-piperidin-4-yl)-butan-2-one

10    The title compound was synthesized by the method of Reference Example 90, Steps A-B.

Step B      2-(3-Cyanophenyl)-3-(1-tert-butyloxycarbonyl-piperidin-4-yl)-1-methylpropylamine

15    The title amine was obtained by the method of Reference Example 2, steps E-G except that di-tert-butyl dicarbonate was not added in Step G. LC-MS, Rt = 2.72 min, m/e = 258 (M-99).

#### REFERENCE EXAMPLE 95



*N*-[3-(4-Chlorophenyl)-2-(3-methylthiophenyl)-1-methylpropyl]amine hydrochloride (Diastereomer  $\alpha$ )

The title compound was prepared following the same procedure as described in

- 5 Reference Example 55 substituting 3,5-dibromopyridine with 3-bromothioanisole at Step A. LC-MS:  $m/e$  306 ( $M + H$ )<sup>+</sup> (2.68 min).

### EXAMPLE 1

Automated Synthesis of a One Dimensional Amide Library

- 10 The following synthesis of a 1-dimensional, single, pure compound library was performed on a Myriad Core System. All reaction vessels were dried under a stream of nitrogen at 120°C for 12 hours prior to use. All solvents were dried over sieves for at least 12 hours prior to use. An appropriate stock solution of *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride (alpha isomer) was
- 15 prepared immediately prior to use in pyridine with 0.05 equivalents (relative to *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride (alpha isomer)) of dimethylaminopyridine added; the diversity carboxylic acids were dissolved immediately prior to use in DMSO. The relative amounts of reactants and coupling reagents are listed in Table 1. Compounds of the present invention that were prepared
- 20 by this method of automated synthesis are listed in Table 2.

Table 1.

Substance	Amount per reaction vessel	MW	Concentration	mmols	Equivs.
Aryl Acid in	1 mL	N/A	0.2 M	0.2	1.67

DMSO					
EDC/HOBt Cocktail in Deuterated Chloroform	0.8 mL	EDC: 191.71 HOBt: 135.13	0.25 M each	0.2 each	1.67 each
Amine in Pyridine with catalytic dimethylaminopyri dine (~0.05 eq.)	0.6 mL	294.227	0.2M	0.12	1.0

#### Procedure

To vessel one of a total of 192 dry, 10 mL fritted Myriad reaction vessels under nitrogen was added the appropriate diversity acid subunit (1.0 mL, 0.2 mmol, 0.2 M in DMSO); this was repeated for the remaining 191 reactions until the diversity acids had been enumerated to all 192 reaction vessels. To each of 192 reaction vessels under nitrogen was then added the EDC/HOBt cocktail (0.8 mL, 0.2 mmol, 0.25 M each in deuterated chloroform). Finally, to each of the 192 reaction vessels was added *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride (alpha isomer) (0.6 mL, 0.12 mmol, 0.2M in pyridine). The reactions were then aged for 4 hours at room temperature (20-25° C) followed by 16 hours at 65°C with nitrogen sparging agitation (1s pulse of nitrogen every 30 minutes.) The crude reactions were analyzed by HPLC-MS Method 1.

#### 15 Analytical LC Method 1:

Column: MetaChem Polaris C-18A, 30 mm X 4.6 mm, 5.0 µm  
 Eluent A: 0.1% TFA in Water  
 Eluent B: 0.1% TFA in Acetonitrile  
 Gradient: 5% B to 95 % B in 3.3 minutes, ramp back to 5% B in 0.3 min  
 Flow: 2.5 mL/min.  
 Column Temp.: 50°C  
 Injection amount: 5 µl of undiluted crude reaction mixture.  
 Detection: UV at 220 and 254 nm.

25 MS: API-ES ionization mode, mass scan range (100-700)

## ELSD: Light Scattering Detector

All 192 crude reactions were purified by preparative HPLC using UV based detection (Preparative method 2). The collected fractions were then analyzed for purity by LC-MS (Analytical method 3); fractions found to be greater than 90% purity were pooled into tared 40 mL EPA vials and lyophilized.

## Preparative LC Method 2:

Column: MetaChem Polaris C-18A, 100 mm X 21.2 mm, 10  $\mu$ m  
Eluent A: 0.1% TFA in Water  
Eluent B: 0.1% TFA in Acetonitrile  
Pre-inject Equilibration: 1.0 min  
Post-Inject Hold: 0.0 min  
Gradient: 10% B to 100 % B in 6.0 minutes, hold at 100% B for an additional 2.0 minutes, ramp back from 100% B to 10% B in 1.5 minutes.  
Flow: 25 mL/min.  
Column Temp.: ambient  
Injection amount: 1.5 ml of undiluted crude reaction mixture.  
Detection: UV at 220 and 254 nm.

## Analytical LC Method 3:

Column: MetaChem Polaris C-18A, 30 mm X 2.0 mm, 3.0  $\mu$ m  
Eluent A: 0.1% TFA in Water  
Eluent B: 0.1% TFA in Acetonitrile  
Gradient: 5% B to 95 % B in 2.0 minutes, ramp back to 5% B in 0.1 min  
Flow: 1.75 mL/min.  
Column Temp.: 60°C  
Injection amount: 5 ul of undiluted fraction.  
Detection: UV at 220 and 254 nm.

MS: API-ES ionization mode, mass scan range (100-700)

ELSD: Light Scattering Detector



## Lyophilization Parameters

Initial Freeze Setpoint: 1 hour at -70°C

Drying Phase Condenser Setpoint: -50°C

## Drying Phase Table:






Shelf Temperature (C)	Duration (minutes)	Vacuum Setpoint (mTorr)
-60°	240	25
-40°	240	25
5°	480	25
20°	1000	25




5







Table 2. The compounds in Table 2 were prepared by automated synthesis (the following compounds are racemic, and the structures imply relative stereochemistry only).



10 Table 2.






Exp. No.	Name	Structure	retention time (min)	HPLC- mass spectrum <i>m/e</i>
-------------	------	-----------	----------------------------	---






1.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-4-carboxamide, trifluoroacetic acid salt		1.094	419.1
2.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(2-pyridyl)-piperidine-3-carboxamide, trifluoroacetic acid salt		1.19	482.1
3.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-methylmorpholine-2-carboxamide, trifluoroacetic acid salt		1.116	421.2
4.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-pyrrolidine-2(S)-carboxamide		1.417	491.1
5.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-2-carboxamide, trifluoroacetic acid salt		1.134	419.1

6.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2,2-dimethyl-tetrahydropyran-4-carboxamide		1.40	434.3
7.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclopentane-carboxamide		1.51	494.3
8.	<i>cis</i> -N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide		1.502	508.2


9.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclohexane-carboxamide		1.455	508.2
10.	<i>cis</i> -N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide		1.53	508.3
11.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-cyclopentanone-3-carboxamide		1.271	404.1
12.	<i>trans</i> -N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide		1.462	508.2
13.	<i>trans</i> -N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide		1.56	508.3
14.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-2-methylazetidine-2-carboxamide		1.479	513.15 (+Na <sup>+</sup> )

15.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzyl-piperidine-2-carboxamide, trifluoroacetic acid salt		1.240	495.1
16.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)pyrrolidine-3-carboxamide		1.415	391.1 (-BOC)

17.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide		1.51	438.2
18.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-pyrrolidine-2(S)-carboxamide		1.529	601.0
19.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-2-methylpyrrolidine-2(S)-carboxamide		1.584	614.1
20.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-methyltetrahydrofuran-2-carboxamide		1.374	406.2
21.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-tetrahydrofuran-2-carboxamide		1.326	392.1

22.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide, trifluoroacetic acid salt		1.194	467.1
23.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-oxopyrrolidine-5-carboxamide		1.181	405.15
24.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-5-methyl-2-oxopyrrolidine-5-carboxamide		1.25	419.2
25.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzylpyrrolidine-2-carboxamide		1.24	481.15
26.	N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(phenylamino)-cyclopentane-carboxamide, trifluoroacetic acid salt		1.57	481.3


EXAMPLE 27



*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopentanecarboxamide

To a solution of phenylcyclopentanecarboxylic acid (Aldrich, 23 mg, 0.12 mmol) in methylene chloride (1 mL) at 0°C was added a drop of dimethylformamide and oxalyl chloride (0.025 mL, 0.29 mmol). After stirring at room temperature for 1 h, the reaction mixture was concentrated on a rotary evaporator and dried under vacuum, and the resulting crude acyl chloride was used without further purification. The crude acyl chloride was dissolved in 1 mL of methylene chloride and was added to a suspension of *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-amine hydrochloride (Reference Example 2) (diastereomer  $\alpha$ , 30 mg, 0.10 mmol) and *N*-methylmorpholine (0.053 mL, 0.48 mmol) in 1 mL of methylene chloride. After stirring at room temperature overnight, the reaction mixture was loaded onto a silica gel column, which was eluted with 10% ethyl acetate to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.45 (d, 2H), 7.35-7.08 (m, 6H), 6.97 (d, 2H), 6.90 (d, 2H), 6.59 (d, 2H), 4.19 (m, 1H), 2.74-2.45 (m, 5H), 2.12-1.65 (m, 6H), 0.79 (d, 3H). LC-MS: *m/e* 432 (*M* + *H*)<sup>+</sup> (4.4 min).

EXAMPLE 28




*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopropanecarboxamide

Phenylcyclopropanecarboxylic acid (Aldrich, 20 mg, 0.12 mmol) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound. <sup>1</sup>H NMR (400



MHz, CD<sub>3</sub>OD):  $\delta$  7.42-7.00 (m, 10 H), 6.95 (d, 2H), 6.85 (d, 2H), 4.17 (m, 1H), 3.02-2.70 (m, 3H), 1.6-1.0 (m, 6H), 0.92 (d, 3H). LC-MS:  $m/e$  404 (M + H)<sup>+</sup> (4.2 min).

### EXAMPLE 29




5

#### N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopropanecarboxamide

1-(3-Pyridyl)cyclopropanecarboxylic acid (prepared following the procedure of Wilt and Philip *J. Org. Chem.* **1959**, 24, 616) (34 mg, 0.21 mmol) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  8.60 (s, 1H), 8.49 (d, 1H), 7.85 (dd, 1H), 7.42 (dd, 1H), 7.22-6.95 (m, 7H), 6.84 (d, 2H), 4.22 (m, 1H), 3.00 (dd, 1H), 2.88-2.70 (m, 2H), 1.64-1.08 (m, 6H), 0.84 (d, 3H). LC-MS:  $m/e$  405 (M + H)<sup>+</sup> (2.7 min).

15

### EXAMPLE 30




#### N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-chlorophenyl)cyclohexanecarboxamide

1-(3-Chlorophenyl)cyclohexanecarboxylic acid (prepared following the procedure of Wilt and Philip *J. Org. Chem.* **1959**, 24, 616) (50 mg, 0.21 mmol) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound. <sup>1</sup>H NMR (400

20

MHz, CD<sub>3</sub>OD):  $\delta$  7.52-6.90 (m, 12H), 6.59 (d, 2H), 4.24 (m, 1H), 2.72-2.42 (m, 5H), 2.0-1.2 (m, 18H), 0.80 (d, 3H). LC-MS: m/e 480 (M + H)<sup>+</sup> (4.7 min).

### EXAMPLES 31 AND 32




5

#### N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-2-phenyltetrahydrofuran-2-carboxamide (Diastereomers I and II)

2-Phenyltetrahydrofuran-2-carboxylic acid (prepared according to procedures described in W. K. Hagmann et al., PCT Patent Application  
 10 WO2001/12183A1 (Feb. 22, 2001); *Chem. Abstracts* **134**(14):193737n) (50 mg, 0.26 mmol) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound as a mixture of diastereomers which were separated by flash column chromatography on silica gel eluted with 10% ethyl acetate in hexane to give a faster eluting isomer and a slower  
 15 eluting isomer.  
 Faster eluting isomer: <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.58 (d, 2H), 7.35-7.02 (m, 10H), 6.84 (d, 2H), 4.2-4.0 (m, 3H), 3.08 (dd, 1H), 2.97 (m, 1H), 2.85-1.92 (m, 5H), 0.82 (d, 3H). LC-MS: m/e 434 (M + H)<sup>+</sup> (4.2 min).  
 Slower eluting isomer: <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.65-6.90 (m, 12H), 6.59 (d,  
 20 2H), 4.2-4.0 (m, 3H), 2.85 (m, 1H), 2.75 (m, 1H), 2.62 (dd, 1H), 2.50 (dd, 1H), 2.18 (m, 1H), 1.95 (m, 2H), 0.90 (d, 3H). LC-MS: m/e 434 (M + H)<sup>+</sup> (4.2 min).

### EXAMPLE 33



N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3,5 difluorophenyl)-cyclopentane-carboxamide

Step A 1-(3,5-Difluorophenyl)cyclopentanecarboxylic acid, methyl ester

- 5 To a solution of 3,5-difluorophenylacetic acid (5.0 g, 29 mmol) in methylene chloride (50 mL) and methanol (50 mL) at 0°C was added trimethylsilyldiazomethane (2 M in hexane) until a yellow color persisted. After stirring at room temperature for 15 min, the reaction mixture was concentrated to dryness, and the residue was dried by azeotroping with toluene to give the corresponding methyl ester (5.5 g), which was
- 10 used without further purification. Thus, to a suspension of potassium hydride (35% in mineral oil, 3.7 g, 32 mmol, washed 2 times with hexane to remove oil) in 5 mL of anhydrous tetrahydrofuran at 0°C was added a solution of the crude methyl ester (2.0 g, 11 mmol) in 40 mL of anhydrous tetrahydrofuran. After the bubbling subsided, the reaction was stirred at room temperature for 15 min. The reaction mixture was then
- 15 cooled to 0°C, and was added 1,4-dibromobutane (3.9 mL, 32 mmol) in 5 mL of tetrahydrofuran. After stirring at room temperature for 20 min, the reaction mixture was partitioned between ethyl acetate (100 mL) and saturated aqueous ammonium chloride (100 mL). The organic layer was separated, washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was
- 20 purified by flash column chromatography on silica gel eluted with 1-5% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 6.95 (m, 2H), 6.81 (m, 1H), 3.62 (s, 3H), 2.60 (m, 2H), 1.84 (m, 2H), 1.75 (m, 4H).

Step B 1-(3,5-Difluorophenyl)cyclopentanecarboxylic acid


- To a solution of 1-(3,5-difluorophenyl)cyclopentanecarboxylic acid methyl ester (Step
- 25 A, 2.0 g, 8.3 mmol) in dimethylsulfoxide (5 mL) was added potassium hydroxide (1.9 g, 33 mol) in 2 mL of water. After stirring at room temperature for 2.5 h, the reaction mixture was partitioned between ether (50 mL) and saturated aqueous hydrochloric acid (2 N, 50 mL). The organic layer was separated and the aqueous layer extracted

with ether (2 x 50 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 6.97 (m, 2H), 6.80 (m, 1H), 2.60 (m, 2H), 1.84 (m, 2H), 1.74 (m, 4H).

5 Step C N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3,5-difluorophenyl)cyclopentanecarboxamide

1-(3,5-Difluorophenyl)cyclopentanecarboxylic acid (Step B, 0.10 g, 0.44 mmol) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound. <sup>1</sup>H NMR (400  
10 MHz, CD<sub>3</sub>OD): δ 7.20-6.75 (m, 10H), 6.65 (d, 2H), 4.18 (m, 1H), 2.76-2.48 (m, 5H), 2.05-1.68 (m, 6H). LC-MS: m/e 468 (M + H)<sup>+</sup> (4.5 min).

EXAMPLE 34



15 N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(4-pyridyl)cyclopentanecarboxamide

Step A Methyl 4-pyridylacetate

To a suspension of methyl 4-pyridylacetate hydrochloride salt (6.0 g, 32 mmol) in ethyl acetate (100 mL) and brine (50 mL) was added sodium carbonate (6.8 g, 64  
20 mmol) in 25 mL of water. After stirring at room temperature for 15 min, the organic layer was separated and the aqueous layer extracted with ethyl acetate (3 x 100 mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give the title compound.

Step B Benzyl 4-Pyridylacetate

25 To a solution of methyl 4-pyridylacetate (Step A, 2.5 g, 16 mmol) and benzyl alcohol (3.4 mL, 33 mmol) in anhydrous tetrahydrofuran (10 mL) at -10 °C was added butyllithium (2 M in cyclohexane, 16 mL, 33 mL), and the reaction was allowed to warm to room temperature overnight. The volatile materials were removed on a rotary evaporator, and the residue was diluted with methylene chloride (50 mL),

washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated to dryness, and the residue was purified by flash column chromatography eluted with 30% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.43 (d, 2H), 7.38 (d, 2H), 7.32 (m, 5 H), 5.16 (s, 2H), 3.99 (s, 2H).

Step C 1-(4-Pyridyl)cyclopentanecarboxylic acid, benzyl ester

Benzyl 4-pyridylacetate (Step B, 0.80 g, 3.5 mmol) was reacted with potassium hydride and 1,4-dibromobutane according to the procedures described in Example 33, Step A to afford the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): 8.42 (d, 2H), 7.39 (d, 2H), 7.28 (m, 3 H), 7.19 (m, 2H), 5.06 (s, 2H), 2.62 (m, 2H), 1.97 (m, 2H), 1.74 (m, 4H). LC-MS: m/e 282 (M + H)<sup>+</sup> (2.2 min).


Step D 1-(4-Pyridyl)cyclopentanecarboxylic acid

To a solution of 1-(4-pyridyl)cyclopentanecarboxylic acid, benzyl ester (Step C, 0.36 g, 1.3 mmol) in methanol (10 mL) was added palladium on carbon (10% w/w, 75 mg). The mixture was degassed and filled with hydrogen using a balloon. After stirring overnight, the reaction mixture was filtered through Celite, and the filtrate was concentrated to dryness to afford the title compound. LC-MS: m/e 192 (M + H)<sup>+</sup> (0.8 min).

Step E N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(4-pyridyl)cyclopentanecarboxamide

1-(4-Pyridyl)cyclopentanecarboxylic acid (Step D) was reacted according to the procedures described in Example 27 substituting for phenylcyclopentanecarboxylic acid to afford the title compound after purification by reverse-phase HPLC eluted with 50-100% water/acetonitrile (0.1% trifluoroacetic acid). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.78 (d, 2H), 7.72 (d, 2H), 7.2-7.1 (m, 3H), 7.00 (d, 2H), 6.75 (d, 2H), 6.64 (d, 2H), 4.21 (m, 1H), 2.8-0.8 (m, 11H), 0.82 (d, 3H) LC-MS: m/e 433 (M + H)<sup>+</sup> (3.0 min).

EXAMPLE 35




*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentane-carboxamide

- The title compound was prepared as a mixture of enantiomers according to the procedures described for Example 34, Steps B through E substituting 4-pyridylacetate with 3-pyridylacetate in Step B. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.65 (d, 1H), 8.40 (d, 1H), 7.92 (dd, 1H), 7.40 (dd, 1H), 7.20-7.06 (m, 3H), 7.00 (d, 2H), 6.93 (d, 2H), 6.62 (d, 2H), 4.20 (m, 1H), 2.75-2.55 (m, 5H), 2.10-1.70 (m, 6H), 0.80 (d, 3H). LC-MS: *m/e* 433 (M + H)<sup>+</sup> (3.0 min).

10


EXAMPLES 36 AND 37



*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentane-carboxamide (Enantiomers A and B)

- 15 The product of Example 35 was separated into its pure enantiomers by chromatography on a ChiralCel OD column eluted with 15% ethanol in hexane. Faster eluting enantiomer (Enantiomer A): Analytical HPLC: retention time = 5.7 min (ChiralCel OD column, flow rate = 0.75 mL/min, 15% ethanol/hexane). LC-MS: *m/e* 433 (M + H)<sup>+</sup> (3.0 min).
- 20 Slower eluting enantiomer (Enantiomer B): Analytical HPLC: retention time = 6.4 min (ChiralCel OD column, flow rate = 0.75 mL/min, 15% ethanol/hexane). LC-MS: *m/e* 433 (M + H)<sup>+</sup> (3.0 min).


EXAMPLE 38



*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl-N-oxide)-cyclopentane-carboxamide

To a solution *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide (Example 37, slower eluting enantiomer B, 9.0 mg, 0.021 mmol) in 0.5 mL of methylene chloride was added *m*-chloroperbenzoic acid (50%, 5.4 mg, 0.031 mmol). After stirring at room temperature for 1 h, the reaction mixture was concentrated to dryness, and the residue was dissolved in dimethylsulfoxide, acetonitrile and water (2:2:1, 2 mL) and loaded onto a reverse-phase semi-preparative HPLC column eluted with a gradient of 75% water/ acetonitrile (0.1% trifluoroacetic acid) to 100% acetonitrile (0.1% trifluoroacetic acid) to yield the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.41 (s, 1H), 8.24 (d, 1H), 7.70-7.08 (m, 5H), 7.02 (d, 2H), 6.96 (d, 2H), 6.72 (d, 2H), 4.20 (m, 1H), 2.82-2.55 (m, 5H), 2.10-1.70 (m, 6H), 0.82 (d, 3H). LC-MS: *m/e* 449 (M + H)<sup>+</sup> (3.2 min).

EXAMPLE 39



*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentane-carboxamide


Step A 1-(2-Pyridyl)cyclopentanecarboxylic acid, benzyl ester

The title compound was prepared as a mixture of enantiomers according to the procedures described for Example 34, Steps B through D substituting 2-pyridylacetate with 3-pyridylacetate in Step B. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.42 (d, 1H), 7.75 (t, 1H), 7.40 (d, 1H), 7.24 (m, 4 H), 7.18 (m, 2H), 5.06 (s, 2H), 2.54 (m, 2H), 2.15 (m, 2H), 1.73 (m, 4H). LC-MS: m/e 282 (M + H)<sup>+</sup> (2.4 min).

Step B N-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide

To a mixture of 1-(2-pyridyl)cyclopentanecarboxylic acid, benzyl ester (Step A, 0.20 g, 0.71 mmol) in acetonitrile (3 mL) and water (1.5 mL) was added lithium hydroxide monohydrate (0.24 g, 5.6 mmol). After stirring at room temperature for 5 h and 50°C overnight, the reaction mixture was concentrated to dryness. The residue was azeotroped with toluene and was suspended in methylene chloride (6 mL) and dimethylformamide (3 mL), and was added N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-amine hydrochloride (Diastereomer α, Reference Example 2, 0.40 g, 0.14 mmol), 4-dimethylaminopyridine (0.10 g, 0.71 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide (38 mg, 0.19 mmol). After stirring at room temperature overnight, the reaction mixture was diluted with ethyl acetate (50 mL), washed with water and brine, dried over anhydrous sodium sulfate and concentrated to dryness. N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide was obtained by flash column chromatography on silica gel eluted with 15-20% ethyl acetate in hexane. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 8.52 (d, 1H), 7.77 (t, 1H), 7.48 (d, 1H), 7.35-7.08 (m, 4H), 7.00 (d, 2H), 6.94 (d, 2H), 6.68 (d, 2H), 4.20 (m, 1H), 2.80 (dd, 1H), 2.73 (m, 1H), 2.60 (m, 2H), 2.45 (m, 1H), 2.30 (m, 1H), 2.18 (m, 1H), 1.85-1.65 (m, 4H), 0.82 (d, 3H). LC-MS: m/e 433 (M + H)<sup>+</sup> (3.2 min).

EXAMPLE 40





*N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(5-bromo-3-pyridyl)-cyclopentanecarboxamide

Step A 1-(5-Bromo-3-pyridyl)-cyclopentanecarboxylic acid, benzyl ester


The title compound was prepared according to the procedures described for Example 34, Steps A through C substituting methyl 4-pyridylacetate with methyl 5-bromo-3-pyridylacetate and was used in the subsequent reaction.

Step B *N*-[3-(4-Chlorophenyl)-2-phenyl-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide

To a solution of 1-(5-bromo-3-pyridyl)-cyclopentanecarboxylic acid, benzyl ester (Step 140 mg, 0.11 mmol) in methanol (1 mL)/tetrahydrofuran (0.5 mL)/water (1.5 mL) was added lithium hydroxide monohydrate (19 mg, 0.44 mmol). After stirring at room temperature for 2 h, the reaction mixture was concentrated to dryness, and the residue was azeotroped with toluene to give the crude acid as the lithium salt, which was used without further modification. The lithium salt was suspended in 2 mL of methylene chloride, and was added *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-amine hydrochloride (Diastereomer  $\alpha$ , 33 mg, 0.11 mmol), *N*-methylmorpholine (0.049 mL, 0.44 mmol) and tris(pyrrolidiny)phosphonium hexafluorophosphate (87 mg, 0.17 mmol). After stirring at room temperature overnight, the reaction mixture was loaded onto a silica gel column eluted with 20% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  8.60 (s, 1H), 8.55 (s, 1H), 8.08 (s, 1H), 7.20-7.10 (m, 3H), 7.01 (d, 2H), 6.95 (d, 2H), 6.64 (d, 2H), 4.20 (m, 1H), 2.80-2.55 (m, 5H), 2.10-1.70 (m, 6H), 0.82 (d, 3H). LC-MS: *m/e* 512 (M + H)<sup>+</sup> (4.1 min).

25

#### EXAMPLE 41



*N*-[2,3-Bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenoxy)-cyclohexanecarboxamide

To a mixture of *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride (Diastereomer  $\alpha$ , Reference Example 1) (60 mg, 0.19 mmol) and 1-(4-chlorophenyl)oxy)cyclohexanecarboxylic acid (60 mg, 0.24 mmol) in 1 mL of methylene chloride was added *N,N*-diisopropylethylamine (0.10 mL, 0.58 mmol) and tris(pyrrolidiny)phosphonium hexafluorophosphate (0.10 g, 0.19 mmol). After stirring at room temperature overnight, the reaction mixture was loaded onto a silica gel column eluted with 6:1 hexane/ether to afford the title compound.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.22 (d, 2H), 7.19 (d, 2H), 7.03 (d, 2H), 6.99 (d, 2H), 6.91 (d, 2H), 6.71 (d, 2H), 4.24 (m, 1H), 2.99 (dd, 1H), 2.80 (ddd, 1H), 2.63 (dd, 1H), 2.3-1.2 (m, 10H), 0.82 (d, 3H). LC-MS:  $m/e$  530 ( $M + H$ ) $^+$  (5.0 min).



The compounds in Table 3 were prepared according to the procedures described in Example 41 substituting 1-(4-chlorophenyl)oxy)cyclohexanecarboxylic acid with the appropriate carboxylic acid.

Table 3.

Exp. No.	Name	Structure	retention time (min)	HPLC-mass spectrum $m/e$
42.	<i>N</i> -[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide		3.9	420
43.	<i>N</i> -[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-( <i>tert</i> -butoxycarbonylamino)cyclohexane-carboxamide		4.4	519
44.	<i>N</i> -[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-( <i>tert</i> -butoxycarbonylamino)cyclopentane-carboxamide		4.3	505

The compounds in Table 4 were prepared from compounds 43 and 44 by reaction with hydrogen chloride in dioxane (4 M) overnight at room temperature to afford the corresponding amine hydrochloride salts (Examples 45 and 46, respectively).

5 Table 4.

Exp. No.	Name	Structure	retention time (min)	HPLC-mass spectrum <i>m/e</i>
45.	<i>N</i> -[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclohexanecarboxamide hydrochloride salt		2.8	419
46.	<i>N</i> -[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclopentanecarboxamide hydrochloride salt		2.8	405

The compounds in Table 5 may be prepared according to the procedures described in Example 41 substituting the appropriate amine from Reference Examples 4-8 for *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride.



10 Table 5.





Exp. No.	Name
47.	<i>N</i> -[2-(3-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
48.	<i>N</i> -[2-(3-fluoropyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
49.	<i>N</i> -[2-(2-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;





50.	<i>N</i> -[2-(4-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
51.	<i>N</i> -[3-(5-chloro-2-pyridyl)-2-phenyl-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;





The compounds in Table 6 were prepared according to the procedures described in Example 41 substituting *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride and 1-(4-chlorophenyloxy)cyclohexanecarboxylic acid with the appropriate amine and carboxylic acid.


Table 6.

Exp. No.	Name	Structure	retention time (min)	HPLC-mass spectrum <i>m/e</i>
52.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide		3.8	536
53.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide		4.1	443

54.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide		4.2	475
55.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide		4.2	475
56.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide		4.3	475
57.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide		4.3	509

58.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide		4.3	491
59.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide		4.5	471
60.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide		4.2	487
61.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide		3.9	420



62.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentane carboxamide		4.2	457
63.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide		3.8	458
64.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexane carboxamide		4.3	471
65.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide		4.3	509

66.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide		4.4	525
-----	---	--	-----	-----






The compounds in Table 7 were prepared according to the procedures described in Example 41 substituting *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-amine hydrochloride and 1-(4-chlorophenyloxy)cyclohexanecarboxylic acid with the appropriate enantiomer of the amine and the appropriate carboxylic acid.





5



Table 7.

Exp. No.	Name	Structure	retention time (min)	HPLC-mass spectrum <i>m/e</i>
67.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclobutanecarboxamide		4.4	418
68.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide		4.1	443




69.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentane carboxamide		3.3	458
70.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-cyclobutanecarboxamide		3.6	367
71.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylcyclobutanecarboxamide		3.7	381
72.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-ethylcyclobutanecarboxamide		3.8	395
73.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-propylcyclobutanecarboxamide		4.0	409






74.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-benzylcyclobutanecarboxamide		4.0	457
75.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-isopropylcyclobutanecarboxamide		4.0	409
76.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-3-azetidine-3-carboxamide		3.8	468
77.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2( <i>S</i> )-azetidinecarboxamide		3.8	468







78.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 1)		4.2	396 (M-Boc) <sup>+</sup>
79.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 2)		4.1	496







The compounds in Table 8 were isolated as single enantiomers from the corresponding racemic material (Table 6) following the procedures described in Example 37 with appropriate modifications of (1) Chiralcel OD or Chiralpak AD column, (2) the eluent composition (4-15% ethanol/hexane), (3) flow rate (6-9 mL/min) and (4) injection volume (200 to 2000  $\mu$ L).







Table 8.


Ex. No.	Name	Structure	retention time (min)	HPLC-mass spectrum <i>m/e</i>	Enantiomer A or B
80.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentane carboxamide		3.8	536	A

81.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentane carboxamide		3.8	536	B
82.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide		4.3	475	A
83.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide		4.3	475	B
84.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide		4.2	475	A
85.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide		4.2	475	B

86.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide		4.3	475	A
87.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide		4.4	475	B
88.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide		4.3	509	A
89.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide		4.3	509	B
90.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide		4.3	491	A
91.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide		4.3	491	B

92.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide		4.3	471	A
93.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide		4.3	471	B
94.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide		4.1	487	A
95.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide		4.1	487	B
96.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide		4.2	457	A
97.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide		4.2	457	B

98.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide		4.3	471	A
99.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide		4.3	471	B
100.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide		4.3	509	A
101.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide		4.3	509	B
102.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide		4.4	525	A
103.	<i>N</i> -[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide		4.4	525	B

EXAMPLE 103

5 *N*-(2*S*,3*S*)-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylsulfonylcyclobutanecarboxamide

Step A Methyl 1-Methylsulfonylcyclobutanecarboxylate

To a solution of methyl methylsulfonylacetate (1.0 g, 6.6 mmol) in 20 mL of acetonitrile was added 1,3-diiodopropane (2.3 mL, 20 mmol) and cesium carbonate (6.4 g, 20 mmol). After stirring at room temperature for 2 h, the reaction was partitioned between water (50 mL) and ethyl acetate (100 mL). The organic layer was separated, dried over anhydrous magnesium sulfate, filtered and concentrated to dryness, and the residue was purified by flash chromatography on silica gel eluting with 20 to 50% ethyl acetate in hexane to give methyl 1-methylsulfonylcyclobutanecarboxylate (1.0 g, 79%).  $\delta$  3.84 (s, 3H), 3.02 (s, 3H), 2.84-2.65 (m, 4H), 2.25-2.02 (m, 2H).


Step B *N*-(2*S*,3*S*)-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylsulfonylcyclobutanecarboxamide

To a solution of methyl 1-methylsulfonylcyclobutanecarboxylate (0.10 g, 0.52 mmol) in 2 mL of dioxane and 2 mL of water was added lithium hydroxide monohydrate (44 mg, 1.0 mmol). After stirring at 100°C for 1 h, the reaction mixture was concentrated to dryness, and the residue was azeotroped with toluene to give the crude lithium 1-methylsulfonylcyclobutanecarboxylate, which was used without further purification. Thus, the crude lithium salt was suspended in 2 mL of anhydrous dimethylformamide, and was added *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-amine hydrochloride (0.15 g, 0.52 mmol), tris(pyrrolidiny)phosphonium hexafluorophosphate (0.41 g, 0.78 mmol) and *N*-methylmorpholine (0.23 mL, 2.1



mmol). After stirring at room temperature overnight, the reaction mixture was partitioned between water (20 mL) and ether (20 mL). The organic layer was separated, and the aqueous layer was extracted with ether (20 mL). The combined extracts were dried over anhydrous magnesium sulfate, filtered and concentrated to dryness, and the residue was purified by flash chromatography on silica gel eluting with 20 to 40% ethyl acetate in hexane to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.57 (m, 1H), 7.49 (s, 1H), 7.43 (m, 2H), 7.10 (d, 2H), 6.92 (d, 2H), 4.38 (m, 1H), 3.35-2.63 (m, 7H), 2.93 (s, 3H), 2.14 (m, 1H), 1.98 (m, 1H), 0.98 (d, 3H). LC-MS: m/e 445 (M + H)<sup>+</sup> (3.5 min).

10

EXAMPLE 104

*N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-3-ethylazetidine-3-carboxamide hydrochloride

15 Step A *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-tert-butoxycarbonyl-3-ethylazetidine-3-carboxamide

The title compound was prepared from 1-tert-butoxycarbonyl-3-ethylazetidine-3-carboxylic acid (Reference Example 12) and *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-amine hydrochloride following the procedure described in Example 103. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 7.54 (m, 1H), 7.48 (s, 1H), 7.42 (m, 2H), 7.12 (d, 2H), 6.95 (d, 2H), 4.32 (m, 1H), 4.12 (m, 2H), 3.74 (m, 2H), 3.16 (dd, 1H), 3.05 (m, 1H), 2.84 (dd, 1H), 1.96 (q, 2H), 1.42 (q, 9H), 0.97 (d, 3H), 0.93 (t, 3H). LC-MS: m/e 518 (M + Na)<sup>+</sup> (3.9 min).



25 Step B *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-3-ethylazetidine-3-carboxamide hydrochloride

To a sample of *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-tert-butoxycarbonyl-3-ethylazetidine-3-carboxamide (Step A, 0.18 g, 0.36 mmol) was

added 4 M hydrogen chloride in dioxane (10 mL). After stirring at room temperature for 1 h, the reaction mixture was concentrated to dryness, and the residue was triturated with ether and dried under vacuum to give the title compound. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.14 (d, 1H), 7.55 (m, 1H), 7.49 (s, 1H), 7.43 (m, 2H), 7.13 (d, 2H), 6.95 (d, 2H), 4.35 (m, 2H), 4.32 (m, 1H), 3.94 (m, 2H), 3.21 (dd, 1H), 3.07 (m, 1H), 2.84 (dd, 1H), 2.08 (q, 2H), 0.99 (d, 3H), 0.94 (t, 3H). LC-MS: m/e 396 (M + H)<sup>+</sup> (2.8 min).

The compounds in Table 9 were prepared from compounds 78 and 79 by reaction with hydrogen chloride in dioxane (4 M) as described in Example 104 to afford the corresponding amine hydrochloride salts.

Table 9.

Exp. No.	Name	Structure	retention time (min)	HPLC-mass spectrum m/e
105.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 1)		2.8	419
106.	<i>N</i> -(2 <i>S</i> ,3 <i>S</i> )-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 2)		2.8	419

Cannabinoid Receptor-1 (CB1) Binding Assay.

Binding affinity determination is based on recombinant human CB1 receptor expressed in Chinese Hamster Ovary (CHO) cells (Felder et al, Mol. Pharmacol. 48: 443-450, 1995). Total assay volume is 250  $\mu$ l (240  $\mu$ l CB1 receptor membrane solution plus 5  $\mu$ l test compound solution plus 5  $\mu$ l [3H]CP-55940 solution). Final concentration of [3H]CP-55940 is 0.6 nM. Binding buffer contains 50mM Tris-HCl, pH7.4, 2.5 mM EDTA, 5mM MgCl<sub>2</sub>, 0.5mg/ml fatty acid free bovine serum albumin and protease inhibitors (Cat#P8340, from Sigma). To initiate the binding reaction, 5  $\mu$ l of radioligand solution is added, the mixture is incubated with gentle shaking on a shaker for 1.5 hours at 30°C. The binding is terminated by using 96-well harvester and filtering through GF/C filter presoaked in 0.05% polyethylenimine. The bound radiolabel is quantitated using scintillation counter. Apparent binding affinities for various compounds are calculated from IC<sub>50</sub> values (DeBlasi et al., Trends Pharmacol Sci 10: 227-229, 1989).

The binding assay for CB2 receptor is done similarly with recombinant human CB2 receptor expressed in CHO cells.

EXAMPLE 108Cannabinoid Receptor-1 (CB1) Functional Activity Assay.

The functional activation of CB1 receptor is based on recombinant human CB1 receptor expressed in CHO cells (Felder et al, Mol. Pharmacol. 48: 443-450, 1995). To determine the agonist activity or inverse agonist activity of any test compound, 50  $\mu$ l of CB1-CHO cell suspension are mixed with test compound and 70  $\mu$ l assay buffer containing 0.34 mM 3-isobutyl-1-methylxanthine and 5.1  $\mu$ M of forskolin in 96-well plates. The assay buffer is comprised of Earle's Balanced Salt Solution supplemented with 5 mM MgCl<sub>2</sub>, 1 mM glutamine, 10 mM HEPES, and 1 mg/ml bovine serum albumin. The mixture is incubated at room temperature for 30 minutes, and terminated by adding 30 $\mu$ l/well of 0.5M HCl. The total intracellular cAMP level is quantitated using the New England Nuclear Flashplate and cAMP radioimmunoassay kit.


To determine the antagonist activity of test compound, the reaction mixture also contains 0.5 nM of the agonist CP55940, and the reversal of the CP55940 effect is quantitated. Alternatively, a series of dose response curves for CP55940 is performed with increasing concentration of the test compound in each of the dose response curves.

The functional assay for the CB2 receptor is done similarly with recombinant human CB2 receptor expressed in CHO cells.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that  
5 various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications for the compounds of the invention indicated above. Likewise, the  
10 specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended,  
15 therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

## WHAT IS CLAIMED IS:

1. A compound of structural formula I:



5

(I)

or a pharmaceutically acceptable salt thereof, wherein:

R<sup>1</sup> is selected from:

- (1) C<sub>1-10</sub>alkyl,
- (2) C<sub>3-10</sub>cycloalkyl,
- 10 (3) cycloheteroalkyl,
- (4) aryl, and
- (5) heteroaryl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

15

R<sup>2</sup> is selected from:

- (1) C<sub>3-10</sub>cycloalkyl,
- (2) cycloheteroalkyl,
- 20 (3) aryl,
- (4) heteroaryl,
- (5) -OR<sup>d</sup>,
- (6) -NRCR<sup>d</sup>, and
- (7) -CO<sub>2</sub>R<sup>d</sup>,

25

wherein each alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>, and each cycloalkyl, and cycloheteroalkyl aryl and heteroaryl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

R<sup>3</sup> is selected from:

- 30 (1) hydrogen, and

(2) C<sub>1-4</sub>alkyl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>;

R<sup>4</sup> is selected from:

- 5 (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>2-10</sub>alkenyl,
- (4) C<sub>2-10</sub>alkynyl,
- (5) -OR<sup>c</sup>,
- 10 (6) -CO<sub>2</sub>R<sup>c</sup>,
- (7) -OCOOR<sup>c</sup>,
- (8) -CONR<sup>c</sup>R<sup>d</sup>,
- (9) -NR<sup>c</sup>R<sup>d</sup>,
- (10) -NR<sup>c</sup>(CO)OR<sup>d</sup>,
- 15 (11) -NR<sup>c</sup>SO<sub>2</sub>R<sup>d</sup>,
- (12) -S(O)<sub>m</sub>R<sup>c</sup>,
- (13) aryl, and
- (14) heteroaryl,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one, two, three or  
20 four substituents independently selected from R<sup>a</sup>, and aryl and heteroaryl are  
optionally substituted with one, two, three or four substituents independently selected  
from R<sup>b</sup>;

R<sup>6</sup> is selected from:

- (1) hydrogen,
- 25 (2) C<sub>1-4</sub>alkyl,
- (3) C<sub>2-4</sub>alkenyl,
- (4) C<sub>2-4</sub>alkynyl,
- (5) -OR<sup>d</sup>,
- (6) halogen,
- 30 (7) -CN,
- (8) -NR<sup>c</sup>R<sup>d</sup>,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four  
substituents independently selected from R<sup>a</sup>;

**A** is a 3- to 8-membered monocyclic saturated ring incorporating the same carbon atom to which  $R^4$  is attached, and optionally containing one to two heteroatoms chosen from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted with one, two, three or four substituents independently selected from  $R^b$ ;

5 each  $R^a$  is independently selected from:

- (1)  $-OR^c$ ,
- (2)  $-NR^cS(O)_mR^d$ ,
- (3)  $-NO_2$ ,
- (4) halogen,
- 10 (5)  $-S(O)_mR^c$ ,
- (6)  $-SR^c$ ,
- (7)  $-S(O)_2OR^c$ ,
- (8)  $-S(O)_mNR^cR^d$ ,
- (9)  $-NR^cR^d$ ,
- 15 (10)  $-O(CR^eR^f)_nNR^cR^d$ ,
- (11)  $-C(O)R^c$ ,
- (12)  $-CO_2R^i$ ,
- (13)  $-CO_2(CR^eR^f)_nCONR^cR^d$ ,
- (14)  $-OC(O)R^c$ ,
- 20 (15)  $-CN$ ,
- (16)  $-C(O)NR^cR^d$ ,
- (17)  $-NR^cC(O)R^d$ ,
- (18)  $-OC(O)NR^cR^d$ ,
- (19)  $-NR^cC(O)OR^d$ ,
- 25 (20)  $-NR^cC(O)NR^cR^d$ ,
- (21)  $-CR^c(N-OR^d)$ ,
- (22)  $CF_3$ ,
- (23)  $-OCF_3$ ,
- (24)  $C_{3-8}$ cycloalkyl,
- 30 (25) cycloheteroalkyl, and
- (26) oxo;

each  $R^b$  is independently selected from:

- (1)  $R^a$ ,
- (2)  $C_{1-10}$ alkyl,

- (3) C<sub>3-8</sub>cycloalkyl,
- (4) cycloheteroalkyl,
- (5) aryl,
- (6) arylC<sub>1-4</sub>alkyl,
- 5 (7) heteroaryl, and
- (8) heteroarylC<sub>1-4</sub>alkyl;

each R<sup>c</sup> and R<sup>d</sup> are independently selected from:

- (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- 10 (3) C<sub>2-10</sub>alkenyl,
- (4) C<sub>2-10</sub>alkynyl,
- (5) cycloalkyl,
- (6) cycloalkyl-C<sub>1-10</sub>alkyl,
- (7) cycloheteroalkyl,
- 15 (8) cycloheteroalkyl-C<sub>1-10</sub>alkyl,
- (9) aryl,
- (10) heteroaryl,
- (11) aryl-C<sub>1-10</sub>alkyl, and
- (12) heteroaryl-C<sub>1-10</sub>alkyl, or

- 20 R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sub>g</sub>,

each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three substituents selected from R<sup>h</sup>;

- 25 each R<sup>e</sup> and R<sup>f</sup> are independently selected from:

- (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>2-10</sub>alkenyl,
- (4) C<sub>2-10</sub>alkynyl;
- 30 (5) cycloalkyl,
- (6) cycloalkyl-C<sub>1-10</sub>alkyl,
- (7) cycloheteroalkyl,
- (8) cycloheteroalkyl-C<sub>1-10</sub>alkyl,
- (9) aryl,



- (10) heteroaryl,
- (11) aryl-C<sub>1-10</sub> alkyl, and
- (12) heteroaryl-C<sub>1-10</sub> alkyl, or

5  $R^e$  and  $R^f$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

each  $R^g$  is independently selected from:

- (1) C<sub>1-10</sub>alkyl,
- (2) C<sub>3-8</sub>cycloalkyl,
- 10 (3) cycloheteroalkyl,
- (4) aryl,
- (5) arylC<sub>1-4</sub>alkyl,
- (6) heteroaryl,
- (7) heteroarylC<sub>1-4</sub>alkyl,
- 15 (8) -S(O)<sub>m</sub> $R^e$ ,
- (9) -C(O) $R^e$ ,
- (10) -CO<sub>2</sub> $R^e$ ,
- (11) -CO<sub>2</sub>(C $R^eR^f$ )<sub>n</sub>CON $R^eR^f$ , and
- (12) -C(O)N $R^eR^f$ ;

20 each  $R^h$  is independently selected from:

- (1) halogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>3-8</sub>cycloalkyl,
- (4) cycloheteroalkyl,
- 25 (5) aryl,
- (6) arylC<sub>1-4</sub>alkyl,
- (7) heteroaryl,
- (8) heteroarylC<sub>1-4</sub>alkyl,
- (9) -O $R^e$ ,
- 30 (10) -N $R^e$ S(O)<sub>m</sub> $R^f$ ,
- (11) -S(O)<sub>m</sub> $R^e$ ,
- (12) -S $R^e$ ,
- (13) -S(O)<sub>2</sub>O $R^e$ ,
- (14) -S(O)<sub>m</sub>N $R^eR^f$ ,
- 35 (15) -N $R^eR^f$ ,

- (16)  $-\text{O}(\text{CR}^{\text{e}}\text{R}^{\text{f}})_n\text{NR}^{\text{e}}\text{R}^{\text{f}}$ ,  
 (17)  $-\text{C}(\text{O})\text{R}^{\text{e}}$ ,  
 (18)  $-\text{CO}_2\text{R}^{\text{e}}$ ,  
 (19)  $-\text{CO}_2(\text{CR}^{\text{e}}\text{R}^{\text{f}})_n\text{CONR}^{\text{e}}\text{R}^{\text{f}}$ ,  
 5 (20)  $-\text{OC}(\text{O})\text{R}^{\text{e}}$ ,  
 (21)  $-\text{CN}$ ,  
 (22)  $-\text{C}(\text{O})\text{NR}^{\text{e}}\text{R}^{\text{f}}$ ,  
 (23)  $-\text{NR}^{\text{e}}\text{C}(\text{O})\text{R}^{\text{f}}$ ,  
 (24)  $-\text{OC}(\text{O})\text{NR}^{\text{e}}\text{R}^{\text{f}}$ ,  
 10 (25)  $-\text{NR}^{\text{e}}\text{C}(\text{O})\text{OR}^{\text{f}}$ ,  
 (26)  $-\text{NR}^{\text{e}}\text{C}(\text{O})\text{NR}^{\text{e}}\text{R}^{\text{f}}$ ,  
 (27)  $-\text{CF}_3$ , and  
 (28)  $-\text{OCF}_3$ ;

each  $\text{R}^{\text{i}}$  is independently selected from:

- 15 (1)  $\text{C}_{1-10}\text{alkyl}$ ,  
 (2)  $\text{C}_{2-10}\text{alkenyl}$ ,  
 (3)  $\text{C}_{2-10}\text{alkynyl}$ ,  
 (4)  $\text{cycloalkyl}$ ,  
 (5)  $\text{cycloalkyl-C}_{1-10}\text{alkyl}$ ,  
 20 (6)  $\text{cycloheteroalkyl}$ ,  
 (7)  $\text{cycloheteroalkyl-C}_{1-10}\text{alkyl}$ ,  
 (8)  $\text{aryl}$ ,  
 (9)  $\text{heteroaryl}$ ,  
 (10)  $\text{aryl-C}_{1-10}\text{alkyl}$ , and  
 25 (11)  $\text{heteroaryl-C}_{1-10}\text{alkyl}$ ,

each  $\text{R}^{\text{i}}$  may be unsubstituted or substituted with one, two or three substituents selected from  $\text{R}^{\text{h}}$ ;

$m$  is selected from 1 and 2; and

$n$  is selected from 1, 2, and 3;

- 30 provided that when  $\text{R}^{\text{l}}$  and  $\text{R}^{\text{2}}$  are aryl and  $\text{A}$  is cyclobutyl, then the cyclobutyl group is either unsubstituted, or is not substituted with an  $\text{R}^{\text{a}}$  substituent selected from the group consisting of  $-\text{OCH}_3$ ,  $-\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$ , and  $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$  and is not substituted with an  $\text{R}^{\text{h}}$  substituent selected from the group consisting of:  $\text{arylC}_{1-4}\text{alkyl}$ ,  $\text{heteroaryl}$ ,  $-\text{NR}^{\text{c}}\text{S}(\text{O})_m\text{R}^{\text{d}}$ ,  $-\text{S}(\text{O})_m\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,  $-\text{O}(\text{CR}^{\text{e}}\text{R}^{\text{f}})_n\text{NR}^{\text{c}}\text{R}^{\text{d}}$ ,

$-C(O)R^c$ , and  $-C(O)NR^cR^d$ ; and

provided that when **A** is cyclopropyl, then the cyclopropyl group is not bis 2,2-dichlorocyclopropyl; and

- 5 provided that when **A** is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl, then  $R^a$  is not selected from the group consisting of  $-NR^cS(O)_mR^d$ ,  $NR^cC(O)R^d$ ,  $-NR^cC(O)OR^d$ , and  $-NR^cC(O)NR^cR^d$ .

2. The compound according to Claim 1 wherein:

10  $R^1$  is selected from:

- (1)  $C_{1-4}$ alkyl,
- (2)  $C_{3-10}$ cycloalkyl-,
- (3) cycloheteroalkyl,
- (4) phenyl, and
- 15 (5) pyridyl,

wherein each alkyl is optionally substituted with one  $R^a$  substituent, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one to three substituents independently selected from  $R^b$ ;

$R^2$  is selected from:

- 20 (1)  $C_{1-10}$ alkyl,
- (2)  $C_{3-10}$ cycloalkyl,
- (3) cycloheteroalkyl,
- (4) aryl,
- (5) heteroaryl,
- 25 (6)  $-OR^d$ ,
- (7)  $-NR^cR^d$ , and
- (8)  $-CO_2R^d$ , and

wherein each alkyl is optionally substituted with one, two or three substituents independently selected from  $R^a$ , and each cycloalkyl,

- 30 cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one, two or three substituents independently selected from  $R^b$ ;  
and pharmaceutically acceptable salts thereof.

3. The compound according to Claim 2 wherein:

R<sup>3</sup> is selected from:

- (1) hydrogen,
- (2) methyl, and
- (3) ethyl;

5 R<sup>4</sup> is selected from:

- (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) -OR<sup>c</sup>,
- (4) -NR<sup>c</sup>R<sup>d</sup>,
- 10 (5) -NH(CO)OR<sup>d</sup>,
- (6) aryl, and
- (7) heteroaryl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>, and aryl and heteroaryl are optionally substituted

15 with one, two, three or four substituents independently selected from R<sup>b</sup>;

R<sup>6</sup> is selected from:

- (1) hydrogen,
- (2) methyl,
- (3) hydroxyl,
- 20 (4) halogen, and
- (5) -CN;

wherein methyl is optionally substituted with one to three R<sup>a</sup> substituents;

each R<sup>a</sup> is independently selected from:

- (1) -OR<sup>c</sup>
- 25 (2) halogen,
- (3) -S(O)<sub>m</sub>R<sup>c</sup>,
- (4) -SR<sup>c</sup>,
- (5) -S(O)<sub>2</sub>OR<sup>c</sup>,
- (6) -S(O)<sub>m</sub>NR<sup>c</sup>R<sup>d</sup>,
- 30 (7) -NR<sup>c</sup>R<sup>d</sup>,
- (8) -C(O)R<sup>c</sup>,
- (9) -CO<sub>2</sub>R<sup>i</sup>,
- (10) -CN,
- (11) -OC(O)NR<sup>c</sup>R<sup>d</sup>,

- (12) CF<sub>3</sub>,  
 (13) -OCF<sub>3</sub>,  
 (14) C<sub>3-8</sub>cycloalkyl,  
 (15) cycloheteroalkyl, and  
 5 (16) oxo;

each R<sup>b</sup> is independently selected from:

- (1) R<sup>a</sup>,  
 (2) C<sub>1-10</sub>alkyl,  
 (3) aryl, and  
 10 (4) heteroaryl;

each R<sup>c</sup> and R<sup>d</sup> is independently selected from:

- (1) hydrogen,  
 (2) C<sub>2-10</sub>alkyl,  
 (3) C<sub>2-10</sub>alkenyl,  
 15 (4) C<sub>2-10</sub>alkynyl,  
 (5) cycloalkyl,  
 (6) cycloalkyl-C<sub>1-10</sub>alkyl,  
 (7) cycloheteroalkyl,  
 (8) cycloheteroalkyl-C<sub>1-10</sub>alkyl,  
 20 (9) aryl,  
 (10) heteroaryl,  
 (11) aryl-C<sub>1-10</sub>alkyl, and  
 (12) heteroaryl-C<sub>1-10</sub>alkyl, or

R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic  
 25 ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected  
 from oxygen, sulfur and N-R<sup>g</sup>,

each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three  
 substituents selected from R<sup>h</sup>;

each R<sup>h</sup> is independently selected from: halogen,

- 30 (2) C<sub>1-10</sub>alkyl,  
 (3) C<sub>3-8</sub>cycloalkyl,  
 (4) cycloheteroalkyl,  
 (5) aryl,  
 (6) heteroaryl-C<sub>1-4</sub>alkyl,

- (7) -OR<sup>e</sup>,
  - (8) -S(O)<sub>m</sub>Re,
  - (9) -SRe,
  - (10) -S(O)<sub>2</sub>OR<sup>e</sup>,
  - 5 (11) -NR<sup>e</sup>R<sup>f</sup>,
  - (12) -CO<sub>2</sub>Re,
  - (13) -CO<sub>2</sub>(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>CONR<sup>e</sup>R<sup>f</sup>,
  - (14) -OC(O)Re,
  - (15) -CN,
  - 10 (16) -NR<sup>e</sup>C(O)R<sup>f</sup>,
  - (17) -OC(O)NR<sup>e</sup>R<sup>f</sup>,
  - (18) -NR<sup>e</sup>C(O)OR<sup>f</sup>,
  - (19) -NR<sup>e</sup>C(O)NR<sup>e</sup>R<sup>f</sup>,
  - (20) -CF<sub>3</sub>, and
  - 15 (21) -OCF<sub>3</sub>;
- or a pharmaceutically acceptable salt thereof.

4. The compound according to Claim 3 wherein:

R<sup>1</sup> is selected from:

- 20 (1) isopropyl,
- (2) isobutyl,
- (3) n-propyl,
- (4) cyclopropyl,
- (5) cyclobutyl,
- 25 (6) cyclopentyl,
- (7) cyclohexyl,
- (8) piperidinyl,
- (9) phenyl, and
- (10) pyridyl,

30 wherein each alkyl is optionally substituted with one R<sup>a</sup> substituent, and each cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is optionally substituted with one to three substituents independently selected from R<sup>b</sup>;

- (1) cyclobutyl,
- (2) cyclopentyl,

- (3) cyclohexyl,  
(4) pyrrolidinyl,  
(5) pyrimidinyl,  
(6) benzoxazolyl,  
5 (7) dihydroindolyl,  
(8) dihydroquinolinyl,  
(9) benzotriazolyl,  
(10) thiophenyl,  
(11) indolyl,  
10 (12) indazolyl,  
(13) pyrrolidinyl,  
(14) pyridazinyl  
(15) triazolyl,  
(16) azaindolyl,  
15 (17) cyclobutylmethoxy,  
(18) phenyl,  
(19) pyridyl,  
(20) -NR<sup>c</sup>R<sup>d</sup>, and  
(21) -CO<sub>2</sub>R<sup>d</sup>,
- 20 wherein each alkyl is optionally substituted with one or two R<sup>a</sup> substituents and each phenyl or pyridyl is independently with one to three R<sup>b</sup> substituents.  
R<sup>3</sup> is methyl;  
R<sup>4</sup> is selected from:
- (1) hydrogen,  
25 (2) methyl,  
(3) ethyl,  
(4) propyl,  
(5) butyl,  
(6) pentyl,  
30 (7) isopropyl,  
(8) -OH,  
(9) -O-aryl,  
(10) -NH<sub>2</sub>,  
(11) -NH-aryl,  
35 (12) -NH(CO)OR<sup>d</sup>,

(13) phenyl, and

(14) pyridyl,

wherein methyl is optionally substituted with one, two or three substituents independently selected from  $R^a$ , and phenyl, aryl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from  $R^b$ ;

$R^6$  is selected from:

(1) hydrogen,

(2) methyl,

10 (3) hydroxyl,

(4) halogen, and

(5)  $-CN$ ;

each  $R^a$  is independently selected from:

(1)  $-OR^c$ ,

15 (2) halogen,

(3)  $-S(O)_mR^c$ ,

(4)  $-C(O)R^c$ , and

(5)  $-CO_2R^i$ ;

each  $R^b$  is independently selected from:

20 (1)  $R^a$ ,

(2) methyl,

(3) ethyl,

(4) propyl,

(5) phenyl,

25 (6) benzyl, and

(7) pyridyl;

each  $R^c$  and  $R^d$  is independently selected from:

(1) hydrogen,

(2)  $C_{2-10}$ alkyl,

30 (3) aryl, and

(4) heteroaryl, or

$R^c$  and  $R^d$  together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N- $R_g$ ,



each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three substituents selected from R<sup>h</sup>;

each R<sup>h</sup> is independently selected from: halogen,

- (2) C<sub>1-10</sub>alkyl,
- 5 (3) C<sub>3-8</sub>cycloalkyl,
- (4) cycloheteroalkyl,
- (5) aryl,
- (6) heteroarylC<sub>1-4</sub>alkyl,
- (7) -CN,
- 10 (8) -CF<sub>3</sub>, and
- (9) -OCF<sub>3</sub>;

or a pharmaceutically acceptable salt thereof.

5. The compound according to Claim 4 wherein:

15 R<sup>1</sup> and R<sup>2</sup> are independently selected from:

- (1) phenyl,
- (2) 4-fluorophenyl,
- (3) 2-chlorophenyl,
- (4) 3-chlorophenyl,
- 20 (5) 4-chlorophenyl,
- (6) 3-cyanophenyl,
- (7) 4-cyanophenyl,
- (8) 4-methylphenyl,
- (9) 4-isopropylphenyl,
- 25 (10) 4-biphenyl,
- (11) 4-bromophenyl,
- (12) 4-iodophenyl,
- (13) 2,4-dichlorophenyl, and
- (14) 2-chloro-4-fluorophenyl;

30 R<sup>4</sup> is selected from:

- (1) hydrogen,
- (2) methyl,
- (3) ethyl,
- (4) propyl,

- (5) butyl,  
(6) pentyl,  
(7) isopropyl,  
(8) -OH,  
5 (9) -O-phenyl,  
(10) -NH<sub>2</sub>,  
(11) -NH-phenyl,  
(12) -NH(CO)O-alkyl,  
(13) phenyl, and  
10 (14) pyridyl,

wherein methyl and alkyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>, and phenyl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

15 each R<sup>a</sup> is independently selected from:

- (1) -O-aryl,  
(2) -OH,  
(3) bromo,  
(4) chloro,  
20 (5) -S(O)<sub>m</sub>-aryl,  
(6) -C(O)-aryl, and  
(7) -CO<sub>2</sub>-alkyl;

each R<sup>c</sup> and R<sup>d</sup> is independently selected from:

- (1) hydrogen,  
25 (2) ethyl,  
(3) propyl,  
(4) butyl, and  
(5) phenyl, or

R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic  
30 ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sup>g</sup>,

each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three substituents selected from R<sup>h</sup>;

or a pharmaceutically acceptable salt thereof.

35

6. The compound according to Claim 1 wherein **A** is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, hexahydroazepiny, morpholiny, tetrahydrofurany, tetrahydropyrany, and tetrahydrothieny, each  
5 optionally substituted with one, two, or three groups independently selected from R<sup>b</sup>.

7. The compound according to Claim 5 wherein **A** is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofurany, and  
10 tetrahydropyrany, each optionally substituted with one or two groups independently selected from oxo and R<sup>b</sup>.

8. A compound selected from:
- 15 (1) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-4-carboxamide;
  - (2) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(2-pyridyl)-piperidine-3-carboxamide;
  - (3) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-methylmorpholine-2-carboxamide;
  - 20 (4) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(t-butyloxycarbonyl)-pyrrolidine-2(S)-carboxamide;
  - (5) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-2-carboxamide;
  - (6) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2,2-dimethyl-  
25 tetrahydropyran-4-carboxamide;
  - (7) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclopentane-carboxamide;
  - (8) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
  - 30 (9) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclohexane-carboxamide;
  - (10) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;
  - 35 (11) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-cyclopentanone-3-carboxamide;

- (12) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;
- (13) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- 5 (14) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(*t*-butyloxycarbonyl)-2-methylazetidine-2-carboxamide;
- (15) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzyl-piperidine-2-carboxamide;
- (16) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(*t*-butyloxycarbonyl)-pyrrolidine-3-carboxamide;
- 10 (17) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide;
- (18) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-pyrrolidine-2(S)-carboxamide;
- 15 (19) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-2-methylpyrrolidine-2(S)-carboxamide;
- (20) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-methyltetrahydrofuran-2-carboxamide;
- (21) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-tetrahydrofuran-2-carboxamide;
- 20 (22) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide;
- (23) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-oxopyrrolidine-5-carboxamide;
- 25 (24) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-5-methyl-2-oxopyrrolidine-5-carboxamide;
- (25) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzylpyrrolidine-2-carboxamide;
- (26) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(phenylamino)-cyclopentanecarboxamide;
- 30 (27) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (28) N-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopropanecarboxamide;

- (29) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopropanecarboxamide;
- (30) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-chlorophenyl)cyclohexanecarboxamide;
- 5 (31) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-2-phenyltetrahydrofuran-2-carboxamide;
- (32) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3,5-difluorophenyl)cyclopentanecarboxamide;
- (33) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(4-pyridyl)cyclopentanecarboxamide;
- 10 (34) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (35) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- 15 (36) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl-*N*-oxide)cyclopentanecarboxamide;
- (37) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (38) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- 20 (39) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (40) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide;
- 25 (41) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclohexane-carboxamide;
- (42) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclopentane-carboxamide;
- (43) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclohexanecarboxamide;
- 30 (44) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclopentanecarboxamide;
- (45) *N*-[2-(3-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;

- (46) *N*-[2-(3-fluoropyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide
- (47) *N*-[2-(2-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- 5 (48) *N*-[2-(4-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide; and
- (49) *N*-[3-(5-chloro-2-pyridyl)-2-phenyl-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (50) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- 10 (51) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide
- (52) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- (53) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- 15 (54) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- (55) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- 20 (56) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- (57) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- 25 (58) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- (59) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (60) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- 30 (61) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (62) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;

- (63) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- (64) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- 5 (65) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- (66) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- (67) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- 10 (68) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-cyclobutanecarboxamide;
- (69) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylcyclobutanecarboxamide;
- 15 (70) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-ethylcyclobutanecarboxamide;
- (71) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-propylcyclobutanecarboxamide;
- (72) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-benzylcyclobutanecarboxamide;
- 20 (73) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-isopropylcyclobutanecarboxamide;
- (74) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-3-azetidine-3-carboxamide;
- 25 (75) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2(*S*)-azetidinecarboxamide;
- (76) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 1);
- (77) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-butoxycarbonyl-2-ethylazetidine-2-carboxamide (isomer 2);
- 30 (78) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (79) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;

- (80) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- (81) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- 5 (82) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- (83) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- (84) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- 10 (85) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- (86) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- 15 (87) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- (88) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- (89) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- 20 (90) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (91) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- 25 (92) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- (93) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (94) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- 30 (95) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- (96) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;



- (97) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- (98) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- 5 (99) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- (100) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- (101) *N*-(2*S*,3*S*)-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylsulfonylcyclobutanecarboxamide;
- 10 (102) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-3-ethylazetidine-3-carboxamide hydrochloride;
- (103) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 1);
- 15 (104) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-ethylazetidinecarboxamide hydrochloride (isomer 2);
- or a pharmaceutically acceptable salt thereof.


9. The compound according to Claim 8 selected from:

- 20 (1) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (2) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclobutanecarboxamide;
- (3) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- 25 (4) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- (5) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- 30 (6) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- (7) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- 35 (8) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;

- (9) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- (10) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- 5 (11) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (12) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (13) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- 10 (14) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- (15) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- 15 (16) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (17) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (18) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- 20 (19) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-fluorophenyl)cyclopentanecarboxamide;
- (20) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- 25 (21) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(3-fluorophenyl)cyclopentanecarboxamide;
- (22) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- (23) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-fluorophenyl)cyclopentanecarboxamide;
- 30 (24) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;
- (25) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-6-fluorophenyl)cyclopentanecarboxamide;

- (26) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- (27) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-chlorophenyl)cyclopentanecarboxamide;
- 5 (28) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (29) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methylphenyl)cyclopentanecarboxamide;
- (30) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(4-methoxyphenyl)cyclopentanecarboxamide;
- 10 (31) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- (32) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- 15 (33) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- (34) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-phenylcyclohexanecarboxamide;
- (35) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- 20 (36) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2-chloro-4-fluorophenyl)cyclopentanecarboxamide;
- (37) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- 25 (38) *N*-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-(2,4-dichlorophenyl)cyclopentanecarboxamide;
- (39) *N*-(2*S*,3*S*)-[3-(4-Chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-1-methylsulfonylcyclobutanecarboxamide;
- 30 (40) *N*-(2*S*,3*S*)-[3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-3-ethylazetidine-3-carboxamide hydrochloride;
- or a pharmaceutically acceptable salt thereof.

10. A compound of structural formula:



or a pharmaceutically acceptable salt thereof, wherein:

R<sup>1</sup> is selected from:

- 5           (1) aryl, and  
              (2) pyridyl,

wherein aryl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

R<sup>2</sup> is selected from:

- 10           (1) aryl, and  
              (2) pyridyl,

wherein aryl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

R<sup>3</sup> is selected from:

- 15           (1) hydrogen, and  
              (2) C<sub>1-4</sub>alkyl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from R<sup>a</sup>;

R<sup>4</sup> is selected from:

- 20           (1) hydrogen,  
              (2) C<sub>1-4</sub>alkyl,  
              (3) C<sub>2-4</sub>alkenyl,  
              (4) C<sub>2-4</sub>alkynyl,  
              (5) -OR<sup>c</sup>,  
25           (6) CO<sub>2</sub>R<sup>c</sup>,  
              (7) -OCOOR<sup>c</sup>,  
              (8) -OCONR<sup>c</sup>R<sup>d</sup>,  
              (9) -NR<sup>c</sup>R<sup>d</sup>,  
              (10) -NR<sup>c</sup>(CO)OR<sup>d</sup>,  
30           (11) -NR<sup>c</sup>SO<sub>2</sub>R<sup>d</sup>,  
              (12) -S(O)<sub>m</sub>R<sup>c</sup>,

(13) aryl, and

(14) heteroaryl,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one, two, three or four substituents independently selected from  $R^a$ , and aryl and heteroaryl are

5 optionally substituted with one, two, three or four substituents independently selected from  $R^b$ ;

**A** is a 3- to 8-membered monocyclic saturated ring incorporating the same carbon atom to which  $R^4$  is attached, and optionally containing one to two heteroatoms  
 10 chosen from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted with one, two, three or four substituents independently selected from oxo and  $R^b$ ;

each  $R^a$  is independently selected from:

- (1)  $-OR^c$ ,
- 15 (2)  $-NR^cS(O)_mR^d$ ,
- (3)  $-NO_2$ ,
- (4) halogen,
- (5)  $-S(O)_mR^c$ ,
- (6)  $-SR^c$ ,
- 20 (7)  $-S(O)_2OR^c$ ,
- (8)  $-S(O)_mNR^cR^d$ ,
- (9)  $-NR^cR^d$ ,
- (10)  $-O(CR^eR^f)_nNR^cR^d$ ,
- (11)  $-C(O)R^c$ ,
- 25 (12)  $-CO_2R^i$ ,
- (13)  $-CO_2(CR^eR^f)_nCONR^cR^d$ ,
- (14)  $-OC(O)R^c$ ,
- (15)  $-CN$ ,
- (16)  $-C(O)NR^cR^d$ ,
- 30 (17)  $-NR^cC(O)R^d$ ,
- (18)  $-OC(O)NR^cR^d$ ,
- (19)  $-NR^cC(O)OR^d$ ,
- (20)  $-NR^cC(O)NR^cR^d$ ,
- (21)  $-CR^c(N-OR^d)$ ,
- 35 (22)  $CF_3$ ,

- (23) -OCF<sub>3</sub>,
- (24) C<sub>3-8</sub>cycloalkyl,
- (25) cycloheteroalkyl, and
- (26) oxo;

5 each R<sup>b</sup> is independently selected from:

- (1) R<sup>a</sup>,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>3-8</sub>cycloalkyl,
- (4) cycloheteroalkyl,
- 10 (5) aryl,
- (6) arylC<sub>1-4</sub>alkyl,
- (7) heteroaryl, and
- (8) heteroarylC<sub>1-4</sub>alkyl;

each R<sup>c</sup> and R<sup>d</sup> are independently selected from:

- 15 (1) hydrogen,
- (2) C<sub>1-10</sub>alkyl,
- (3) C<sub>2-10</sub>alkenyl,
- (4) C<sub>2-10</sub>alkynyl,
- (5) cycloalkyl,
- 20 (6) cycloalkyl-C<sub>1-10</sub>alkyl,
- (7) cycloheteroalkyl,
- (8) cycloheteroalkyl-C<sub>1-10</sub>alkyl,
- (9) aryl,
- (10) heteroaryl,
- 25 (11) aryl-C<sub>1-10</sub>alkyl, and
- (12) heteroaryl-C<sub>1-10</sub>alkyl, or

R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sup>g</sup>,

30 each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three substituents selected from R<sup>h</sup>;

each R<sup>e</sup> and R<sup>f</sup> are independently selected from:

- (1) hydrogen,

- 5 (2) C<sub>1-10</sub>alkyl,  
 (3) C<sub>2-10</sub>alkenyl,  
 (4) C<sub>2-10</sub>alkynyl;  
 (5) cycloalkyl,  
 (6) cycloalkyl-C<sub>1-10</sub> alkyl,  
 (7) cycloheteroalkyl,  
 (8) cycloheteroalkyl-C<sub>1-10</sub> alkyl,  
 (9) aryl,  
 (10) heteroaryl,  
 10 (11) aryl-C<sub>1-10</sub> alkyl, and  
 (12) heteroaryl-C<sub>1-10</sub> alkyl, or

R<sup>e</sup> and R<sup>f</sup> together with the carbon to which they are attached form a ring of 5 to 7  
 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and  
 nitrogen;

15

each R<sup>g</sup> is independently selected from:

- (1) C<sub>1-10</sub>alkyl,  
 (2) C<sub>3-8</sub>cycloalkyl,  
 (3) cycloheteroalkyl,  
 20 (4) aryl,  
 (5) arylC<sub>1-4</sub>alkyl,  
 (6) heteroaryl,  
 (7) heteroarylC<sub>1-4</sub>alkyl,  
 (8) -S(O)<sub>m</sub>R<sup>e</sup>,  
 25 (9) -C(O)R<sup>e</sup>,  
 (10) -CO<sub>2</sub>R<sup>e</sup>,  
 (11) -CO<sub>2</sub>(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>CONR<sup>e</sup>R<sup>f</sup>, and  
 (12) -C(O)NR<sup>e</sup>R<sup>f</sup>;

30 each R<sup>h</sup> is independently selected from:

- (1) halogen,  
 (2) C<sub>1-10</sub>alkyl,  
 (3) C<sub>3-8</sub>cycloalkyl,  
 (4) cycloheteroalkyl,

- (5) aryl,  
 (6) arylC<sub>1-4</sub>alkyl,  
 (7) heteroaryl,  
 (8) heteroarylC<sub>1-4</sub>alkyl,  
 5 (9) -OR<sup>e</sup>,  
 (10) -NR<sup>e</sup>S(O)<sub>m</sub>R<sup>f</sup>,  
 (11) -S(O)<sub>m</sub>R<sup>e</sup>,  
 (12) -SR<sup>e</sup>,  
 (13) -S(O)<sub>2</sub>OR<sup>e</sup>,  
 10 (14) -S(O)<sub>m</sub>NR<sup>e</sup>R<sup>f</sup>,  
 (15) -NR<sup>e</sup>R<sup>f</sup>,  
 (16) -O(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>NR<sup>e</sup>R<sup>f</sup>,  
 (17) -C(O)R<sup>e</sup>,  
 (18) -CO<sub>2</sub>R<sup>e</sup>,  
 15 (19) -CO<sub>2</sub>(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>CONR<sup>e</sup>R<sup>f</sup>,  
 (20) -OC(O)R<sup>e</sup>,  
 (21) -CN,  
 (22) -C(O)NR<sup>e</sup>R<sup>f</sup>,  
 (23) -NR<sup>e</sup>C(O)R<sup>f</sup>,  
 20 (24) -OC(O)NR<sup>e</sup>R<sup>f</sup>,  
 (25) -NR<sup>e</sup>C(O)OR<sup>f</sup>,  
 (26) -NR<sup>e</sup>C(O)NR<sup>e</sup>R<sup>f</sup>,  
 (27) -CF<sub>3</sub>, and  
 (28) -OCF<sub>3</sub>;  
 25 each R<sup>i</sup> is independently selected from:  
 (1) C<sub>1-10</sub>alkyl,  
 (2) C<sub>2-10</sub> alkenyl,  
 (3) C<sub>2-10</sub>alkynyl,  
 (4) cycloalkyl,  
 30 (5) cycloalkyl-C<sub>1-10</sub>alkyl,  
 (6) cycloheteroalkyl,  
 (7) cycloheteroalkyl-C<sub>1-10</sub> alkyl,  
 (8) aryl,  
 (9) heteroaryl,  
 35 (10) aryl-C<sub>1-10</sub>alkyl, and



(11) heteroaryl-C<sub>1-10</sub>alkyl,

each R<sup>i</sup> may be unsubstituted or substituted with one, two or three substituents selected from R<sup>h</sup>;

m is selected from 1 and 2; and

5 n is selected from 1, 2, and 3;

provided that when R<sup>1</sup> and R<sup>2</sup> are aryl and **A** is cyclobutyl, the cyclobutyl group is either unsubstituted, or is not substituted with an R<sup>a</sup> substituent selected from the group consisting of -OCH<sub>3</sub>, -C(O)NR<sup>c</sup>R<sup>d</sup>, and -NR<sup>c</sup>C(O)NR<sup>c</sup>R<sup>d</sup> and is not

10 substituted with an R<sup>h</sup> substituent selected from the group consisting of: arylC<sub>1-4</sub>alkyl, heteroaryl, -NR<sup>c</sup>S(O)<sub>m</sub>R<sup>d</sup>, -S(O)<sub>m</sub>NR<sup>c</sup>R<sup>d</sup>, -O(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>NR<sup>c</sup>R<sup>d</sup>, -C(O)R<sup>c</sup>, and -C(O)NR<sup>c</sup>R<sup>d</sup>; and

provided that when **A** is cyclopropyl, the cyclopropyl group is not bis 2,2-dichlorocyclopropyl; and

15 provided that when **A** is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl, R<sup>a</sup> is not selected from the group consisting of -NR<sup>c</sup>S(O)<sub>m</sub>R<sup>d</sup>, NR<sup>c</sup>C(O)R<sup>d</sup>, -NR<sup>c</sup>C(O)OR<sup>d</sup>, and -NR<sup>c</sup>C(O)NR<sup>c</sup>R<sup>d</sup>.

11. The compound of claim 10 wherein:

20 R<sup>1</sup> and R<sup>2</sup> are phenyl, each optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;  
and pharmaceutically acceptable salts thereof.

12. The compound of claim 11 wherein:

25 R<sup>3</sup> is selected from:

- (1) hydrogen,
- (2) methyl, and
- (3) ethyl;

R<sup>4</sup> is selected from:

- 30
- (1) hydrogen,
  - (2) C<sub>1-4</sub>alkyl,
  - (3) -OR<sup>c</sup>,
  - (4) -NR<sup>c</sup>R<sup>d</sup>,
  - (5) -NH(CO)OR<sup>d</sup>,

(6) aryl, and

(7) heteroaryl,

wherein alkyl is optionally substituted with one, two, three or four substituents independently selected from  $R^a$ , and aryl and heteroaryl are optionally substituted

5 with one, two, three or four substituents independently selected from  $R^b$ ;

each  $R^a$  is independently selected from:

- (1)  $-OR^c$
- (2) halogen,
- (3)  $-S(O)_mR^c$ ,
- 10 (4)  $-SR^c$ ,
- (5)  $-S(O)_2OR^c$ ,
- (6)  $-S(O)_mNR^cR^d$ ,
- (7)  $-NR^cR^d$ ,
- (8)  $-C(O)R^c$ ,
- 15 (9)  $-CO_2R^i$ ,
- (10)  $-CN$ ,
- (11)  $-OC(O)NR^cR^d$ ,
- (12)  $CF_3$ ,
- (13)  $-OCF_3$ ,
- 20 (14)  $C_3$ -8cycloalkyl,
- (15) cycloheteroalkyl, and
- (16) oxo;

each  $R^b$  is independently selected from:

- (1)  $R^a$ ,
- 25 (2)  $C_{1-10}$ alkyl,
- (3) aryl, and
- (4) heteroaryl;

each  $R^c$  and  $R^d$  is independently selected from:

- (1) hydrogen,
- 30 (2)  $C_{2-10}$ alkyl,
- (3)  $C_{2-10}$ alkenyl,
- (4)  $C_{2-10}$ alkynyl,
- (5) cycloalkyl,
- (6) cycloalkyl- $C_{1-10}$ alkyl,
- 35 (7) cycloheteroalkyl,

- (8) cycloheteroalkyl-C<sub>1-10</sub> alkyl,  
 (9) aryl,  
 (10) heteroaryl,  
 (11) aryl-C<sub>1-10</sub>alkyl, and  
 5 (12) heteroaryl-C<sub>1-10</sub>alkyl, or
- R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sup>g</sup>,  
 each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three  
 10 substituents selected from R<sup>h</sup>;  
 each R<sup>h</sup> is independently selected from: halogen,  
 (2) C<sub>1-10</sub>alkyl,  
 (3) C<sub>3-8</sub>cycloalkyl,  
 (4) cycloheteroalkyl,  
 15 (5) aryl,  
 (6) heteroarylC<sub>1-4</sub>alkyl,  
 (7) -OR<sup>e</sup>,  
 (8) -S(O)<sub>m</sub>R<sup>e</sup>,  
 (9) -SR<sup>e</sup>,  
 20 (10) -S(O)<sub>2</sub>OR<sup>e</sup>,  
 (11) -NR<sup>e</sup>R<sup>f</sup>,  
 (12) -CO<sub>2</sub>R<sup>e</sup>,  
 (13) -CO<sub>2</sub>(CR<sup>e</sup>R<sup>f</sup>)<sub>n</sub>CONR<sup>e</sup>R<sup>f</sup>,  
 (14) -OC(O)R<sup>e</sup>,  
 25 (15) -CN,  
 (16) -NR<sup>e</sup>C(O)R<sup>f</sup>,  
 (17) -OC(O)NR<sup>e</sup>R<sup>f</sup>,  
 (18) -NR<sup>e</sup>C(O)OR<sup>f</sup>,  
 (19) -NR<sup>e</sup>C(O)NR<sup>e</sup>R<sup>f</sup>,  
 30 (20) -CF<sub>3</sub>, and  
 (21) -OCF<sub>3</sub>;
- or a pharmaceutically acceptable salt thereof.

13. The compound of claim 12 wherein:

R<sup>1</sup> and R<sup>2</sup> are independently selected from:

- (1) phenyl,
- (2) 4-fluorophenyl,
- (3) 2-chlorophenyl,
- 5 (4) 3-chlorophenyl,
- (5) 4-chlorophenyl,
- (6) 4-cyanophenyl,
- (7) 4-methylphenyl,
- (8) 4-isopropylphenyl,
- 10 (9) 4-biphenyl,
- (10) 4-bromophenyl,
- (11) 4-iodophenyl,
- (12) 2,4-dichlorophenyl, and
- (13) 2-chloro-4-fluorophenyl;

15 R<sup>3</sup> is methyl;

R<sup>4</sup> is selected from:

- (1) hydrogen,
- (2) methyl,
- (3) -OH,
- 20 (4) -O-aryl,
- (5) -NH<sub>2</sub>,
- (6) -NH-aryl,
- (7) -NH(CO)OR<sup>d</sup>,
- (8) phenyl, and
- 25 (9) pyridyl,

wherein methyl is optionally substituted with one, two or three substituents independently selected from R<sup>a</sup>, and phenyl, aryl and pyridyl are optionally substituted with one, two, three or four substituents independently selected from R<sup>b</sup>;

30 each R<sup>a</sup> is independently selected from:

- (1) -OR<sup>c</sup>,
- (2) halogen,
- (3) -S(O)<sub>m</sub>R<sup>c</sup>,
- (4) -C(O)R<sup>c</sup>, and

(5)  $-\text{CO}_2\text{R}^i$ ;

each  $\text{R}^b$  is independently selected from:

- 5       (1)  $\text{R}^a$ ,  
      (2) methyl,  
      (3) ethyl,  
      (4) propyl,  
      (5) phenyl,  
      (6) benzyl, and  
      (7) pyridyl;

10    each  $\text{R}^c$  and  $\text{R}^d$  is independently selected from:

- (1) hydrogen,  
      (2)  $\text{C}_2$ -10alkyl,  
      (3) aryl, and  
      (4) heteroaryl, or

15     $\text{R}^c$  and  $\text{R}^d$  together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N- $\text{R}^g$ ,

each  $\text{R}^c$  and  $\text{R}^d$  may be unsubstituted or substituted with one, two or three substituents selected from  $\text{R}^h$ ;

20    each  $\text{R}^h$  is independently selected from: halogen,

- (2)  $\text{C}_1$ -10alkyl,  
      (3)  $\text{C}_3$ -8cycloalkyl,  
      (4) cycloheteroalkyl,  
      (5) aryl,  
25    (6) heteroaryl $\text{C}_1$ -4alkyl,  
      (7)  $-\text{CN}$ ,  
      (8)  $-\text{CF}_3$ , and  
      (9)  $-\text{OCF}_3$ ;

or a pharmaceutically acceptable salt thereof.

30

14. The compound of Claim 13 wherein:

$\text{R}^1$  and  $\text{R}^2$  are independently selected from phenyl and 4-chlorophenyl;

$\text{R}^4$  is selected from:

- (1) hydrogen,

- (2) methyl,  
(3) -OH,  
(4) -O-phenyl,  
(5) -NH<sub>2</sub>,  
5 (6) -NH-phenyl,  
(7) -NH(CO)O-alkyl,  
(8) phenyl, and  
(9) pyridyl,

wherein methyl and alkyl are optionally substituted with one, two, three or  
10 four substituents independently selected from R<sup>a</sup>, and phenyl and pyridyl are  
optionally substituted with one, two, three or four substituents independently  
selected from R<sup>b</sup>;

each R<sup>a</sup> is independently selected from:

- (1) -O-aryl,  
15 (2) -OH,  
(3) bromo,  
(4) chloro,  
(5) -S(O)<sub>m</sub>-aryl,  
(6) -C(O)-aryl, and  
20 (7) -CO<sub>2</sub>-alkyl;

each R<sup>c</sup> and R<sup>d</sup> is independently selected from:

- (1) hydrogen,  
(2) ethyl,  
(3) propyl,  
25 (4) butyl, and  
(5) phenyl, or

R<sup>c</sup> and R<sup>d</sup> together with the atom(s) to which they are attached form a heterocyclic  
ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected  
from oxygen, sulfur and N-R<sup>g</sup>,

- 30 each R<sup>c</sup> and R<sup>d</sup> may be unsubstituted or substituted with one, two or three  
substituents selected from R<sup>h</sup>;  
or a pharmaceutically acceptable salt thereof.

15. The compound of Claim 10 wherein **A** is selected from the  
35 group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl,

azetidiny, pyrrolidiny, piperidiny, piperaziny, hexahydroazepiny, morpholiny, tetrahydrofurany, tetrahydropyrany, and tetrahydrothieny, each optionally substituted with one, two, or three groups independently selected from oxo and R<sup>b</sup>.

5                    16.    The compound of Claim 10 wherein **A** is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofurany, and tetrahydropyrany, each optionally substituted with one or two groups independently selected from oxo and R<sup>b</sup>.

10

17.    A compound selected from:

- (1) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-4-carboxamide;
- 15    (2) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(2-pyridyl)-piperidine-3-carboxamide;
- (3) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-methylmorpholine-2-carboxamide;
- (4) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(*t*-butoxycarbonyl)-pyrrolidine-2(*S*)-carboxamide;
- 20    (5) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-methylpiperidine-2-carboxamide;
- (6) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2,2-dimethyl-tetrahydropyran-4-carboxamide;
- (7) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclopentane-carboxamide;
- 25    (8) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- (9) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-3-benzoyl-cyclohexane-carboxamide;
- 30    (10) *cis*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;
- (11) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-cyclopentanone-3-carboxamide;
- 35    (12) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-4-benzoyl-cyclohexane-carboxamide;

- (13) *trans*-N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexane-carboxamide;
- (14) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(*t*-butyloxycarbonyl)-2-methylazetidine-2-carboxamide;
- 5 (15) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzyl-piperidine-2-carboxamide;
- (16) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(*t*-butyloxycarbonyl)-pyrrolidine-3-carboxamide;
- (17) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide;
- 10 (18) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-pyrrolidine-2(S)-carboxamide;
- (19) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-((3,5-dichloro)benzenesulfonyl)-2-methylpyrrolidine-2(S)-carboxamide;
- 15 (20) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-methyltetrahydrofuran-2-carboxamide;
- (21) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-tetrahydrofuran-2-carboxamide;
- (22) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide;
- 20 (23) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-oxopyrrolidine-5-carboxamide;
- (24) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-5-methyl-2-oxopyrrolidine-5-carboxamide;
- 25 (25) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-benzylpyrrolidine-2-carboxamide;
- (26) N-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(phenylamino)-cyclopentanecarboxamide;
- (27) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopentanecarboxamide;
- 30 (28) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-phenylcyclopropanecarboxamide;
- (29) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopropanecarboxamide;



- (30) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-chlorophenyl)cyclohexanecarboxamide;
- (31) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-2-phenyltetrahydrofuran-2-carboxamide;
- 5 (32) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3,5-difluorophenyl)cyclopentanecarboxamide;
- (33) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(4-pyridyl)cyclopentanecarboxamide;
- (34) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- 10 (35) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl)cyclopentanecarboxamide;
- (36) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(3-pyridyl-*N*-oxide)cyclopentanecarboxamide;
- 15 (37) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(2-pyridyl)cyclopentanecarboxamide;
- (38) *N*-[3-(4-chlorophenyl)-2-phenyl-1-methylpropyl]-1-(5-bromo-3-pyridyl)cyclopentanecarboxamide;
- (39) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- 20 (40) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide;
- (41) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclohexanecarboxamide;
- 25 (42) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclopentanecarboxamide;
- (43) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclohexanecarboxamide;
- (44) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-aminocyclopentanecarboxamide;
- 30 (45) *N*-[2-(3-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;
- (46) *N*-[2-(3-fluoropyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide

- (47) *N*-[2-(2-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;  
(48) *N*-[2-(4-pyridyl)-3-(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide; and  
5 (49) *N*-[3-(5-chloro-2-pyridyl)-2-phenyl-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;  
or a pharmaceutically acceptable salt thereof.

18. The compound according to Claim 17 selected from:  
10 (1) *cis*-*N*-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-benzoyl-cyclohexanecarboxamide;  
(2) *N*-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-2-phenylcyclopropane-carboxamide;  
(3) *N*-(2,3-bis(4-chlorophenyl)-1-methylpropyl)-1-(3-pyridyl)-cyclopentane-carboxamide;  
15 (4) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(4-chlorophenyloxy)-cyclohexanecarboxamide;  
(5) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-hydroxycyclohexanecarboxamide;  
(6) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclohexane-carboxamide;  
20 (7) *N*-[2,3-bis(4-chlorophenyl)-1-methylpropyl]-1-(*tert*-butoxycarbonylamino)cyclopentane-carboxamide;  
or a pharmaceutically acceptable salt thereof.

19. A composition comprising a compound according to Claim 1  
25 and a pharmaceutically acceptable carrier.

20. A composition comprising a compound according to Claim 8  
and a pharmaceutically acceptable carrier.

- 30 21. A method of preventing obesity in a person at risk for obesity comprising administration to said person of about 0.001 to about 100 mg/kg of a compound according to Claim 1.

22. A method of preventing obesity in a person at risk for obesity comprising administration to said person of about 0.001 to about 100 mg/kg of a compound according to Claim 8.

5                   23. A method of treating a disease mediated by the Cannabinoid-1 receptor comprising administration of a therapeutically effective amount of a compound of Claim 1 to a patient in need of such treatment.

10                   24. The method according to Claim 23 wherein the disease mediated by the Cannabinoid-1 receptor is selected from: psychosis, memory deficit, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders, cerebral vascular accidents, head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, schizophrenia, substance abuse disorders, constipation, chronic intestinal pseudo-obstruction, cirrhosis of the liver, asthma, obesity, and other eating disorders  
15 associated with excessive food intake.

20                   25. The method according to Claim 24 wherein the disease mediated by the Cannabinoid-1 receptor is an eating disorder associated with excessive food intake.

26. The method according to Claim 25 wherein the eating disorder associated with excessive food intake is selected from obesity, bulimia nervosa, and compulsive eating disorders.

25                   27. The method according to Claim 26 wherein the eating disorder associated with excessive food intake is obesity.

30                   28. The use of a compound according to Claim 1 for the manufacture of a medicament useful for the treatment of a disease mediated by the Cannabinoid-1 receptor in a human patient in need of such treatment.

35                   29. The use according to Claim 28 wherein the disease mediated by the Cannabinoid-1 receptor is selected from: psychosis, memory deficit, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders, cerebral vascular accidents, head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease,

schizophrenia, substance abuse disorders, constipation, chronic intestinal pseudo-obstruction, cirrhosis of the liver, asthma, obesity, and other eating disorders associated with excessive food intake.

5                    30.    The use according to Claim 29 wherein the disease mediated by the Cannabinoid-1 receptor is an eating disorder associated with excessive food intake.

10                   31.    The use according to Claim 30, wherein the eating disorder associated with excessive food intake is selected from obesity, bulimia nervosa, and compulsive eating disorders.

15                   32.    The use according to Claim 31 wherein the eating disorder associated with excessive food intake is obesity.

                    33.    The use of a compound according to Claim 1 for the manufacture of a medicament for the prevention of obesity in a person at risk therefor.

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number  
**WO 2003/082190 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/16**,  
31/335, 31/34, 31/44, 31/445, C07C 233/57, C07D 205/00,  
207/08, 211/30, 211/32, 211/68, 211/80, 307/02, 315/00

(21) International Application Number:  
PCT/US2003/008722

(22) International Filing Date: 21 March 2003 (21.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/367,655 26 March 2002 (26.03.2002) US

(71) Applicant (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HAGMANN, William, K.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **LIN, Linus, S.** [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **SHAH, Shrenik, K.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **GOULET, Mark, T.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **JEWELL, James, P.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
19 February 2004

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **SPIROCYCLIC AMIDES AS CANNABINOID RECEPTOR MODULATORS**

(57) Abstract: Novel compounds of structural formula (I) are antagonists and/or inverse agonists of the Cannabinoid-1 (CB1) receptor and are useful in the treatment, prevention and suppression of diseases mediated by the CB1 receptor. The compounds of the present invention are useful as psychotropic drugs in the treatment of psychosis, memory deficits, cognitive disorders, migraine, neuropathy, neuro-inflammatory disorders including multiple sclerosis and Guillain-Barre syndrome and the inflammatory sequelae of viral encephalitis, cerebral vascular accidents, and head trauma, anxiety disorders, stress, epilepsy, Parkinson's disease, movement disorders, and schizophrenia. The compounds are also useful for the treatment of substance abuse disorders, the treatment of obesity or eating disorders, as well as, the treatment of asthma, constipation, chronic intestinal pseudo-obstruction, and cirrhosis of the liver.

WO 2003/082190 A3

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/08722

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/318, 330, 357, 449, 461, 613; 546/193, 225, 336; 548/567, 953; 549/126, 487

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS—structure

EAST/West—image

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	US 5,462,931 A (SHABER et al.) 31 October 1995, see entire document, especially col. 17-18, table 1, compound 71.	1-2, 6-7 ----- 3
A,P	US 2003/0105132 A1 (CHALLENGER et al.) 05 June 2003, see entire document.	1-33
Y	US 4,150,052 A (WATSON et al.) 17 April 1979, see entire document, especially col. 6, line 19.	1-2, 6-7
Y	Database CAPLUS on STN (Columbus, OH, USA) No. 133:53157, 'Development of a novel class of monocyclic and bicyclic alkyl amides....' Berglund et al. Drug Design and Discovery, vol. 16, pages 281-294 (2000), see RN 278176-97-9 or 278176-98-0.	1-2, 6-7

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 26 SEPTEMBER 2003	Date of mailing of the international search report 09 DEC 2003
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer CELIA CHANG <i>Janice Ford</i> Telephone No. (703) 308-1235 <i>for</i>

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/08722

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

IPC-7 A61K 31/16, 31/335, 31/34, 31/44, 31/445; C07C 233/57; C07D 205/00, 207/08, 211/30, 211/32, 211/68, 211/80, 307/02, 315/00

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/318, 330, 357, 449, 461, 613; 546/193, 225, 336; 548/567, 953; 549/426, 487

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claims 1-17 (in part), 18, 19-23 (in part), drawn to compounds, compositions and method of using material wherein A is C<sub>3-6</sub>cycloalkyl.

Group II, claims 1-17 (in part), 19-23 (in part), drawn to compounds, compositions and method of using material wherein A is one oxygen containing monocyclic saturated ring.

Group III, claims 1-17 (in part), 19-23 (in part), drawn to compounds, compositions and method of using material wherein A is one sulfur containing monocyclic saturated ring.

Group IV, claims 1-17 (in part), 19-23 (in part), drawn to compounds, compositions and method of using material wherein A is one nitrogen containing monocyclic saturated ring.

Group V, claims 1-17 (in part), 19-23 (in part), drawn to compounds, compositions and method of using material wherein A is two heteroatom containing monocyclic saturated ring.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reason:

In PCT administration instruction Annex B, Markush practice, section (f)(i)(B)(2)(v) it was stated that "when dealing with alternatives, if it can be shown that at least one Markush alternative is not novel over the [prior art, the question of unity of invention shall be reconsidered...."

In the instant case, it was found, *at least one alternative is not novel* since at least one prior art anticipated the claims, see Ohkubo et al. CA 123:285788, see RN 169497-82-9 anticipated claim 1 when R1 is aryl, R2 is COORD, Rd is H, R6 is H, R3 is H, R4 is H, A is monocyclic saturated one nitrogen containing ring substituted with C(O)Rc, Rc is cyclohexyl-C<sub>3</sub>alkyl.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US03/08722

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	Database CAPLUS on STN (Columbus, OH, USA) No. 138:24957, 'Amino acid conjugates providing for sustained....' Gallop et al. WO 2002100344, see entire document, especially RN 478186-94-6.	1-2, 6-7